



Correlates and assessment of excess cardiovascular risk in bronchiectasis

Aarash D. Saleh, Bessie Kwok, Jeremy S. Brown and John R. Hurst

Affiliation: Centre for Inflammation and Tissue Repair, UCL Respiratory Medicine, Royal Free Campus, University College London, London, UK.

Correspondence: Aarash Dawood Saleh, Centre for Inflammation and Tissue Repair, UCL Respiratory, Royal Free Campus, University College London, London, NW3 2PF, UK. E-mail: aarash.saleh@nhs.net

 @ERSpublications

Bronchiectasis is associated with elevated cardiovascular risk, which is greatest in frequent exacerbators <http://ow.ly/pFgg30eXEXY>

Cite this article as: Saleh AD, Kwok B, Brown JS, *et al.* Correlates and assessment of excess cardiovascular risk in bronchiectasis. *Eur Respir J* 2017; 50: 1701127 [<https://doi.org/10.1183/13993003.01127-2017>].

ABSTRACT Patients with bronchiectasis are at increased risk of cardiovascular disease. We aimed to identify factors associated with elevated cardiovascular risk in bronchiectasis, measured using aortic stiffness and cardiac biomarkers. In addition, we sought to compare these direct measures against calculated QRISK2 scores.

Aortic stiffness, cardiac biomarkers and systemic inflammation were measured in 101 adults with stable bronchiectasis. In addition, clinical and demographic data were collected to allow calculation of QRISK2 score and the bronchiectasis severity index (BSI) for each patient.

The BSI score correlated with measured cardiovascular risk assessments, partly due to greater exacerbation frequency and lower forced expiratory volume in 1 s. Pulse-wave velocity was significantly higher in frequent exacerbators (≥ 3 events-year⁻¹) than infrequent exacerbators (< 3 events-year⁻¹; 10.5 *versus* 9.2 m·s⁻¹, $p=0.01$). In addition, frequent exacerbators had elevated serum C-reactive protein concentration, suggesting increased systemic inflammation (4.8 *versus* 2.2 mg·L⁻¹, $p=0.005$). QRISK2 systematically underestimated cardiovascular risk in this population (median change in relative risk 1.29). Underestimation was associated with frequent exacerbations and male sex.

Patients with bronchiectasis have greater cardiovascular risk than published reference populations. Excess cardiovascular risk is associated with exacerbation frequency and impaired lung function. Cardiovascular risk assessment in bronchiectasis should be individualised, as calculation tools are likely to underestimate the risk in this population.

Received: June 06 2017 | Accepted after revision: Aug 16 2017

Support statement: This study was supported by National Institute for Health Research funding to the University College London Hospitals Biomedical Research Centre (fast-track grant F180). Funding information for this article has been deposited with the Crossref Funder Registry.

Conflict of interest: None declared.

Copyright ©ERS 2017

Introduction

Bronchiectasis is defined as the persistent abnormal dilatation of bronchi in association with chronic purulent cough and/or frequent chest infections, called exacerbations [1]. This pathological end-point can result from a variety of diseases that affect mucociliary clearance and immunity. Consequent frequent or chronic lung infection further exacerbates mucosal damage and perpetuates the classic vicious cycle [2]. Long neglected, UK mortality from bronchiectasis is currently ~1000 deaths per year and rising [3].

A large cohort analysis has demonstrated the significant burden of comorbidity carried by patients with bronchiectasis and the substantial impact this has on mortality [4]. Of interest, cardiovascular disease is likely to be causally linked to bronchiectasis, as it is to other diseases that feature increased systemic inflammation [5]. We have previously defined a relationship between higher exacerbation frequency in chronic obstructive pulmonary disease (COPD) and elevated cardiovascular risk [6]. In addition, bronchiectasis is associated with increased systemic inflammation [7] and studies have reported that patients with bronchiectasis are at greater cardiac risk than matched controls [8, 9]. In order to reduce this excess cardiovascular risk in bronchiectasis, work is needed to better understand its correlates and how to identify bronchiectasis patients at higher risk of developing cardiovascular disease. The UK National Institute for Health and Care Excellence recommends the use of the QRISK2 cardiovascular risk tool to aid decisions on blood pressure management and use of statins, but recognises that QRISK2 may not adequately assess risk in patients with systemic inflammatory disorders [10]. The QRISK2 algorithm uses epidemiological data, selected comorbidities, cholesterol, blood pressure and body mass index (BMI) to calculate the 10-year risk of myocardial infarction and stroke [11]. For example, rheumatoid arthritis is included in QRISK2 as a comorbidity, but bronchiectasis is not. We hypothesised that use of the QRISK2 score might systematically underestimate cardiovascular risk in patients with bronchiectasis, and therefore restrict the use of effective protective interventions.

The American College of Cardiology and American Heart Association 2013 guideline on the assessment of cardiovascular risk highlights the potential for technology such as cardiac biomarkers to better inform individual risk in patients otherwise classed as “intermediate” [12]. One example is arterial stiffness, an independent predictor of cardiovascular risk in a number of patient groups [13–15]. As well as being correlated with advancing age and atherosclerotic burden, elevated arterial stiffness (measured by pulse-wave velocity (PWV)) has a causal relationship with cardiovascular mortality through increased left ventricular afterload and impaired coronary perfusion [16]. PWV is a gold-standard assessment of arterial stiffness providing additional predictive value beyond estimates based on existing cardiovascular risk factors [17]. In addition, circulating cardiac biomarkers are useful in better defining individual cardiovascular risk. Troponins are markers of myocardial injury, and occult cardiac disease can result in minor elevation of circulating troponins. Troponin concentration predicts future cardiac events [18] and all-cause mortality [19], even when adjusted for standard risk factors. Brain natriuretic peptide (BNP) is another marker of myocardial and circulatory stress that can help predict future cardiac events and death in asymptomatic populations after adjustment for clinical risk factors [20]. An alternative approach to assessing cardiovascular risk using biomarkers is to assay markers of systemic inflammation, for example C-reactive protein (CRP) or fibrinogen, which also correlate with risk of cardiovascular events [5].

We hypothesised that patients with bronchiectasis would have elevated cardiovascular risk, not adequately assessed using QRISK2. The aim of this study was to identify which disease factors were significantly associated with elevated cardiovascular risk measurements in bronchiectasis, and furthermore, to determine whether measured risk was being predicted accurately by the calculated QRISK2 score.

Methods

Study subjects

101 patients with bronchiectasis were recruited as a convenience sample from clinics at the Royal Free London and University College London Hospitals National Health Service foundation trusts. The study was approved by the local research ethics committee (reference 10/H0720/43) and written informed consent was obtained from all participants.

Bronchiectasis was confirmed on computed tomography (CT). Clinically significant bronchiectasis was defined as the presence of either chronic productive cough or recurrent respiratory infections together with a compatible CT scan [1]. Patients were investigated to determine the aetiology of bronchiectasis in accordance with the British Thoracic Society 2010 guideline [1]. Those with a primary diagnosis of COPD were excluded, because of the recognised association of COPD with increased cardiovascular risk and since airway wall thickening and dilatation are common incidental radiological findings in COPD [21]. Patients were excluded if they were experiencing an acute worsening of respiratory symptoms, receiving treatment for an exacerbation or had completed treatment for exacerbation within the past 2 weeks.

Clinical assessment

We collated variables allowing calculation of the bronchiectasis severity index (BSI), available in 96 patients [22]. This composite score comprises age, BMI, forced expiratory volume in 1 s (FEV₁), exacerbations within the past year, hospitalisations within the past 2 years, Medical Research Council dyspnoea score, sputum colonisation with *Pseudomonas aeruginosa* or other potentially pathogenic micro-organisms and the number of lobes involved on CT. Disease is categorised as mild (≤ 4 points), moderate (5–8 points) or severe (> 8 points). Exacerbation history was defined according to healthcare utilisation and was based on patient recall [23]. Not all the variables were available in all patients: existing CT scans were available in 99 patients, sputum data were available from 77 patients and spirometry was available in 97 patients. In the absence of sputum culture results, noncolonisation was assumed, but where other data were missing BSI was not calculated. Diameter of the pulmonary artery and ascending aorta were measured at the pulmonary artery bifurcation using an established method [24]. 86 CT scans were suitable for this measurement.

PWV

Carotid–femoral PWV was measured in all patients as in our previous report [5], by one of two operators using the Vicorder system (Skidmore Medical, Bristol, UK). Values were measured in triplicate and the mean value used for analysis [6]. A pilot study performed on 15 healthy volunteers by the two operators (ADS and BK) demonstrated good individual and interobserver consistency assessed using intraclass correlation coefficients (ICC): operator 1 ICC=0.975, operator 2 ICC=0.901 and agreement between operators ICC=0.969.

Reference ranges for PWV have been published by the Reference Values for Arterial Stiffness Collaboration using large European populations of healthy subjects [25]. This is a composite analysis of multiple studies normalised to reflect values detected using the Sphygmocor device (AtCor Medical, West Ryde, NSW, Australia). These have been demonstrated to correlate well with measurements made using the Vicorder when path length is calculated using the midpoint of the femoral cuff [26].

Serum analysis

CRP, N-terminal (NT)-proBNP and troponin T were measured in 100, 97 and 98 patients, respectively, on the day of the study visit using a Modular Analytics E 170 module (Roche Diagnostics, Burgess Hill, UK) with limits of detection of 1 mg·L⁻¹, 2 pg·mL⁻¹ and 0.003 mg·L⁻¹, respectively. Plasma fibrinogen was analysed in 78 patients using the Clauss method (IL ACL Top Coagulation Analyzer; Instrumentation Laboratory, Lexington, MA, USA), with a limit of detection at 1.5 g·L⁻¹. As fibrinogen concentration was strongly correlated with CRP concentration, the missing 23 fibrinogen values were imputed by regression against CRP. Results for CRP and fibrinogen have been reported previously [27].

Statistical analysis

Analysis was performed using SPSS (version 22; IBM Corporation, Armonk, NY, USA). Normality was assessed using the Kolmogorov–Smirnov test. Normally distributed data are expressed as mean \pm SD and skewed data as median (interquartile range (IQR)). Pearson or Spearman rank correlations were applied when analysing two continuous variables, as appropriate. Paired and unpaired t-tests and Mann–Whitney U-tests were used when comparing means/ranks between two groups and one-way ANOVA or Kruskal–Wallis test were employed for comparing means/ranks between multiple groups. Multivariate regression analysis was used to determine which components of BSI explained most variance in serum biomarkers. We assumed $p \leq 0.05$ as statistically significant, unless specifically stated. Hochberg correction was performed for the multiple comparisons reported in table 2 [28]. This method controls the false discovery rate with a lower risk of type 2 error than the highly conservative Bonferroni correction [29].

Results

101 patients with bronchiectasis were recruited. Their baseline characteristics, bronchiectasis severity indices and the results of aortic stiffness (PWV) and cardiac biomarker measurements (PWV, troponin T, NT-proBNP, fibrinogen and CRP) are reported in table 1. As would be expected, the majority of subjects (n=63) were female, and the median age was 61 years. While idiopathic and post-infectious disease were common (as would be seen in most cohorts), the single commonest aetiology in our subjects was primary immune deficiency (mainly common variable immunodeficiency), reflecting local clinical practice.

Correlates of elevated cardiovascular risk in bronchiectasis

The relationships between patient or disease characteristics and cardiovascular risk are reported in table 2. There was a significant relationship between the overall severity of bronchiectasis as assessed by BSI score and all of the cardiac risk assessments. However, as age is a component of BSI and is associated with

TABLE 1 Baseline characteristics, bronchiectasis severity indices and assessment of pulse-wave velocity and cardiac biomarkers in 101 patients with bronchiectasis

Subjects	101
Age years	61.0 (46.1–71.6)
Male	38 (38)
Current smoker	2 (2)
Ex-smoker	28 (28)
Pack-years (current/ex-smokers)	3.5 (1–15)
Body mass index kg·m⁻²	23.0 (20.9–25.7)
Systolic blood pressure mmHg	126 (115–139)
Diastolic blood pressure mmHg	80 (70–85)
Total cholesterol mmol·L⁻¹	4.8±0.4
Aetiology	
Primary immunodeficiency [#]	28 (28)
Idiopathic	21 (21)
Post-infective	19 (19)
Allergic bronchopulmonary aspergillosis	9 (9)
Other [¶]	24 (24)
Long-term macrolide	19 (19)
Long-term nebulised antibiotic	3 (3)
Inhaled corticosteroid	42 (42)
FEV₁ % predicted	72.7±27.0
FEV₁ L	1.94±0.88
Exacerbation frequency events·year⁻¹	2 (1–3)
Time since last exacerbation days	171 (62–388)
Lobes involved on CT n	4 (3–6)
Pulmonary artery diameter mm	25.5±4.0
Pulmonary artery:aorta ratio	0.77 (0.70–0.84)
MRC dyspnoea score	2 (1–3)
Bronchiectasis severity index	5.5 (3–8)
Pulse-wave velocity m·s⁻¹	9.3 (8.1–10.9)
Troponin T µg·L⁻¹	0.006 (0.003–0.009)
NT-proBNP pmol·L⁻¹	10.0 (4.5–17.0)
Fibrinogen g·L⁻¹	3.31 (2.69–4.12)
C-reactive protein mg·L⁻¹	3.2 (1.0–7.7)

Data are presented as n, median (interquartile range), n (%) or mean±sd. FEV₁: forced expiratory volume in 1 s; CT: computed tomography; MRC: Medical Research Council; NT-proBNP: N-terminal pro-brain natriuretic peptide. [#]: common variable immunodeficiency (n=23), IgG deficiency (n=4), X-linked agammaglobulinaemia (n=1); [¶]: rheumatoid arthritis (n=5), aspiration (n=3), Kartagener's syndrome (n=3), HIV (n=2), graft *versus* host disease (n=2), sarcoidosis (n=2), systemic lupus erythematosus (n=1), secondary immunodeficiency (n=1), surgery-related (n=1), post-radiotherapy (n=1), non-tuberculous mycobacteria (n=1), inflammatory bowel disease (n=1), Mounier-Kuhn syndrome (n=1).

increasing cardiovascular risk (confirmed in the table) we went on to explore relationships between cardiovascular risk and key indices of bronchiectasis severity, including exacerbation frequency, lung function impairment and extent of disease on CT.

Cardiovascular risk and systemic inflammation were both associated with exacerbation frequency. Patients with three or more exacerbations per year had a significantly higher median (IQR) CRP concentration (4.8 (2.0–11.8) *versus* 2.2 (0.5–5.4) mg·L⁻¹, p=0.005), as illustrated in figure 1, higher mean PWV (10.5 *versus* 9.2 m·s⁻¹, p=0.01) and elevated troponin, suggesting subclinical myocardial injury. Multivariable analysis assessing which components of BSI predicted CRP confirmed an independent relationship with exacerbation frequency (β=0.38, p=0.001) and FEV₁ % predicted (β=-0.23, p=0.029).

There was a correlation between radiological disease severity as assessed by the number of lobes involved on CT and the NT-proBNP concentration (r=0.24, p=0.02), although this was not statistically significant after Hochberg correction for multiple comparisons. To investigate whether any association between radiological extent and cardiovascular strain (NT-proBNP) might be caused by pulmonary hypertension, we measured pulmonary artery diameter and the ratio between diameter of the pulmonary artery and the ascending aorta (PA:A ratio), both of which correlate with invasive measures of pulmonary hypertension [30, 31]. As only four patients had a PA:A ratio >1, we analysed the relationship between pulmonary artery diameter or PA:A ratio and BNP. There were no significant correlations.

TABLE 2 Relationships between assessments of cardiovascular risk and BSI/selected individual components

	PWV $\text{m}\cdot\text{s}^{-1}$		Troponin $\mu\text{g}\cdot\text{L}^{-1}$		NT-proBNP $\text{pmol}\cdot\text{L}^{-1}$		Fibrinogen $\text{g}\cdot\text{L}^{-1}$		CRP $\text{mg}\cdot\text{L}^{-1}$	
	r/ ρ	p-value	r/ ρ	p-value	r/ ρ	p-value	r/ ρ	p-value	r/ ρ	p-value
Subjects n	101		98		97		100		100	
BSI	0.45	<0.001[#]	0.52	<0.001[#]	0.34	0.001[#]	0.39	0.001[#]	0.24	0.017[¶]
Age	0.77	<0.001[#]	0.59	<0.001[#]	0.47	<0.001[#]	0.28	0.012[¶]	0.10	0.323
Annual exacerbation frequency										
≥3 events·year ⁻¹ (n=40)	10.5±2.7	0.011[¶]	0.008 (0.004–0.013)	0.047[¶]	11.0 (5.0–18.5)	0.293	3.5 (2.7–4.3)	0.253	4.8 (2.0–11.8)	0.005[¶]
<3 events·year ⁻¹ (n=61)	9.2±1.7		0.005 (0.003–0.008)		9.0 (3.0–17.0)		3.2 (2.7–3.9)		2.2 (0.5–5.4)	
FEV ₁ L	-0.34	0.001[#]	-0.42	<0.001[#]	-0.34	0.001[#]	-0.53	<0.001[#]	-0.45	<0.001[#]
FEV ₁ % predicted	-0.02	0.832	-0.28	0.006[¶]	-0.11	0.301	-0.43	<0.001[#]	-0.34	0.001[#]
CT lobes involved	0.14	0.159	0.17	0.091	0.24	0.02[¶]	-0.07	0.570	0.07	0.482

Data are presented as mean±SD or median (interquartile range), unless otherwise stated. Significant findings are presented in bold. BSI: bronchiectasis severity index; PWV: pulse-wave velocity; NT-proBNP: N-terminal pro-brain natriuretic peptide; CRP: C-reactive protein; FEV₁: forced expiratory volume in 1 s; CT: computed tomography. [#]: significant after Hochberg adjustment for multiple comparisons; [¶]: significant at $\alpha<0.05$.

Lung function impairment (FEV₁ % pred) was inversely correlated with troponin ($r=-0.28$, $p=0.006$), CRP ($r=-0.34$, $p=0.001$) and fibrinogen ($r=-0.43$, $p<0.001$). All of the markers were significantly and inversely related to absolute FEV₁, such that patients with lower FEV₁ have elevated markers of cardiovascular risk, and increased systemic inflammation.

The interrelationships between the four assessments of cardiovascular risk are reported in table 3. All were significantly correlated with each other, with the exception of NT-proBNP with both CRP and fibrinogen. This suggests that assay of cardiac biomarkers, readily available in many healthcare settings, may be able to reflect increased cardiovascular risk when compared to the “gold standard” assessment of PWV.

Assessment of cardiovascular risk using QRISK2

Our cohort had significantly greater cardiovascular risk assessed using PWV than that published for age-matched controls [25]. 69 (68%) patients had a measured PWV higher than the healthy, age-adjusted mean. The mean difference between the expected and observed PWV of $0.9\pm 1.6 \text{ m}\cdot\text{s}^{-1}$ was highly statistically significant ($p<0.001$), and associated with a median (IQR) change in relative risk of adverse cardiovascular outcome of 1.29 (0.96–1.61) [32].

The QRISK2 calculator includes age, sex, ethnicity, postal (zip) code (as a marker of socioeconomic status), family history of ischaemic heart disease, presence of diabetes, chronic kidney disease, atrial fibrillation and/or rheumatoid arthritis, systolic blood pressure, cholesterol and BMI. It estimates the 10-year absolute risk of myocardial infarction and stroke and the relative risk of these outcomes compared to a healthy individual of the same age, sex and ethnicity [11]. Compared to a healthy individual, 30

FIGURE 1 Exacerbation frequency in bronchiectasis is associated with increased systemic inflammation in the stable state as assessed by C-reactive protein [CRP] concentration (n=100, $2.2 \text{ mg}\cdot\text{L}^{-1}$ versus $4.8 \text{ mg}\cdot\text{L}^{-1}$ for fewer than three exacerbations per year and three or more exacerbations per year, respectively; $p=0.005$).




TABLE 3 Relationships between individual assessments of cardiovascular risk in 101 patients with bronchiectasis

	Troponin $\mu\text{g}\cdot\text{L}^{-1}$		NT-proBNP $\text{pmol}\cdot\text{L}^{-1}$		Fibrinogen $\text{g}\cdot\text{L}^{-1}$		CRP $\text{mg}\cdot\text{L}^{-1}$	
	r/ ρ	p-value	r/ ρ	p-value	r/ ρ	p-value	r/ ρ	p-value
Subjects n	98		97		78		100	
Pulse-wave velocity $\text{m}\cdot\text{s}^{-1}$	0.49	<0.001	0.27	0.008	0.24	0.036	0.20	0.043
CRP $\text{mg}\cdot\text{L}^{-1}$	0.20	0.045	0.01	0.258	0.55	<0.001		
Fibrinogen $\text{g}\cdot\text{L}^{-1}$	0.37	0.001	0.19	0.090				
NT-proBNP $\text{pmol}\cdot\text{L}^{-1}$	0.44	<0.001						

Significant findings are presented in bold. NT-proBNP: N-terminal pro-brain natriuretic peptide; CRP: C-reactive protein.

(30%) patients had their risk increased and 65 (64%) patients had their risk decreased after adjustment for these variables. Risk was unchanged in six patients.

Next, we compared the QRISK2-derived relative risk with a relative risk estimate calculated from the patient's measured PWV, as follows. Meta-analysis data of cardiovascular end-points in relation to PWV has concluded that each standard deviation increase in \log_e PWV is associated with a 1.45-fold increase in relative risk of cardiovascular events (myocardial infarction and stroke) at 5 years [32]. Using this assumption, the baseline relative risk of cardiovascular event was recalculated for each subject according to their PWV. For example, a subject aged 56 years with a PWV of $10.2 \text{ m}\cdot\text{s}^{-1}$ has a \log_e PWV of 2.32, which lies 1.2 SD above the median \log_e PWV for healthy patients aged 50–60 years. Their relative risk therefore increases to 1.56 ($1.45^{1.2}$).

Calculated QRISK2 estimated lower relative risk of adverse cardiovascular outcomes than measured PWV in our population of patients with bronchiectasis. There was a mean \pm SD difference of 0.42 ± 0.78 between the relative risk calculated from QRISK2 and relative risk calculated from PWV ($p<0.001$). The different distribution of cardiovascular risk measured by PWV compared to that calculated by QRISK2 is illustrated in figure 2. 73 (74%) patients had a higher cardiovascular risk measured by PWV than calculated by QRISK2.

Finally, we went on to examine whether there were specific patient demographic or bronchiectasis characteristics that were associated with underestimation of risk using QRISK2 compared to PWV. We examined this by comparing these variables against the magnitude of QRISK2 risk underestimation. The data are presented in table 4, which demonstrates that underestimation of cardiovascular risk by QRISK2 compared to PWV was significantly associated with male sex and increased exacerbation frequency.

Discussion

The key findings of our study are that patients with bronchiectasis have greater measured cardiovascular risk than published reference control populations, that excess cardiovascular risk in bronchiectasis is associated with markers of bronchiectasis severity, specifically exacerbation frequency and impaired lung




FIGURE 2 Relative risk of adverse cardiovascular events calculated by QRISK2 compared to risk conferred by measured pulse-wave velocity (PWV), both in relation to that expected for age-, sex-, ethnicity- and socioeconomic status-matched controls. Mean \pm SD difference in risk 0.42 ± 0.78 ($p<0.001$).

TABLE 4 Relationships between patient and bronchiectasis indices, and the magnitude of cardiovascular risk underestimation using QRISK2 compared with pulse-wave velocity

	Degree of relative risk underestimate by QRISK2	p-value
Age	-0.16	0.12
Sex		
Male	0.62±0.85	0.037
Female	0.29±0.70	
BSI	0.06	0.59
FEV₁	0.06	0.54
FEV₁ % predicted	-0.12	0.23
Lobes involved on CT	-1.0	0.34
Exacerbation frequency	0.26	0.008
MRC score	0.04	0.68
Systolic blood pressure	-0.03	0.78
Fibrinogen	0.01	0.94
Troponin	-0.05	0.65
NT-proBNP	-0.06	0.59
CRP	0.17	0.09

Data are presented as r, ρ or mean \pm SD, unless otherwise stated. Significant findings are presented in bold. BSI: bronchiectasis severity index; FEV₁: forced expiratory volume in 1 s; CT: computed tomography; MRC: Medical Research Council; NT-proBNP: N-terminal pro-brain natriuretic peptide; CRP: C-reactive protein.

function and that standard calculators of cardiovascular risk such as QRISK2 may underestimate measured cardiovascular risk in bronchiectasis, especially in males and in patients susceptible to frequent exacerbations. Given the biological plausibility of exacerbations driving cardiovascular damage through mechanisms such as plaque rupture [33] and microthrombosis tendency [34], it follows that exacerbation prevention may result in fewer cardiovascular events in these patients. These findings suggest that cardiovascular risk assessment in bronchiectasis should be individualised and not rely solely on calculation tools suitable for populations without the disease.

This is the largest study of individual cardiovascular risk in bronchiectasis ever reported. The first suggestion that bronchiectasis may be associated with cardiac dysfunction was a report by ALZEER *et al.* [35] in 2008, demonstrating right ventricular impairment in association with pulmonary hypertension in patients with more severe, cystic disease. We have recently used population database analysis to demonstrate a higher risk of ischaemic cardiovascular events in patients with bronchiectasis [9] and we therefore wished to investigate cardiovascular risk in a carefully phenotyped cohort of patients with bronchiectasis. PWV is a gold-standard assessment of arterial stiffness that we have previously reported as stable over time and reproducible [6]. PWV provides additional predictive value beyond estimates based on existing cardiovascular risk factors [17]. Our findings are consistent with those published in cystic fibrosis [36] as well as those of GALE *et al.* [8] who reported a higher aortic PWV in 20 patients with bronchiectasis compared to controls. Greater aortic stiffness directly impacts cardiovascular events, as there is premature return of the reflected pulse wave during late systole, which impairs normal diastolic filling of the coronary arteries. In addition, stiffer central arteries impose a greater afterload on the heart and increase myocardial work, which over time may result in ventricular dysfunction [37]. In support of this, we found a significant correlation between aortic stiffness and the concentration of NT-proBNP.

We next went on to explore which features of bronchiectasis were associated with increased cardiovascular risk. While the composite BSI related strongly to aortic stiffness, both are age-dependent. Regarding specific features of bronchiectasis, the factors significantly associated with aortic stiffness were exacerbations and FEV₁. The elevation in pulse-wave velocity of 1.3 m·s⁻¹ in patients experiencing three or more exacerbations annually, *versus* infrequent exacerbators, equates to a clinically significant excess risk of 12–18% [38]. Therefore, this difference is large enough to be associated with adverse outcomes in patients susceptible to frequent exacerbations. We used a threshold of three exacerbations per year to define frequent exacerbations, as this is used in the BSI score [22] and bronchiectasis guidelines [1]. It is known that exacerbations affect quality of life in bronchiectasis [39] and our findings extend the rationale for exacerbation prevention to include possible reduction in adverse cardiovascular outcomes.

We have previously defined an association between exacerbation frequency and cardiovascular risk in COPD [6]. Although this association in COPD may be confounded by smoking, chronic inflammatory conditions such as rheumatoid arthritis are also associated with elevated vascular risk, supporting chronic

inflammation itself as a driving force [40]. Similarly, in systemic lupus erythematosus cardiovascular disease accounts for 50% of mortality and is strongly associated with markers of inflammation including fibrinogen and CRP [41]. Our finding of higher baseline CRP in frequent exacerbators supports a mechanism of chronic inflammation underlying the elevated cardiovascular risk. While the concept of an independent exacerbation susceptibility phenotype associated with poor clinical outcomes is accepted in COPD [42], further work is required to establish whether the same is true in bronchiectasis. However, we have reported that increased systemic inflammation is seen even in community-treated exacerbations [43].

In addition, greater impairment in lung function, as assessed by FEV₁, was associated with both increased systemic inflammation and elevation of cardiac risk. It has previously been reported that smaller lungs are associated with increased arterial stiffness and cardiovascular death in a general population, which reflects our finding in this disease group [44]. The extent of disease on CT was associated with elevation of BNP (uncorrected analysis), in support of the study described earlier in which more severe cystic disease was associated with ventricular dysfunction [35]. We postulated that pulmonary hypertension may be the mechanism underlying this association and performed a univariate analyses between PA:A ratio and pulmonary artery diameter, both established surrogates of pulmonary hypertension [30, 31], and BNP and CT lobar extent. Similarly to DEVARAJ *et al.* [45], we found no correlation between radiological extent and CT measures of pulmonary hypertension. We hypothesise that the link between BNP, a measure of cardiac impairment, and CT extent of disease is through systemic inflammation causing cardiac strain, rather than specifically through pulmonary vascular disease.

While CT extent was not related to arterial stiffness, the extent of disease did relate to cardiac impairment, and lung function to markers of myocardial damage and systemic inflammation. These findings suggest that preservation of lung function and minimisation of disease extent are additional important goals in bronchiectasis management with regard to the mitigation of future cardiovascular disease.

Having identified increased cardiovascular risk in bronchiectasis, the next question we addressed was how this should best be assessed. We demonstrated that the QRISK2 score systematically underestimated cardiovascular risk in bronchiectasis compared to calculating risk using PWV data, particularly in male patients, and those experiencing frequent exacerbations. We have shown that the latter group has elevated systemic inflammation in the stable state. We speculate that including inflammatory conditions such as bronchiectasis in cardiovascular risk calculators may improve predictive value. We suggest that an individual approach to cardiovascular risk assessment is required, with the use of additional tools where calculated risk is intermediate. This might include assessment of cardiac biomarkers (which correlated with PWV in this study) or PWV itself, or indeed alternative approaches such as coronary artery calcium score. The latter is attractive given the need for a chest CT scan to make the diagnosis of bronchiectasis. Biochemical analysis of cardiac biomarkers is probably the easiest measure to perform in routine practice, but age-specific values from healthy populations are not available and we defined significant relationships between both troponin and BNP with age in this study. We cannot comment on whether other risk scores underestimate cardiovascular risk in bronchiectasis; an idealised prospective study would be to measure risk with sufficient follow-up to record a significant number of future cardiovascular events.

Our results emphasise the importance of exacerbation prevention strategies in bronchiectasis, not only to improve quality of life and to reduce lung morbidity, but also to potentially prevent cardiovascular disease events. Our data also suggest that anti-inflammatory therapies may have the additional benefit of reducing cardiovascular risk, which should be considered as an exploratory end-point in future clinical trials. Although there are currently no licensed anti-inflammatory interventions available in bronchiectasis, we have recently demonstrated that specific components of systemic inflammation in bronchiectasis are related to disease severity and aetiology, and elucidating the underlying pathways may identify novel therapeutic approaches [27].

There were some limitations to this study. Patients with bronchiectasis related to primary immunodeficiency are overrepresented in this sample, reflecting our local referrals; this is a limitation of any single-centre study in bronchiectasis and there can never be a “typical” population. The 28 subjects with primary immunodeficiency-related bronchiectasis were removed for an exploratory subanalysis in order to determine whether overrepresentation of this group may have skewed results. PWV remained significantly greater in frequent exacerbators (≥ 3 events-year⁻¹) compared to infrequent exacerbators (10.8 *versus* 9.4 m·s⁻¹, $p=0.005$), as did CRP (3.9 (2.0–11.0) *versus* 2.3 (0.9–5.2) mg·L⁻¹, $p=0.019$). Relative risk remained greater when measured by PWV compared to QRISK score (0.86 (0.69–1.04) *versus* 1.33 (0.96–1.67), $p<0.001$). This demonstrates that, despite reduced statistical power, our main results are robust to the removal of patients with primary immunodeficiency from the analysis.

In other limitations, the study may have been underpowered to detect weaker correlations, and we did not perform an *a priori* power calculation. We cannot exclude the possibility that the relationship between

frequent exacerbators and arterial stiffness represents, in part, elevated PWV persisting from the preceding exacerbation. We used a novel approach to calculate the excess cardiovascular risk attributable to elevated arterial stiffness. This relied on the use of a reference range for PWV and assumes that the relative risk increase attributable to a given increase in arterial stiffness is attributable to our study population. BEN-SHLOMO *et al.* [32] made PWV-based adjustments to the relative risk of cardiovascular events over 5 years, whereas QRISK calculates a 10-year risk. We made the assumption that changes in PWV would adjust the relative risk of cardiovascular events over 5 and 10 years in a linear way.

In conclusion, we confirm that bronchiectasis is associated with increased cardiovascular risk, and define that this is related in part to the frequency of exacerbations and impairment in lung function. Assessment of cardiovascular risk using tools such as QRISK2 could underestimate risk in some patients, especially males and those susceptible to frequent exacerbations. Therefore, an individual approach should be used for patients with intermediate risk. Awareness and mitigation of existing cardiovascular risk factors, and exacerbation prevention and anti-inflammatory therapy in bronchiectasis all have the potential to reduce cardiovascular events and require further study if we are to improve clinical outcomes in this neglected disease.

Acknowledgements

We are grateful for advice received from Reecha Sofat (University College London, London, UK) on cardiovascular risk scores. We thank the patients with bronchiectasis for their willingness to take part in our studies and clinic staff at both Trusts for their help in facilitating our research. We thank Anant Patel (Royal Free London NHS Foundation Trust, London) for assistance with aortic pulse-wave velocity training.

References

- 1 Pasteur MC, Bilton D, Hill AT. British Thoracic Society guideline for non-CF bronchiectasis. *Thorax* 2010; 65: Suppl. 1, i1–i58.
- 2 Cole PJ. Inflammation: a two-edged sword – the model of bronchiectasis. *Eur J Respir Dis Suppl* 1986; 147: 6–15.
- 3 Roberts HJ, Hubbard R. Trends in bronchiectasis mortality in England and Wales. *Respir Med* 2010; 104: 981–985.
- 4 McDonnell MJ, Aliberti S, Goeminne PC, *et al.* Comorbidities and the risk of mortality in patients with bronchiectasis: an international multicentre cohort study. *Lancet Respir Med* 2016; 4: 969–979.
- 5 Kaptoge S, Di Angelantonio E, Pennells L, *et al.* C-reactive protein, fibrinogen, and cardiovascular disease prediction. *N Engl J Med* 2012; 367: 1310–1320.
- 6 Patel AR, Kowlessar BS, Donaldson GC, *et al.* Cardiovascular risk, myocardial injury, and exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2013; 188: 1091–1099.
- 7 Wilson CB, Jones PW, O’Leary CJ, *et al.* Systemic markers of inflammation in stable bronchiectasis. *Eur Respir J* 1998; 12: 820–824.
- 8 Gale NS, Bolton CE, Duckers JM, *et al.* Systemic comorbidities in bronchiectasis. *Chron Respir Dis* 2012; 9: 231–238.
- 9 Navaratnam V, Millett ER, Hurst JR, *et al.* Bronchiectasis and the risk of cardiovascular disease: a population-based study. *Thorax* 2017; 72: 161–166.
- 10 Rabar S, Harker M, O’Flynn N, *et al.* Lipid modification and cardiovascular risk assessment for the primary and secondary prevention of cardiovascular disease: summary of updated NICE guidance. *BMJ* 2014; 349: g4356.
- 11 Hippisley-Cox J, Coupland C, Vinogradova Y, *et al.* Predicting cardiovascular risk in England and Wales: prospective derivation and validation of QRISK2. *BMJ* 2008; 336: 1475–1482.
- 12 Goff DC Jr, Lloyd-Jones DM, Bennett G, *et al.* 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014; 63: 2935–2959.
- 13 Meaume S, Benetos A, Henry OF, *et al.* Aortic pulse wave velocity predicts cardiovascular mortality in subjects >70 years of age. *Arterioscler Thromb Vasc Biol* 2001; 21: 2046–2050.
- 14 Laurent S, Boutouyrie P, Asmar R, *et al.* Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. *Hypertension* 2001; 37: 1236–1241.
- 15 Blacher J, Guerin AP, Pannier B, *et al.* Impact of aortic stiffness on survival in end-stage renal disease. *Circulation* 1999; 99: 2434–2439.
- 16 Davies JI, Struthers AD. Pulse wave analysis and pulse wave velocity: a critical review of their strengths and weaknesses. *J Hypertens* 2003; 21: 463–472.
- 17 Mitchell GF, Hwang SJ, Vasan RS, *et al.* Arterial stiffness and cardiovascular events: the Framingham Heart Study. *Circulation* 2010; 121: 505–511.
- 18 Benetos A, Thomas F, Bean KE, *et al.* Why cardiovascular mortality is higher in treated hypertensives *versus* subjects of the same age, in the general population. *J Hypertens* 2003; 21: 1635–1640.
- 19 Saunders JT, Nambi V, de Lemos JA, *et al.* Cardiac troponin T measured by a highly sensitive assay predicts coronary heart disease, heart failure, and mortality in the Atherosclerosis Risk in Communities Study. *Circulation* 2011; 123: 1367–1376.
- 20 Wang TJ, Larson MG, Levy D, *et al.* Plasma natriuretic peptide levels and the risk of cardiovascular events and death. *N Engl J Med* 2004; 350: 655–663.
- 21 Patel IS, Vlahos I, Wilkinson TM, *et al.* Bronchiectasis, exacerbation indices, and inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004; 170: 400–407.
- 22 Chalmers JD, Goeminne P, Aliberti S, *et al.* The bronchiectasis severity index. An international derivation and validation study. *Am J Respir Crit Care Med* 2014; 189: 576–585.

- 23 Quint JK, Donaldson GC, Hurst JR, *et al.* Predictive accuracy of patient-reported exacerbation frequency in COPD. *Eur Respir J* 2011; 37: 501–507.
- 24 Wells JM, Washko GR, Han MK, *et al.* Pulmonary arterial enlargement and acute exacerbations of COPD. *N Engl J Med* 2012; 367: 913–921.
- 25 Reference Values for Arterial Stiffness Collaboration. Determinants of pulse wave velocity in healthy people and in the presence of cardiovascular risk factors: ‘establishing normal and reference values’. *Eur Heart J* 2010; 31: 2338–2350.
- 26 Hickson SS, Butlin M, Broad J, *et al.* Validity and repeatability of the Vicorder apparatus: a comparison with the SphygmoCor device. *Hypertens Res* 2009; 32: 1079–1085.
- 27 Saleh AD, Chalmers JD, De Soyza A, *et al.* The heterogeneity of systemic inflammation in bronchiectasis. *Respir Med* 2017; 127: 33–39.
- 28 Hochberg Y. A sharper Bonferroni procedure for multiple tests of significance. *Biometrika* 1988; 75: 800–802.
- 29 Bender R, Lange S. Adjusting for multiple testing – when and how? *J Clin Epidemiol* 2001; 54: 343–349.
- 30 Pérez-Enguix D, Morales P, Tomás JM, *et al.* Computed tomographic screening of pulmonary arterial hypertension in candidates for lung transplantation. *Transplant Proc* 2007; 39: 2405–2408.
- 31 Ng CS, Wells AU, Padley SP. A CT sign of chronic pulmonary arterial hypertension: the ratio of main pulmonary artery to aortic diameter. *J Thorac Imaging* 1999; 14: 270–278.
- 32 Ben-Shlomo Y, Spears M, Boustred C, *et al.* Aortic pulse wave velocity improves cardiovascular event prediction: an individual participant meta-analysis of prospective observational data from 17,635 subjects. *J Am Coll Cardiol* 2014; 63: 636–646.
- 33 Bazaz R, Marriott HM, Francis SE, *et al.* Mechanistic links between acute respiratory tract infections and acute coronary syndromes. *J Infect* 2013; 66: 1–17.
- 34 Maclay JD, McAllister DA, Johnston S, *et al.* Increased platelet activation in patients with stable and acute exacerbation of COPD. *Thorax* 2011; 66: 769–774.
- 35 Alzeer AH, Al-Mobeirek AF, Al-Otair HA, *et al.* Right and left ventricular function and pulmonary artery pressure in patients with bronchiectasis. *Chest* 2008; 133: 468–473.
- 36 Hull JH, Garrod R, Ho TB, *et al.* Increased augmentation index in patients with cystic fibrosis. *Eur Respir J* 2009; 34: 1322–1328.
- 37 Vlachopoulos C, Alexopoulos N, Stefanadis C. Fast in the aorta, slow in the coronaries. *Cardiology* 2010; 116: 257–260.
- 38 Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol* 2010; 55: 1318–1327.
- 39 Wilson CB, Jones PW, O’Leary CJ, *et al.* Validation of the St. George’s Respiratory Questionnaire in bronchiectasis. *Am J Respir Crit Care Med* 1997; 156: 536–541.
- 40 del Rincón ID, Williams K, Stern MP, *et al.* High incidence of cardiovascular events in a rheumatoid arthritis cohort not explained by traditional cardiac risk factors. *Arthritis Rheum* 2001; 44: 2737–2745.
- 41 Gustafsson JT, Simard JF, Gunnarsson I, *et al.* Risk factors for cardiovascular mortality in patients with systemic lupus erythematosus, a prospective cohort study. *Arthritis Res Ther* 2012; 14: R46.
- 42 Hurst JR, Vestbo J, Anzueto A, *et al.* Susceptibility to exacerbation in chronic obstructive pulmonary disease. *N Engl J Med* 2010; 363: 1128–1138.
- 43 Brill SE, Patel AR, Singh R, *et al.* Lung function, symptoms and inflammation during exacerbations of non-cystic fibrosis bronchiectasis: a prospective observational cohort study. *Respir Res* 2015; 16: 16.
- 44 Zureik M, Benetos A, Neukirch C, *et al.* Reduced pulmonary function is associated with central arterial stiffness in men. *Am J Respir Crit Care Med* 2001; 164: 2181–2185.
- 45 Devaraj A, Wells AU, Meister MG, *et al.* Pulmonary hypertension in patients with bronchiectasis: prognostic significance of CT signs. *AJR Am J Roentgenol* 2011; 196: 1300–1304.