



## Early View

Editorial

### **Asthma and COVID-19: do we finally have answers?**

Katrien Eger, Elisabeth H. Bel

Please cite this article as: Eger K, Bel EH. Asthma and COVID-19: do we finally have answers?. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.04451-2020>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©ERS 2020. This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.

Editorial

## **Asthma and COVID-19: do we finally have answers?**

Katrien Eger MD<sup>1</sup>, Elisabeth H. Bel MD, PhD<sup>1</sup>

<sup>1</sup>Department of Respiratory Medicine, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Corresponding author:

Name: Elisabeth Bel

Address: Amsterdam UMC, University of Amsterdam

Dept. of Respiratory Medicine F5-152

Meibergdreef 9

1105 AZ Amsterdam, The Netherlands

Phone: +31205664356

Fax: +31205669001

Email: [e.h.bel@amsterdamumc.nl](mailto:e.h.bel@amsterdamumc.nl)

Word count body: 1365

Conflict of Interest Statement: KE has nothing to declare. EHB Dr. Bel reports grants and personal fees from AstraZeneca, grants and personal fees from GSK, grants and personal fees from Novartis, personal fees from Sanofi/Regeneron, grants and personal fees from Teva, personal fees from Sterna, personal fees from Chiesi, outside the submitted work.

## Text

The 2019 coronavirus disease (COVID-19) pandemic has already claimed the lives of nearly 1.5 million people and the virus is continuing to spread across the world.[1] The disease can affect anyone, and although SARS-CoV-2 infection leads to mild COVID-19 in the majority of cases, the proportion of patients developing severe pneumonia is of great concern.

It is now well-recognized that older age, obesity, cardiovascular disease and diabetes are risk factors of poor COVID-19 outcome.[2–4] What is not yet clear is whether chronic respiratory diseases like asthma are amongst the risk factors as well. The many studies that have addressed this question show discrepant results and point towards numerous factors that may play a role in the susceptibility and severity of COVID 19 in asthma patients.[5–10] These include the severity of asthma itself, the asthma phenotype (allergic or non-allergic), asthma medication (corticosteroids or no corticosteroids), and co-morbidities.[11–15] Because of this complex interplay between numerous factors involved, there is a need for large-scale studies that allow adjustment for confounders, making it possible to evaluate the true impact of asthma on the susceptibility and outcome of COVID-19.

This is exactly the approach taken by Choi *et al.* and Izquierdo *et al.* in the two real-life studies in this issue of the European Respiratory Journal.[16,17] While Choi *et al.* included all COVID-19 cases in Korea (n=7,590) by using a national claims database, Izquierdo *et al.* analyzed medical record data from 71,182 patients with asthma who attended medical services from a region in Spain. Choi *et al.* found a higher frequency of asthma in the COVID-19 population compared to the general population (2.9 vs 1.6-2.2%), while Izquierdo *et al.* found a higher frequency of COVID-19 amongst asthma patients compared to the general population (1.41 vs 0.86%). In addition, Izquierdo *et al.* showed an even higher incidence of COVID-19 in patients on biologic therapy (2.31%). Thus, both studies suggest that the susceptibility

for contracting COVID-19 in asthma patients is higher than in the general population, especially in those with severe asthma on biologic therapy.

Regarding the severity of COVID-19, Izquierdo *et al.* reported equal hospitalization rates in asthma and non-asthma patients with COVID-19 (in both cases 26%). Likewise, Choi *et al.* found that COVID-19 severity as illustrated by ICU admission rate and duration were similar between COVID-19 patients with and without asthma. Although mortality rates were higher in patients with asthma, asthma and asthma severity did not show to be significant predictors of COVID-19 related mortality after adjustment for age, sex and co-morbidities. A higher prevalence of ICS users in patients who were hospitalized for COVID-19 was found by Izquierdo *et al.*, but Choi *et al.* did not find any association between asthma medications and COVID-19 outcome.

The studies by Choi *et al.* and Izquierdo *et al.* are both unique and important because of their impressive sample size and innovative approach (big data analytics and artificial intelligence in the study by Izquierdo *et al.*). But have these large-scale studies provided us with definitive answers as to whether asthma, the severity of asthma and asthma medications affect COVID-19 susceptibility and severity? The higher incidence of COVID-19 among asthma patients reported in both studies is more or less consistent with other large (but smaller) studies.[8,18–20]. However, there are still some outstanding issues that may have influenced the results. First, the method by which asthma was diagnosed was very strict in the study by Choi *et al.* and rather vague in the study by Izquierdo *et al.*, possibly leading to under- or over diagnosis, respectively. This may also explain differences in proportions of patients using ICS between the two studies (23% vs 41%). Another bias factor may have been the methods by which COVID-19 was diagnosed. Choi *et al.* exclusively included PCR confirmed COVID-19 cases, while Izquierdo *et al.* also included patients with suspected COVID-19 based on clinical parameters. In addition, infection rates may also have been affected by testing policies or shielding advices, for example, if older patients,

patients with co-morbidities like asthma, or patients with more severe symptoms tested more frequently or better protected themselves. Thus, with such inaccuracies in case definition and variations in local conditions, it remains difficult to determine with certainty whether asthma patients are more susceptible to getting COVID-19 or not.

How about the risk of poor outcome or death from COVID-19? Severity and outcome of COVID-19 are highly dependent on age, as children experience less severe COVID-19 than elderly people.[22] Age is therefore an important confounder in the assessment of risk of contracting severe COVID-19. Izquierdo *et al.* showed that asthma patients without COVID-19 were younger and more likely to have eczema and rhinitis, while those with COVID-19 were older and more likely to have co-morbidities like hypertension and diabetes. These results could very well be confounded by age. Choi *et al.* solved this issue by adjusting for age and co-morbidities in a multivariate analysis, which would have been of additive value in the study by Izquierdo *et al.*

What do the two studies teach us about safety of asthma medication, in particular inhaled corticosteroids with respect to COVID-19 susceptibility and outcome? There is much debate about the risk-increasing or protective effects of asthma medication in the course of COVID-19 disease.[23] In such risk-assessment, asthma severity and phenotype are important confounders. Choi *et al.* found higher health-care related costs in patients using oral short-acting beta agonists (SABA). However, less than 4% of patients used oral SABA in the 2 months before COVID-19 diagnosis, suggesting that not oral SABA itself, but other factors, such as the type of patients that is prescribed oral SABA, played a role in these increased costs. Not surprisingly, regression analysis showed that asthma medications were not independently associated with poor outcome of COVID-19. Choi *et al.* also found a longer hospital stay in patients on Step 5 treatment (defined as  $\geq 90$  days oral corticosteroids in the previous year), but firm conclusions cannot be drawn as only 4 patients were on Step 5 treatment. Izquierdo *et al.* conclude that

ICS use is “safe”, because ICS users in their study were less frequently hospitalized for COVID-19 than non-ICS-users. This finding is however in contrast with a recent observational study in the UK showing an association between higher ICS doses and risk of COVID-19 related death in asthma patients.[13] The studies by Choi *et al.* and Izquierdo *et al.* did not take dosage of ICS into account, which may partly explain the discrepancy between the three studies regarding the influence of ICS use on COVID-19 outcome.

How about the use of asthma biologics and the risk of COVID-19? Currently available asthma biologics block pathways of type 2 inflammation. This type of inflammation, in particular allergic inflammation, has been suggested to have a protective effect through down-regulating the angiotensin converting enzyme (ACE)2 receptor used by the virus to enter host cells, or hypothetically, by counterbalancing the exaggerated anti-viral immune response observed in severe COVID-19 patients.[24,25] Izquierdo *et al.* analysed a large number of patients on biologic therapy (n=865), and found a relatively high incidence of COVID-19 of 2.3% in these patients, while this was 1.4% in the general asthma population. Only two out of 20 infected patients on asthma biologics (10%) were hospitalized. While these numbers are small, they are consistent with other reports suggesting that there is no increased risk of a poor outcome in asthma patients on biologics.[21,26]

In summary, these large-scale studies have confirmed previous findings about the risk for asthma patients to develop (severe) COVID-19. Asthma patients appear to be slightly more susceptible to contracting COVID-19, but severe disease progression does not seem to be related to medication use, including asthma biologics, but rather to older age and co-morbidities.

However, no definitive conclusions can be drawn yet as many factors can influence the reported incidences of (severe) COVID-19 (Figure 1). These potential bias factors have not been taken into account in the published studies so far and therefore many questions remain unanswered. Similar large-

scale, preferably multinational real-life studies with detailed information on asthma phenotype and medication usage in patients with a confirmed diagnosis of COVID-19 would be an ideal next step to further build on this new evidence.

Legends to figure

**Title Figure 1:** Factors that may influence reported incidences of (severe) COVID-19 in asthma patients

**Legend Figure 1:** The reported incidences of (severe) COVID-19 cases among asthma patients are not determined by patient-related factors alone. Also local factors and the applied methodologies can play an important role.

Abbreviations: ACE2, angiotensin converting enzyme 2; COVID-19, coronavirus disease 2019; ICS, inhaled corticosteroids; OCS, oral corticosteroids.



## References

1. WHO. Weekly epidemiologic update. <https://www.who.int/publications/m/item/weekly-epidemiological-update---1-december-2020>. Date last updated: December 1 2020. Date last accessed: December 3 2020.
2. Wu Z, McGoogan JM. Characteristics of and Important Lessons from the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323(13):1239–42.
3. Guan W-j, Liang W-h, Zhao Y, Liang H-r, Chen Z-s, Li Y-m, Liu X-q, Chen R-c, Tang C-l, Wang T, Ou C-q, Li L, Chen P-y, Sang L, Wang W, Li J-f, Li C-c, Ou L-m, Cheng B, Xiong S, Ni Z-y, Xiang J, Hu Y, Liu L, Shan H, Lei C-l, Peng Y-x, Wei L, Liu Y, Hu Y-h, Peng P, Wang J-m, Liu J-y, Chen Z, Li G, Zheng Z-j, Qiu S-q, Luo J, Ye C-j, Zhu S-y, Cheng L-l, Ye F, Li S-y, Zheng J-p, Zhang N-f, Zhong N-s, He J-x. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J.* 2020; 55: 2000547
4. Garg S, Kim L, Whitaker M, O'Halloran A, Cummings C, Holstein R, Prill M, Chai SJ, Kirley PD, Alden NB, Kawasaki B, Yousey-Hindes K, Niccolai L, Anderson EJ, Openo KP, Weigel A, Monroe M, Ryan P, Henderson J, Kom S, Como-Sabetti K, Lynfield R, Sosin D, Torres S, Muse A, Bennet N, Billing L, Sutton M, West N, Schaffner W, Talbot K, Aquino C, George A, Budd A, Brammer L, Langley G, Hall A, Fry A. Hospitalization Rates and Characteristics of Patients Hospitalized with Laboratory-Confirmed Coronavirus Disease 2019 — COVID-NET, 14 States, March 1–30, 2020. *Morb Mortal Wkly Report, US Dep Heal Hum Serv Dis Control Prev.* 2020;69(15):458–64.
5. Choi H-g, Wee JH, Kim SY, Kim J-h, Kim HI, Park J-y, Park J-y, Park S, Hwang YI, Jang SY, Jung K-s. Association between asthma and clinical mortality/morbidity in COVID-19 patients using clinical epidemiologic data from Korean Disease Control & Prevention. *Allergy.* 2020. Online ahead of print.
6. Zhu Z, Hasegawa K, Ma B, Fujiogi M, Camargo CA, Liang L. Association of asthma and its genetic predisposition with the risk of severe COVID-19. *J Allergy Clin Immunol.* 2020;146(2):327-329.e4.
7. Green I, Merzon E, Vinker S, Golan-cohen A, Magen E. COVID-19 Susceptibility in Bronchial Asthma. *J Allergy Clin Immunol Pract.* 2020. Online ahead of print.

8. Mahdavinia M, Foster KJ, Jauregui E, Moore D, Adnan D, Andy-Nweye AB, Khan S, Bishehsari F. Asthma prolongs intubation in COVID-19. *J Allergy Clin Immunol Pract*. 2020;8(7):2388–91.
9. Lieberman-Cribbin W, Rapp J, Alpert N, Tuminello S, Taioli E. The Impact of Asthma on Mortality in Patients With COVID-19. *Chest*. 2020;194:2019–20.
10. Avdeev S, Moiseev S, Brovko M, Yavorovskiy A, Umbetova K, Akulkina L, Tsareva N, Merzhoeva Z, Gainitdinova V. Low prevalence of bronchial asthma and chronic obstructive lung disease among intensive care unit patients with COVID-19. *Allergy*. 2020 Oct;75(10):2703-2704.
11. Yang JM, Koh HY, Moon SY, Yoo IK, Ha EK, You S, Kim SY, Yon DK, Lee SW. Allergic disorders and susceptibility to and severity of COVID-19: A nationwide cohort study. *J Allergy Clin Immunol*. 2020;146(4):790–8.
12. Beurnier A, Jutant E-M, Jevnikar M, Boucly A, Pichon J, Preda M, Frank M, Laurent J, Richard C, Monnet X, Duranteau J, Harrois A, Chaumais M-C, Bellin M-F, Noël N, Bulifon S, Jaïs X, Parent F, Seferian A, Savale L, Sitbon O, Montani D, Humbert M. Characteristics and outcomes of asthmatic patients with COVID-19 pneumonia who require hospitalisation. *Eur Respir J*. 2020. In press.
13. Schultze A, Walker AJ, MacKenna B, Morton CE, Bhaskaran K, Brown JP, Rentsch CT, Williamson E, Drysdale H, Croker R, Bacon S, Hulme W, Bates C, Curtis HJ, Mehrkar A, Evans D, Inglesby P, Cockburn J, McDonald HI, Tomlinson L, Mathjur R, Wing K, Wong AYS, Forbes H, Parry J, Hester F, Harper S, Evans SJQ, Quint J, Smeeth L, Douglas IJ, Goldacret B. Risk of COVID-19-related death among patients with chronic obstructive pulmonary disease or asthma prescribed inhaled corticosteroids: an observational cohort study using the OpenSAFELY platform. *Lancet Respir Med*. 2020 Nov;8(11):1106-1120.
14. Peters MC, Sajuthi S, Deford P, Christenson S, Rios CL, Montgomery MT, Woodruff PG, Mauger DT, Erzurum SC, Johansson MW, Denlinger LC, Jarjour NN, Castro M, Hastie AT, Moore W, Ortega HE, Bleecker ER, Wenzel SE, Israel E, Levy BD, Seibold MA, Fahy JV. COVID-19 Related Genes in Sputum Cells in Asthma: Relationship to Demographic Features and Corticosteroids. *Am J Respir Crit Care Med*. 2020 Jul 1;202(1):83-90.
15. Camiolo M, Gauthier M, Kaminski N, Anuradha R WS. Expression of SARS-CoV-2 receptor ACE2 and coincident host response signature varies by asthma inflammatory phenotype. *J Allergy Clin Immunol*. 2020 Aug;146(2):315-324.e7.

16. Choi YJ, Park J-Y, Lee HS, Suh J, Song JY, Byun MK, Cho JH, Kim HJ, Lee JH, Park JW, Park HJ. Effect of Asthma and Asthma Medication on the Prognosis of Patients with COVID-19. *Eur Respir J*. 2020. In press.
17. Izquierdo JL, Almonacid C, González Y, Del Rio-Bermúdez C, Ancochea J, Cárdenas R, Soriano JB. The Impact of COVID-19 on Patients with Asthma. *Eur Respir J*. 2020. In press.
18. Lovinsky-Desir S, Deshpande DR, De A, Murray L, Stingone JA, Chan A, Patel N, Rai N, DiMango E, Milner J, Kattan M.. Asthma among hospitalized patients with COVID-19 and related outcomes. *J Allergy Clin Immunol*. 2020 Nov;146(5):1027-1034.e4.
19. Cchiba KD, Patel GB, Vu THT, Chen MM, Guo, A, Kudlaty E, Mai Q, Yeh C, Muhammed LF, Harris KE, Bochner BS, Grammer LC, Greenberger PA, Kalhan R, Kuang FL, Saltoun CA, Schleimer RP, Stevens WW, Peters AT. Prevalence and characterization of asthma in hospitalized and non-hospitalized patients with COVID-19. *J Allergy Clin Immunol*. 2020 Aug;146(2):307-314.e4.
20. Wang L, Foer D, Bates WD, Boyce JA ZL. Risk factors for hospitalization, intensive care, and mortality among patients with asthma and COVID-19. *J Allergy Clin Immunol*. 2020 Oct; 146(4): 808–812.
21. Hanon S, Brusselle G, Deschamphelire M, Louis R, Michils A, Peché R, Pilette C, Rummens P, Schuermans D, Simonis H, Vandenplas O, Schleich F. Early View COVID-19 and biologics in severe asthma : data from the Belgian Severe Asthma Registry. *Eur Respir J*. 2020 Oct 22;2002857.
22. Götzinger F, Santiago-García B, Noguera-Julián A, Lanaspá M, Lancella L, Calò Carducci FI, Gabrovská N, Velizarova S, Prunk P, Osterman V, Krivec U, Lo Vecchio A, Shingadia D, Soriano-Arandes A, Melendo S, Lanari M, Pierantoni L, Wagner N, L’Huillier AG, Heiningner U, Ritz N, Bandi S, Krajcar N, Roglic S, Santos S, Christiaens C, Creuven M, Buonsenso D, Welch SB, Bogyi M, Brinkmann F, Tebruegge M. COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *Lancet Child Adolesc Heal*. 2020;4(9):653–61.
23. Halpin DMG, Singh D, Hadfield RM. Inhaled corticosteroids and COVID-19: A systematic review and clinical perspective. *Eur Respir J*. 2020;55(5).
24. Jackson DJ, Busse WW, Bacharier LB, Kattan M, O’Connor GT, Wood RA, Visness CM, Durham SR, Larson D, Esnault S, Ober C, Gergen PJ, Becker P, Togias A, Gem JE, Altman MC.. Association of

Respiratory Allergy, Asthma and Expression of the SARS-CoV-2 Receptor, ACE2. *J Allergy Clin Immunol.* 2020 Jul;146(1):203-206.e3.

25. Carli G, Cecchi L, Stebbing J, Parronchi P, Farsi A. Carli Allergy 2020 Is asthma protective against COVID-19. *Allergy.* 2020. Online ahead of print.
26. Heffler E, Detoraki A, Contoli M, Papi A, Paoletti G, Malipiero G, et al. COVID-19 in Severe Asthma Network in Italy (SANI) patients: Clinical features, impact of comorbidities and treatments. *Allergy.* 2020 Aug 1:10.1111/all.14532.




Figure 1