



Early View

Research letter

Leveraging aging models of pulmonary fibrosis; the efficacy of nintedanib in aging

Kosuke Kato, Yoon-Joo Shin, Sunny Palumbo, Ioannis Papageorgiou, Seongmin Hahn, Joseph D. Irish, Skye P. Rounseville, Robert T. Krafty, Lutz Wollin, Maor Sauler, Louise Hecker

Please cite this article as: Kato K, Shin Y-J, Palumbo S, *et al.* Leveraging aging models of pulmonary fibrosis; the efficacy of nintedanib in aging. *Eur Respir J* 2021; in press (<https://doi.org/10.1183/13993003.00759-2021>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Leveraging Aging Models of Pulmonary Fibrosis; The Efficacy of Nintedanib in Aging

Kosuke Kato¹, Yoon-Joo Shin¹, Sunny Palumbo², Ioannis Papageorgiou¹, Seongmin Hahn², Joseph D. Irish², Skye P. Rounseville², Robert T. Krafty³, Lutz Wollin⁴, Maor Sauler⁵, and Louise Hecker^{1, 6*}

Affiliations:

¹Division of Pulmonary, Allergy and Critical Care and Sleep Medicine, Department of Medicine, Emory University, Atlanta, GA, USA

²Division of Pulmonary, Allergy and Critical Care and Sleep Medicine, Department of Medicine, University of Arizona, Tucson, AZ, USA

³Department of Biostatistics and Bioinformatics, Rollins School of Public Health, Emory University, Atlanta, GA, USA

⁴Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach, Germany

⁵Department of Medicine, Yale School of Medicine, New Haven, CT, USA

⁶Atlanta VA Healthcare System, Atlanta, GA, USA

*To whom correspondence should be addressed:

Louise Hecker, Ph.D.

Telephone: (404) 727-3689

Department of Medicine, Division of Pulmonary, Allergy, and Critical Care and Sleep Medicine
Emory University, Whitehead Biomedical Research Building

615 Michael Street NE, Lab #135, Atlanta, GA 30322 USA

E-mail: louise.hecker@emory.edu

Take-Home Message:

Although nintedanib is overwhelmingly prescribed to elderly patients, this is the first study to demonstrate that aging does not impact the efficacy of nintedanib. This study sheds light on the utility of aged animal models in pulmonary fibrosis.

To the Editor:

Nintedanib is one of two FDA-approved treatments for idiopathic pulmonary fibrosis (IPF). The clinical efficacy of nintedanib for inhibiting the progression of lung fibrosis is well-established [1]. However, although nintedanib is overwhelmingly prescribed to elderly patients, the impact of aging on its efficacy is difficult to discern from clinical data due to the magnitude of confounding variables that exist among human subjects (genetics, gender, comorbidities, disease stage at the onset of treatment, etc.). A recent post-hoc meta analysis of five IPF clinical trials suggested that the effect of nintedanib in reducing the rate of forced vital capacity decline is consistent across patients with age (patients >75 vs. patients <75 years of age) [2]. However, it is important to note that the average age of IPF diagnosis is 66 years and the average patient ages in these cohorts were 78 (>75) vs. 64 (<75) years. Further, one could argue that patients in both cohorts represent the elderly population. This study highlights the complexity of evaluating the impact of aging on efficacy in a clinical setting. To date, all pre-clinical efficacy studies with nintedanib have been performed in young animals. We therefore sought to determine whether aging impacts the efficacy of nintedanib for inhibiting the development of lung fibrosis. Bleomycin-induced lung injury in young (2 month) and aged (18 month) mice was followed by treatment with nintedanib or vehicle from day 10-21 (**Fig. 1A**), using a previously described protocol [3]. We previously demonstrated in this injury model that the severity of lung fibrosis is identical in young and aged mice, in terms of the net increase in total lung collagen following injury [4]. Although some prior studies have reported seemingly contradictory results, indicating increased severity of fibrosis in aged mice [5, 6], this discrepancy could be attributed to increased baseline levels of collagen in aged mice and the methodology/analyses used for fibrosis assessment, as the net increase in collagen appear to be similar in both young and aged mice [5, 6]. In line with our previous findings, both young and aged vehicle-treated mice demonstrated

similar levels of fibrosis severity and a similar decline in lung function at 3 weeks post-injury (**Fig. 1B-D, G-H**). Also consistent with numerous prior reports [7, 8], we found that in young mice, nintedanib demonstrated efficacy for inhibiting the development of fibrosis (**Figs. 1B-G**) and led to improved lung function (**Fig. 1H**). Interestingly, nintedanib also significantly inhibited the development of lung fibrosis in aged mice, to a similar extent as young cohorts (**Figs. 1B-G**). Although nintedanib treatment resulted in lung functional improvement to a similar extent in both young (49%) and aged (57%) mice (**Fig. 1H**), results did not reach statistical significance in aged mice. Of note, there is less than 47% power to detect mean differences between the aged-vehicle and aged-nintedanib groups given the observed effect and sample sizes of aged mice; the trending *p-value* of 0.06 is displayed to provide a better understanding of the results. No significant differences in survival rate were observed between nintedanib- vs. vehicle-treated groups for both young (68% vs. 72%, respectively) and aged mice (83% vs. 76%, respectively) during this treatment period (day 10-21). Overall, these data indicate that aging does not impact the efficacy of nintedanib in terms of its ability to inhibit the development of de novo lung fibrosis.

Most IPF patients exhibit progressive lung fibrosis. Although this bleomycin model is not typically considered to be a model of “progressive fibrosis”, one could argue that the development of de novo fibrosis is a component of progressive fibrosis. This is the first study to suggest that age alone does not impact the efficacy of nintedanib for inhibiting the development of de novo lung fibrosis. This is reassuring from a clinical standpoint, given that nintedanib is predominantly administered to elderly patients [1]. It is tempting to speculate that the efficacy of nintedanib in aging (as described here) may have been a fortuitous feature that contributed to its success in clinical trials, particularly since numerous drug candidates demonstrating efficacy in young mice have failed in clinical trials with elderly patients. No studies have evaluated whether aging impacts

the efficacy of pirfenidone for inhibiting the development of fibrosis. However, if age-dependent differences in the efficacy of pirfenidone are identified, such findings could have clinical implications for the selection of therapeutics based on patient age.

It is well established that nintedanib inhibits progressive lung decline in IPF patients [1]. The prevailing understanding has been that nintedanib inhibits fibrosis progression. However, no studies have evaluated whether nintedanib can reverse established fibrosis. Such studies are not feasible in young mice, as young mice demonstrate a self-limited fibrotic response where fibrosis spontaneously resolves following peak injury [4, 9]. Conversely, aged mice exhibit a persistent fibrotic response, with little to no resolution of fibrosis at 4 months post-injury [4]. This aging model of persistent lung fibrosis offers a more clinically relevant testing protocol, where the therapeutic efficacy of agents can be evaluated for their ability to resolve age-dependent persistent fibrosis; no studies have previously evaluated the efficacy of nintedanib or reversing age-dependent established fibrosis. We therefore administered nintedanib or vehicle to aged mice daily from 3-6 weeks post-injury (when aged mice demonstrate established/persistent fibrosis) [4] (**Fig. 1I**). Like our previously published studies [4], we found that aged vehicle-treated mice exhibited persistent fibrosis during this time period (from 3-6 weeks) (**Fig. 1J**). However, nintedanib treatment failed to rapidly promote the resolution of age-dependent established lung fibrosis during the 3-week treatment period (**Fig. 1J-P**). Further, no significant survival benefit was observed between vehicle and nintedanib treated aged mice (73.9% vs. 78.2%, respectively) during this treatment period (3-6 weeks). It is possible that increased duration of nintedanib treatment may be required in order to definitively evaluate the efficacy of nintedanib for reversing age-dependent established fibrosis. Senescent fibroblasts secrete elevated levels of PDGF (a target of nintedanib) [10], and aged mice exhibit a persistent senescence response associated with non-resolving fibrosis

for up to 4 months post-injury [4]. To date, no therapeutic has demonstrated the ability to reverse age-associated established fibrosis – which may represent the holy grail for therapeutic strategies to more effectively treat and/or cure IPF.

Although the FDA-approval of 2 therapeutic agents represents a significant breakthrough by providing the first available treatment options for IPF patients, these therapies do not cure IPF or significantly improve the quality of life for IPF patients. Thus, the search for more effective therapies continues. One possible explanation for the limited therapies available is that age-dependent pathologic mechanisms remain largely unexploited in the drug development process, despite that aging has been strongly implicated in the pathogenesis of IPF [11, 12]. Numerous studies have demonstrated the presence of premature aging hallmarks, including the accumulation of senescent cells, in the lungs of IPF patients [4, 13]. However, senescence can be pro- or anti-fibrotic, depending on the context of age [4, 14]. Following injury in young mice, senescent cells undergo apoptosis, which promotes fibrosis resolution [4, 14]. Whereas in aged mice, senescence leads to the acquisition of an apoptosis-resistant myofibroblast phenotype, which promotes fibrosis persistence [4]. With the emergence of senolytics (senescence-targeting therapeutic agents) as a promising treatment strategy for IPF [15], age-relevant pre-clinical efficacy models will be critical to accurately evaluate their potential. Key opinion leaders recommend that aging models should be employed when evaluating age-related targets [16-18], as testing of senolytics in young animal models alone may not provide a fully comprehensive assessment of their potential. Overall, this study sheds light on the potential utility of aged animal models, which provide opportunities to identify/validate age-dependent pathologic mechanisms, address specific research questions, and/or employ more rigorous efficacy testing protocols for evaluating therapeutic agents. Incorporating aging models at the appropriate stages of research and/or therapeutic development

is likely to provide novel insight that would accelerate the successful translation of improved therapies for IPF, a disease that disproportionately afflicts the elderly population.

Support statement: This work was funded by Boehringer Ingelheim Pharma GmbH & Co. KG.


Author contributions: LH conceived of the project, provided funding, and supervised all studies. LH, KK, LW, and MS designed and/or performed analyses for experiments. All authors contributed intellectual input to the overall project, technical support for conducting studies, and/or manuscript preparation.

References:

1. Lancaster L, Crestani B, Hernandez P, Inoue Y, Wachtlin D, Loaiza L, Quaresma M, Stowasser S, Richeldi L: **Safety and survival data in patients with idiopathic pulmonary fibrosis treated with nintedanib: pooled data from six clinical trials.** *BMJ Open Respir Res* 2019, **6**(1):e000397.
2. Glaspole I, Bonella F, Bargagli E, Glassberg MK, Caro F, Stansen W, Quaresma M, Orsatti L, Bendstrup E: **Efficacy and safety of nintedanib in patients with idiopathic pulmonary fibrosis who are elderly or have comorbidities.** *Respir Res* 2021, **22**(1):125.
3. Wollin L, Maillet I, Quesniaux V, Holweg A, Ryffel B: **Antifibrotic and anti-inflammatory activity of the tyrosine kinase inhibitor nintedanib in experimental models of lung fibrosis.** *J Pharmacol Exp Ther* 2014, **349**(2):209-220.
4. Hecker L, Logsdon NJ, Kurundkar D, Kurundkar A, Bernard K, Hock T, Meldrum E, Sanders YY, Thannickal VJ: **Reversal of persistent fibrosis in aging by targeting Nox4-Nrf2 redox imbalance.** *Sci Transl Med* 2014, **6**(231):231ra247.
5. Redente EF, Jacobsen KM, Solomon JJ, Lara AR, Faubel S, Keith RC, Henson PM, Downey GP, Riches DW: **Age and sex dimorphisms contribute to the severity of bleomycin-induced lung injury and fibrosis.** *Am J Physiol Lung Cell Mol Physiol* 2011, **301**(4):L510-518.
6. Sueblinvong V, Neujahr DC, Mills ST, Roser-Page S, Ritzenthaler JD, Guidot D, Rojas M, Roman J: **Predisposition for disrepair in the aged lung.** *Am J Med Sci* 2012, **344**(1):41-51.
7. Wollin L, Distler JHW, Redente EF, Riches DWH, Stowasser S, Schlenker-Herceg R, Maher TM, Kolb M: **Potential of nintedanib in treatment of progressive fibrosing interstitial lung diseases.** *Eur Respir J* 2019, **54**(3).
8. Wollin L, Wex E, Pautsch A, Schnapp G, Hostettler KE, Stowasser S, Kolb M: **Mode of action of nintedanib in the treatment of idiopathic pulmonary fibrosis.** *Eur Respir J* 2015, **45**(5):1434-1445.
9. Izbicki G, Segel MJ, Christensen TG, Conner MW, Breuer R: **Time course of bleomycin-induced lung fibrosis.** *Int J Exp Pathol* 2002, **83**(3):111-119.
10. Demaria M, Ohtani N, Youssef SA, Rodier F, Toussaint W, Mitchell JR, Laberge RM, Vijg J, Van Steeg H, Dolle ME *et al*: **An essential role for senescent cells in optimal wound healing through secretion of PDGF-AA.** *Dev Cell* 2014, **31**(6):722-733.
11. Kato K, Hecker L: **NADPH oxidases: Pathophysiology and therapeutic potential in age-associated pulmonary fibrosis.** *Redox Biol* 2020:101541.
12. Hecker L: **Mechanisms and consequences of oxidative stress in lung disease: therapeutic implications for an aging populace.** *Am J Physiol Lung Cell Mol Physiol* 2018, **314**(4):L642-L653.
13. Schafer MJ, White TA, Iijima K, Haak AJ, Ligresti G, Atkinson EJ, Oberg AL, Birch J, Salmonowicz H, Zhu Y *et al*: **Cellular senescence mediates fibrotic pulmonary disease.** *Nat Commun* 2017, **8**:14532.
14. Krizhanovsky V, Yon M, Dickins RA, Hearn S, Simon J, Miething C, Yee H, Zender L, Lowe SW: **Senescence of activated stellate cells limits liver fibrosis.** *Cell* 2008, **134**(4):657-667.
15. Merkt W, Bueno M, Mora AL, Lagares D: **Senotherapeutics: Targeting senescence in idiopathic pulmonary fibrosis.** *Semin Cell Dev Biol* 2020, **101**:104-110.

16. Jenkins RG, Moore BB, Chambers RC, Eickelberg O, Konigshoff M, Kolb M, Laurent GJ, Nanthakumar CB, Oltman MA, Pardo A *et al*: **An Official American Thoracic Society Workshop Report: Use of Animal Models for the Preclinical Assessment of Potential Therapies for Pulmonary Fibrosis**. *Am J Respir Cell Mol Biol* 2017, **56**(5):667-679.
17. Yanagihara T, Chong SG, Vierhout M, Hirota JA, Ask K, Kolb M: **Current models of pulmonary fibrosis for future drug discovery efforts**. *Expert Opin Drug Discov* 2020, **15**(8):931-941.
18. Bonniaud P, Fabre A, Frossard N, Guignabert C, Inman M, Kuebler WM, Maes T, Shi W, Stampfli M, Uhlig S *et al*: **Optimising experimental research in respiratory diseases: an ERS statement**. *Eur Respir J* 2018, **51**(5).

FIGURE LEGENDS









I**J****K****L****M****N****O****P**

Figure 1: Nintedanib inhibits pulmonary fibrosis development in aged mice but does not reverse age-dependent established fibrosis. (A-H) C57BL/6 young (2 month) and aged (18 month) male mice received intratracheal instillation of bleomycin (0.02875 U/mouse) were treated with nintedanib (60 mg/kg dissolved in de-ionized water) or vehicle (de-ionized water) via oral gavage daily from days 10-21. All mice were evaluated at 21 days post-injury. (A) Schematic diagram illustrating treatment protocol. (B) Lung tissue was assessed by H&E staining for histopathology and Masson's trichrome staining for collagen. (C) Total lung collagen was determined by quantitative Sircol assay. Data are presented as % change at 3 weeks post-injury relative to respective uninjured control group. (D) Total lung hydroxyproline was analyzed by quantitative hydroxyproline assay. Data are presented as % change at 3 weeks post-injury relative to respective uninjured control group. (E-F) Collagen-1 α 1 expression was assessed by Western blot (E) and (F) densitometric analysis. (G) Lung weight at day 0 and day 21 (pre- and post-injury/treatment, respectively). (H) Static lung compliance was determined by FlexiVent. (I-P) Aged (18 month) C57BL/6 female mice were administered bleomycin (0.03125 U/mouse) via intratracheal instillation. Starting at 3 weeks post-injury (when aged mice demonstrate established/persistent fibrosis), nintedanib (60 mg/kg) or vehicle was administered daily from 3-6 weeks via oral gavage. (I) Schematic diagram illustrating treatment protocol. (J) Lung tissue was assessed at 6 weeks post-injury by H&E staining for histopathology and Masson's trichrome staining for collagen. (K) Total lung collagen at 6 weeks post-injury was determined by quantitative Sircol assay. Data are presented as % change at 6 weeks relative to 3 weeks post-injury. (L) Total lung hydroxyproline was analyzed by quantitative hydroxyproline assay. Data are presented as % change at 6 weeks relative to 3 weeks post-injury. (M-N) Collagen-1 α 1 expression was assessed by Western blot (M) and densitometric analysis (N). (O) Lung weight at

6 weeks post-injury is shown. (**P**) Ashcroft scoring was evaluated by HistoWiz (Brooklyn, NY). All graphed data are shown as a dot plot \pm SEM. Student's two-tailed t-test was used for comparisons between two groups and one-way ANOVA was used to compare three or more groups; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, and *ns* = not significant.