



Early View

Correspondence

The curious incident of Long-Covid symptoms, from imaginary condition to recognised syndrome. A “small victory”

Valentina E. Di Mattei, Gaia Perego, Francesca Milano, Thomas Eric Hill, Sergio Alfonso Harari

Please cite this article as: Di Mattei VE, Perego G, Milano F, *et al.* The curious incident of Long-Covid symptoms, from imaginary condition to recognised syndrome. A “small victory”. *Eur Respir J* 2022; in press (<https://doi.org/10.1183/13993003.00653-2022>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©The authors 2022. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

The curious incident of Long-Covid symptoms, from imaginary condition to recognized syndrome. A “small victory”.

Valentina E. Di Mattei ^{1,2}, Gaia Perego ³, Francesca Milano ², Thomas Eric Hill ⁴, & Sergio Alfonso Harari ^{5,6}

¹ School of Psychology, Vita-Salute San Raffaele University, Milan, Italy

² Clinical and Health Psychology Unit, IRCCS San Raffaele Scientific Institute, Milan, Italy

³ Department of Psychology, University of Milano-Bicocca, Milan, Italy

⁴ School of Medicine, Vita-Salute San Raffaele University, Milan, Italy

⁵ Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy

⁶ Department of Medicine, IRCCS Ospedale San Giuseppe MultiMedica, Milan, Italy

Over the last two years, the scientific community has acquired a better understanding about COVID-19 as an acute disease, but there is still much to learn about COVID-19, especially when considering its chronic aspect. In the early stages of the pandemic, healthcare workers focused on COVID-19 acute symptoms, often overlooking its long-term and chronic implications. In October 2021, the World Health Organization recognized these long term complications as Long-COVID syndrome, which probably affected more than 62% of patients [1]. Patients with Long-COVID syndrome experienced the fear of living forever with those symptoms, arising from the lack of medical information on this issue [2], combined with the feeling of being ignored and disbelieved. Several patients felt they were neglected the appropriate health care, as they perceived that healthcare workers disregarded their symptoms and judged them as unreal, often considering them secondary to psychiatric issues [3].

Beyond the healthcare field, patients found little support or understanding even in their social networks, with a post-COVID syndrome resulting in a deterioration in their ability to perform daily activities, family relationships and working capacity [1]. For instance, family members were frequently unable to provide enough emotional support or accept the patient’s increased need for rest due to symptoms [2]. In the workplace, people were often unable to perform their duties as well as usual, thus worked shorter shifts due to fatigue and a fear of making mistakes caused by brain fog [3]. This led to isolation, anxiety, self-doubt, changed professional identity [3,4], and to a dangerous feeling of loneliness.

Originally these “invisible symptoms” seemed to be noticed only in online communities which reported symptoms primarily through anecdotes. Patients’ illness narratives drew public and scientific/medical attention on Long-COVID syndrome, leading to the recognition of long-covid symptoms as a physical illness rather than an imaginary condition [4]. This belief and attention by clinicians and relatives provided patients with a sense of relief and recognition, which can be defined as “a small victory” [3]. Long-COVID syndrome is consistently regarded as the first disease collectively constructed and identified by patients [5].

The impression of being understood and believed is particularly important for patients, as illness makes people feel vulnerable and in need of assistance. Such debilitating conditions put patients in a position of dependence on healthcare workers, fostering the activation of a set of defense mechanisms, unconscious strategies used by our mind to protect the ego from stress, fear, or trauma. One of the most used during illness is the so called “regression”, in which an individual, regardless of age, copes with stressful situations by retreating to earlier stages of development and abandoning forms of gratification appropriate to them. For example, an adult who is hospitalized after being diagnosed with a medical issue may deal with their circumstances by curling up on the bed in the fetal position and clutching a stuffed animal dog. A reversible regression is an important adaptive tool during illness. In the short term, it may have a positive function by helping patients to accept the conditions imposed by illness, but it may hinder their independence and empowerment in the long term. Indeed, the patient-clinician relationship could easily activate reversible mechanisms, which are frequently expressed in terms of dependent child - caring/rewarding parent [6]. Consistently, healthcare workers’ recognition and acceptance of symptoms are critical in this field.

In addition, the literature shows that the attachment styles of patients and healthcare workers influences their relationship [7]. Attachment styles are different ways of interacting and behaving in intimate relationships, which are heavily influenced by self-worth and interpersonal trust. Theoretically, the degree of attachment security in adults is directly related to how well they bonded to others as children. Interestingly, patients’ attachment style influences their health-seeking behaviors as well as their acceptance of being helped and cared for [8]. This is also relevant for healthcare workers as attachment style influences their ability to communicate well with the patient and not only get a more complete understanding of the disease but also build a solid patient-doctor relationship [7]. However, the effort of being empathetic and open to the patient’s experience may be hampered by the medium and

high burnout levels measured among physicians and nurses, especially as was noticed during the pandemic [9].

In conclusion, given the importance of the doctor-patient relationship in Long-COVID management, we advocate for a “relationship-based care” [10]: general practitioners must offer consideration, empathy, and support to Long-COVID patients and their families, to assist them in overcoming the disease.

Moreover, several studies recommend face-to-face patient assessment and the adoption of a multidisciplinary approach at each stage of the disease [2;3]. This would involve the integration and training of mental health professionals and occupational health teams, which could help reduce burnout among physicians and nurses while also allowing for continuity of care and personalized rehabilitation.

Conflict of interest

The authors declare they have no conflict of interest.

Bibliography

1. Harari S, Ripamonti L, Marveggio P, Mannucci PM. Long COVID: A patient perspective. *European journal of internal medicine* 2022; 95, 104-105.
2. Kingstone T, Taylor AK, O'Donnell CA, Atherton H, Blane DN, Chew-Graham CA. Finding the “right” GP: a qualitative study of the experiences of people with long-COVID. *BJGP Open* 2020; 4(5), bjgpopen20X101143. <https://doi.org/10.3399/bjgpopen20X101143>
3. Callan C, Ladds E, Husain L, Pattinson K, Greenhalgh T. “I can't cope with multiple inputs”: Qualitative study of the lived experience of ‘brain fog’ after Covid-19. *MedRxiv: The Preprint Server for Health Sciences* 2021 <https://doi.org/https://doi.org/10.1101/2021.08.07.21261740>
4. Baz S, Fang C, Carpentieri JD, Sheard L. Understanding the lived experience of Long Covid: A rapid literature review. (Convalescence Long-COVID Study Briefing Reports). University College London 2021.
5. Callard F, Perego E. How and why patients made Long Covid. *Social Science and Medicine*. 2021; 268. doi: 10.1016/j.socscimed.2020.113426.
6. Kernberg OF. Severe personality disorders: Psychotherapeutic strategies. Yale University Press; 1984.

7. Ardenghi S, Rampoldi G, Bani M, Strepparava MG. Attachment styles as predictors of self-reported empathy in medical students during pre-clinical years. *Patient Education and Counseling* 2020; 103(5), 965-970.
8. Hooper LM, Tomek S, Newman CR. Using attachment theory in medical settings: implications for primary care physicians. *Journal of Mental Health* 2012; 21(1), 23-37.
9. Di Mattei VE, Perego G, Milano F, Mazzetti M, Taranto P, Di Pierro R, De Panfilis C, Madeddu F, Preti E. The “healthcare workers’ wellbeing (Benessere operatori)” project: A picture of the mental health conditions of italian healthcare workers during the first wave of the covid-19 pandemic. *International Journal of Environmental Research and Public Health* 2021; 18(10). <https://doi.org/10.3390/ijerph18105267>
10. Atherton H, Briggs T, Chew-Graham C. Long COVID and the importance of the doctor–patient relationship. *British Journal of General Practice* 2021; 71(703), 54-55.