

Clinical efficacy of low-dose inhaled budesonide once or twice daily in children with mild asthma not previously treated with steroids

G. Jónasson*, K-H. Carlsen[†], P. Blomqvist[‡]

Clinical efficacy of low-dose inhaled budesonide once or twice daily in children with mild asthma not previously treated with steroids. G. Jónasson, K-H. Carlsen, P. Blomqvist. ©ERS Journals Ltd 1998.

ABSTRACT: The aim of the present study was to examine the efficacy of low-dose inhaled budesonide (BUD) administered via Turbuhaler® once or twice daily on symptoms, lung function and bronchial hyperreactivity in children with mild asthma.

One hundred and sixty-three children (mean age 9.9 yrs, 56 females/107 males) with mild asthma (forced expiratory volume in one second (FEV₁) 103% of predicted, morning peak expiratory flow (PEF) 87% pred, reversibility in FEV₁ 3%, fall in FEV₁ after exercise 10.4% from pre-exercise value) and not previously treated with inhaled steroids, were included in a double-blind, randomized, parallel-group study. After a two-week run-in period, the children received inhaled BUD 100 µg or 200 µg once daily in the morning, 100 µg twice daily or placebo for 12 weeks. Exercise and methacholine challenges were performed before and at the end of treatment.

After 12 weeks of therapy, the fall in FEV₁ after an exercise test was significantly less in all three BUD groups (4.3–5.1%) than in the placebo group (8.6%). Bronchial hyperreactivity to methacholine with the provocative dose causing a 20% fall in FEV₁ decreased significantly in the BUD 100 µg twice-daily group compared with placebo (ratio at the end of treatment 156%). Changes in baseline lung function (FEV₁ and PEF) were less marked than changes in bronchial responsiveness.

In conclusion, low doses of inhaled budesonide, given once or twice daily, provided protection against exercise-induced bronchoconstriction in children with mild asthma and near normal lung function.

Eur Respir J 1998; 12: 1099–1104.

*Paediatric Section for Allergy and Pulmonology, Ullevål Hospital, Oslo, Norway. [†]Voksentoppen, Centre of Allergy and Asthma in Children, Oslo, Norway. [‡]Clinical Research and Development, Astra Draco, Lund, Sweden.

Correspondence: G. Jónasson
Paediatric Section for Allergy and Pulmonology
Ullevål Hospital
N-0407 Oslo
Norway
Fax: 47 22118663

Keywords: Asthma
bronchial hyperreactivity
budesonide
children
exercise-induced asthma
inhaled corticosteroids

Received: February 6 1998
Accepted after revision June 28 1998

The goal of asthma therapy in children is to allow them to be involved in normal everyday activities, including full participation in exercise and sports. There should not be excessive absence from school and children should be free from symptoms day and night. Lung function should be normalized, with low diurnal variation in peak expiratory flow (PEF) [1]. At present, glucocorticosteroids are the most effective anti-inflammatory drugs in the treatment of asthma and inhaled steroids are now recommended as first-line therapy in children with severe or moderate asthma. Treatment recommendations for children with mild, persistent asthma include either a low dose of the inhaled steroid cromolyn or nedocromil [2]. Long-term studies in children with asthma have shown that early intervention may prevent the development of irreversible airway obstruction [3, 4]. In the past few years, inhaled steroids have been introduced in the treatment of mild asthma with a recommended dose of 50–200 µg twice daily [5]. However, there is evidence to suggest that the lowest possible dose should be used to control asthma symptoms, since systemic effects of glucocorticosteroids are dose-dependent [6–8]. Previous studies have shown that low doses (100–200 µg) of inhaled steroids may be effective in controlling asthma symptoms in children, without causing apparent systemic side-effects [9–11].

In the present paper, a randomized, double-blind, placebo-controlled trial of low doses of inhaled budesonide (BUD) given once or twice daily in children with mild asthma is described.

Patients and methods

Study design

The study was a double-blind, single-centre trial. A two-week open run-in period, which served as a baseline and during which the children were introduced to the Turbuhaler® inhaler (Astra Draco, Lund Sweden), a peak flow meter and completing a diary, was followed by a 12-week double-blind treatment period with five visits to the clinic (at enrolment, at randomization and at four-week intervals). Patients were randomized into four parallel groups in balanced blocks. At randomization, each patient received two Turbuhaler inhalers, one to be used in the morning and the other in the evening. Group I patients were given BUD 100 µg once daily in the morning and placebo in the evening. Group II received BUD 200 µg once daily in the morning and placebo in the evening.

Group III received BUD 100 µg twice daily. The remaining group, group IV, received placebo twice daily.

The patients kept daily records of peak flow values, symptoms, use of β₂-agonists and intake of study medication. Clinical assessments and lung function tests (flow-volume curve and plethysmography) were performed at four-week intervals. Exercise tests were performed at enrolment and at the end of the 12-week treatment period. Measurement of the provocative dose of methacholine causing a 20% fall in the forced expiratory volume in one second (PD₂₀) was performed at randomization and after 12 weeks of treatment. After completing the baseline lung function tests and the exercise test at enrolment, a reversibility test to inhaled terbutaline was performed. Pulmonary function testing was performed at approximately the same time of the day for each patient. The exercise test and the methacholine provocation were performed on separate days.

The study was approved by the Regional Medical Ethics Committee in Oslo.

Patients

A total of 163 children with mild asthma were included in the study (107 male and 56 female, aged 7–16 yrs). A total of 166 patients were enrolled in the run-in period of the study and three discontinued prior to randomization. Two of these were excluded because their asthma deteriorated and one patient because of noncompliance. Patients were selected from the outpatient clinic at the Section of Allergy and Pulmonology, Dept of Paediatrics, Ullevål University Hospital in Oslo.

Inclusion criteria were the diagnosis of asthma, based on the definition in the International Consensus report [12] and in the Nordic Consensus report [13], with three previous obstructive episodes or one previous obstructive episode with atopy. At least one of these episodes had to have occurred within the year prior to the first visit. The patients did not use inhaled steroids within 2 months, or cromoglycate and/or nedocromil within 4 weeks of entry. No patient had any lower respiratory tract infection or exacerbation of asthma requiring an emergency room visit and/or hospitalization in the 4 weeks prior to entry.

Methods

BUD (daily dose 100 µg or 200 µg) or placebo was inhaled from Turbuhaler in the morning and evening during the 12-week treatment period. The patients were instructed to rinse their mouths out with water after each use. All patients were supplied with terbutaline (Bricanyl Turbuhaler® 250 µg), to be used as needed during the study. Children were required to demonstrate that they were competent in using Turbuhaler before enrolment, after the run-in period, and thereafter at four-week intervals during the study. A Vitalograph® peak flow meter (Vitalograph, Ennis, Ireland) was dispensed at enrolment. Patients measured their PEF every morning and evening throughout the entire study period. The highest PEF value from three exhalations was recorded on each occasion. At each clinic visit, staff checked that the children could use the PEF meter correctly. Patients were instructed to fill in a diary

twice daily, recording intake of study medication as well as rescue medication, PEF values, and symptom score. Asthma symptoms were scored on a four-point scale, which ranged from 0 (no symptoms) to 3 (severe symptoms). The diary sheet was collected at every visit.

Exercise-induced bronchoconstriction. This was determined at enrolment and at the end of the 12-week treatment period by the use of a motor-driven treadmill, where the children were instructed to run for 6 min with a submaximal exercise load [14]. The inclination of the treadmill was 5.5% and the speed was adjusted to a submaximal load to achieve a steady-state cardiac frequency of 170–180 beats·min⁻¹. The cardiac frequency was recorded electronically (Sport-Tester® PE 3000 with memory function: Polar Electro KY, Kempele, Finland). The submaximal run on the treadmill was performed at the same speed (exercise load) on both test occasions (at randomization and after 12 weeks of treatment) for each individual child. Forced expiratory volume in one second (FEV₁) was measured before running, immediately after, and 3, 6, 10 and 15 min after running. The maximum percentage fall in FEV₁ after the exercise test was calculated by:

$$\frac{(\text{Pre-exercise FEV}_1 - \text{Minimum post-exercise FEV}_1) \times 100}{\text{Pre-exercise FEV}_1}$$

Bronchial hyperreactivity. This was measured at randomization and at the end of the 12-week treatment period by a methacholine chloride inhalation test for the determination of PD₂₀, [15]. Methacholine was inhaled in doubling doses from 0.128 µmol up to a maximum dose of 16.32 µmol or until FEV₁ had fallen ≥20% compared with FEV₁ after an initial saline (0.9%) inhalation. The bronchial aerosol provocation system (APS) (Jaeger, Würzburg, Germany) with the Sandoz nebulizer (Jaeger) were used for this procedure. The Sandoz nebulizer was calibrated to nebulize 5 µL per nebulization and was triggered by inspiration on tidal ventilation. Each nebulization lasted for 0.5s. Lung function was measured 1 min after each inhaled dose. PD₂₀ was determined by linear interpolation on a semilogarithmic scale [15, 16].

If the patient had inhaled a β₂-agonist within 8 h prior to the lung function tests, the challenge was postponed until the next day.

Lung function. This was measured at every visit using maximal forced expiratory flow-volume curves and whole-body plethysmography (Masterlab Body; Jaeger) under body temperature, barometric pressure, saturated with water vapour (BTPS) conditions. Predicted lung function values were calculated using reference values from ZAPLETAL *et al.* [17]. At least three independent flow-volume curves were obtained on each visit and the curve with the highest FEV₁ was chosen. Values for the forced expiratory flow at 25% of the forced vital capacity (FEF_{25%}), FEF_{50%} and FEF_{75%} were also obtained from selected flow-volume curves. Calibrations were carried out daily on all lung function measuring equipment used in the study. Reversibility was tested at the entry visit, 15 min after administration of 500 µg of inhaled terbutaline from Turbuhaler.

Adverse events. Information was obtained at each visit by asking the patients or guardian "Have you/your child had any health problems or symptoms not usually associated with your/his/her asthma since the last visit?" The onset, severity and outcome of each adverse event were recorded.

Sample size

It was assumed that the standard deviation of a change in morning PEF was 30 L·min⁻¹ and with 40 patients per treatment group there would have been an 80% chance of detecting a difference of 20 L·min⁻¹ between two treatment effects when using a two-sided t-test with a significance level of $p < 0.05$.

Statistical evaluation

The change from baseline to the end of treatment was subjected to analysis of variance (ANOVA) with treatment as a factor and baseline value as a covariate. The ANOVA was used to obtain estimates of the pairwise differences between treatment effects and the standard deviation of each pairwise difference between treatment effects. An additive model was used when analysing diary variables, lung function variables and the maximum fall in FEV₁ after the exercise test, whereas a multiplicative model was used when analysing plethysmography variables and PD₂₀. The efficacy of BUD 100 µg once daily and placebo was compared using a two-sided t-test at a $p < 0.05$ significance level. The effect of BUD 200 µg once daily and BUD 100 µg once daily was compared in the same manner. The statistical analysis was based on the All Patients Treated approach; thus, all patients who had received at least one dose of study medication were included. Missing values were handled by applying the Last Value Extended principle. For diary variables, this was accomplished by extending the period means.

Results

The 163 randomized patients had a mean age of 9.9 yrs, a mean weight of 37.7 kg and a mean height of 143.4 cm

Table 1. – Patient characteristics at baseline for each treatment group

Characteristics	BUD 100 <i>b.i.d.</i>	BUD 200 <i>o.d.</i>	BUD 100 <i>o.d.</i>	Placebo
Number of patients	40	42	41	40
Males/Females	26/14	23/19	31/10	27/13
Age yrs	10.2	10.0	9.8	9.6
Height cm	145.4	144.9	142.1	141.2
Atopics n	26	25	31	28
FEV ₁ L	2.34	2.21	2.15	2.08
FEV ₁ % pred	105 (14)	101 (11)	103 (12)	102 (14)
Reversibility in FEV ₁ %	2.6 (5.1)	1.9 (7.2)	4.2 (4.7)	3.3 (5.4)
PEF morning L·min ⁻¹	260 (86)	272 (81)	245 (71)	237 (63)
PEF morning % pred	86 (17)	91 (16)	86 (14)	85 (14)
Methacholine PD ₂₀ µmol	3.67	4.14	3.74	3.02
Maximum fall in FEV ₁ after exercise test %	8.4	11	12.3	9.6

Results are given as mean values with SD in parentheses. BUD: budesonide; FEV₁: forced expiratory value in one second; PEF: peak expiratory flow; PD₂₀: provocative dose causing a 20% fall in FEV₁.

(table 1). The mean FEV₁ was 103% predicted and the mean reversibility in FEV₁ was 3%. Atopy was diagnosed in 67.5% of the children. Three patients were withdrawn from the study after randomization (all from the BUD 100 µg once-daily group), two of these because their asthma deteriorated and one due to noncompliance. At entry, the treatment groups were comparable regarding the number of patients, mean age, mean height and number of atopic patients. The proportion of females and males varied between the groups, as did baseline mean values for PEF and FEV₁, with the placebo group having the lowest mean values (table 1).

Exercise-induced bronchoconstriction

The mean maximum fall in FEV₁ (% fall from pre-exercise value) after the exercise test at baseline is presented in table 1. After 12 weeks of treatment, the corresponding values were: 4.3% for the BUD 100 µg twice-daily group, 5.1% for the BUD 200 µg once-daily group, 5.0% for the BUD 100 µg once daily group and 8.6% for the placebo group (fig. 1). All pairwise comparisons of active treatment *versus* placebo revealed statistically significant differences, with no differences between the active treatments (table 2).

Methacholine hyperreactivity

The mean percentage increase in PD₂₀ (µmol) from baseline to end of treatment was 110% in the BUD 100 µg twice-daily group, 40% in the BUD 200 µg once-daily group, 60% in the BUD 100 µg once-daily group, and 40% in the placebo group.

Pairwise comparison of active treatments *versus* placebo revealed a statistically significant difference between BUD 100 µg twice daily and placebo, with an estimated treatment effect ratio of 156% in favour of the BUD group (table 3).

Lung function

Baseline morning PEF values varied considerably among the four treatment groups (table 1). However, changes




Fig. 1. – Comparison between different treatment effects on maximum fall in the forced expiratory volume in one second (FEV₁). B: budesonide after exercise test (% fall from pre-exercise value) from baseline to week 12. ■: Baseline; □: week 12.

Table 2. – Comparison between different treatment effects, on maximum fall in forced expiratory volume in one second (FEV₁) after exercise test (% fall from pre exercise value) after 12 weeks of therapy

Treatment group	Estimated difference in fall FEV ₁ %	CI	p-value
BUD 100 <i>b.i.d.</i> /Placebo	-4.19	-7.6–-0.8	0.01
BUD 100 <i>o.d.</i> /Placebo	-4.69	-8.1–-1.3	<0.01
BUD 200 <i>o.d.</i> /Placebo	-4.15	-7.5–-0.8	0.015
BUD 200 <i>o.d.</i> /BUD 100 <i>b.i.d.</i>	-0.05	-3.3–3	NS
BUD 200 <i>o.d.</i> /BUD 100 <i>o.d.</i>	0.54	-2.8–3.9	NS

CI: 95% confidence interval; BUD: budesonide; NS: nonsignificant.

from baseline did not differ between the groups throughout the treatment period. A pairwise comparison between the different treatment groups showed no significant differences (table 4). An analysis of change in PEF (% pred) gave consistent results as compared with the change in actual PEF values.

A statistically significant difference in FEV₁ was only found between BUD 100 µg twice daily and placebo (0.1 L, *p*=0.015). At the end of the treatment period, no statistically significant difference was found between the treatment groups in FEF₂₅%, whereas there was a significant difference in both FEF₅₀% (0.3 L, *p*=0.001) and FEF₇₅% (0.1 L, *p*=0.039) between BUD 100 µg twice daily and placebo, and also between BUD 100 µg twice daily and BUD 200 µg once daily (FEF₅₀%; 0.3 L, *p*=0.002; FEF₇₅%; 0.1 L, *p*=0.04) in favour of the BUD 100 µg twice-daily group. Moreover, FEV₁ (% pred) showed a difference between BUD 100 µg twice daily and placebo (5.2%, *p*=0.008) and between BUD 100 µg twice daily and BUD 200 µg once daily (4.1%, *p*=0.035) in favour of the BUD 100 µg twice-daily group.

Resistance of the respiratory system (*R*_{rs}) and airway conductance (*G*_{aw}) differed significantly between BUD 100 µg twice daily and placebo (estimated ratio *R*_{rs} 89.3%, *p*=0.03; estimated ratio *G*_{aw} 115.2%, *p*=0.006). In a pairwise comparison between the groups, no other significant differences for *G*_{aw} and *R*_{rs} were found.

Symptom scores

Mean values for asthma symptoms were low throughout the study for all treatment groups. The mean symptom scores were already low at baseline, with a daytime mean symptom score of 0.39 and a night-time mean symptom score of 0.16.

Table 3. – Provocative dose of methacholine (µmol) causing a 20% fall in the forced expiratory volume in one second: estimated ratio between treatment effects after 12 weeks of therapy

Treatment group	Estimated ratio %	CI	p-value
BUD 100 <i>b.i.d.</i> /Placebo	156	101.5–239.1	0.04
BUD 100 <i>o.d.</i> /Placebo	121	78.3–186.7	NS
BUD 200 <i>o.d.</i> /Placebo	107	70.0–163.5	NS
BUD 100 <i>b.i.d.</i> /BUD 200 <i>o.d.</i>	146	95.4–222.2	NS
BUD 200 <i>o.d.</i> /BUD 100 <i>o.d.</i>	89	57.7–135.9	NS

CI: 95% confidence interval; BUD: budesonide; NS: nonsignificant.

Table 4. – Comparison between different treatment effects on change in peak expiratory flow after 12 weeks of therapy

Treatment group	Estimated difference L·min ⁻¹	95% CI	p-value
BUD 100 <i>b.i.d.</i> /Placebo	5.8	-8.1–19.7	NS
BUD 100 <i>o.d.</i> /Placebo	3.0	-10.8–16.8	NS
BUD 200 <i>o.d.</i> /Placebo	2.9	-11.0–16.7	NS
BUD 200 <i>o.d.</i> /BUD 100 <i>b.i.d.</i>	2.9	-10.8–16.6	NS
BUD 200 <i>o.d.</i> /BUD 100 <i>o.d.</i>	-0.1	-13.9–13.6	NS

Adjusted means from the analysis of variance and 95% confidence interval (CI). BUD: budesonide; NS: nonsignificant.

Statistically significant differences in symptoms during the night were found between BUD 200 µg once daily and placebo (-0.11, *p*=0.047), as well as between BUD 200 µg once daily and BUD 100 µg once daily (-0.11, *p*=0.040). This applied to both the mean symptom score and the frequency of symptom-free nights.

At baseline, the children used a mean number of 0.49 β₂-agonist inhalations during the day and 0.11 inhalations during the night. There were no significant differences in mean change from baseline between the treatment groups.

Adverse events

The adverse event profile observed in the three active treatment groups was similar to that reported for the placebo group. The most common adverse events were respiratory infection, coughing, and headache. One adverse event was regarded as serious: one patient receiving placebo experienced pharyngitis and high fever and was admitted to hospital for 3 days. A causal relationship to the study drug was judged unlikely. None of the patients discontinued the study because of adverse events.

Discussion

The present study in children with asthma shows that a low dose of BUD, administered once or twice daily for 12 weeks, protects against an exercise-induced reduction in lung function.

The once-daily morning regimens were shown to be as effective as the twice-daily dosing regimen with regard to the protective effect against exercise-induced asthma.

The majority of the patients selected for the present study had mild asthma, mild being defined as low-grade symptoms that did not interfere with sleep and lifestyle, or episodes of cough and wheeze occurring less than once a month [1]. Patients were selected on the basis of known asthma symptoms in the past year and not because of previously documented bronchial hyperresponsiveness. The fact, that mean reversibility was only 3% after inhalation of a bronchodilator and that FEV₁ at baseline showed a mean value of >100% pred indicate that the patients had near-normal lung function at baseline with little room for improvement. The mean bronchodilator use at baseline was 0.6 inhalations·24 h⁻¹. This relatively high use in this group of patients with mild disease can partly be explained by the fact that many of the children were using a bronchodilator routinely before exercise.

PEF, which is often used when evaluating the effect of inhaled steroid treatment, was shown to be a less sensitive measure than an exercise test in detecting treatment effects in the present study in children with near normal lung function and mild symptoms. The results of the peak flow measurements failed to demonstrate any statistically significant difference between the active treatments and placebo.

In a study on children with moderate to severe asthma (PEF 55% pred), PEDERSEN and HANSEN [11] found that an exercise test is a more sensitive parameter than PEF and FEV₁ in detecting differences between various doses of BUD. Even low doses of BUD, approximately 100 µg·day⁻¹ showed marked effects. This also proved to be true in this study in patients with mild asthma, where all active treatments provided better protection against exercise-induced bronchoconstriction than did placebo.

The results are important and useful for future studies, since protection against exercise-induced asthma may be used as a sensitive measure of the efficacy of inhaled steroids, which may reveal differences between different treatments that are not apparent when standard lung function tests are used. The result of the PEF data obtained in this study is supported by a study by UWYED *et al.* [18], where it was concluded that PEF monitoring is not sensitive enough to register meaningful changes in children and young adults with mild asthma (FEV₁ 82% pred and PEF 87% pred).

WAALKENS *et al.* [19] compared the effects of treatment with budesonide 200 µg twice daily and terbutaline with the effects of terbutaline alone in a placebo-controlled, double-blind study in 27 children with mild asthma (FEV₁ 93% pred) over 8 weeks. They found that mean FEV₁ did not change in either group and that there was no significant change in morning PEF, although they found a significant difference in favour of BUD for evening and nocturnal PEF. Airway hyperresponsiveness (provocative concentration of histamine causing a 20% fall in FEV₁ (PD₂₀)) decreased in the BUD/terbutaline-treated patients compared with the placebo/terbutaline group.

In the present study, the methacholine PD₂₀ was significantly larger in the children receiving BUD 100 µg twice daily than in those receiving placebo. The differences in PD₂₀ in the two other treatment groups were not statistically significant, compared with placebo, although there were numerical increases in all treatment groups. VAN ESSEN-ZANDVLIET *et al.* [20] showed that a reduction in the direct nonspecific bronchial responsiveness (PD₂₀ for histamine) increases gradually and first stabilizes after 20 months of treatment with inhaled budesonide in children with asthma. These findings can partly explain the weak response in the methacholine challenge test in the present study, with a treatment period of only 3 months.

With respect to spirometry, the high baseline FEV₁ values (>100% pred) and poor reversibility gave little room for improvement. Although statistically significant, the increase in FEV₁ in the twice-daily group was small. As pointed out by FERGUSON [21], FEF₂₅₋₇₅ is a more sensitive indicator of airflow than PEF or FEV₁. He suggested that frequent assessment of FEF₂₅₋₇₅ is required for optimal assessment of the effectiveness of therapy. In the present study this is demonstrated by the finding of a significant difference between BUD 100 µg twice daily and placebo as well as between BUD 100 µg twice daily and

BUD 200 µg once daily for FEF_{50%} and FEF_{75%}, otherwise the results were consistent when compared to FEV₁ or PEF.

The effect of switching from twice-daily to once-daily treatment with budesonide in well-controlled asthmatic children was studied by McCARTHY [22] and MÖLLER *et al.* [23]. They found that PEF, FEV₁, symptom scores and bronchodilator use during once-daily treatment were not different from those during twice-daily treatment.

In summary, the present study showed that low doses of inhaled budesonide administered at a dose of 100 µg or 200 µg once daily or 100 µg twice daily *via* Turbuhaler® provided protection against exercise-induced bronchoconstriction in children with mild asthma and that all three treatment regimens were well tolerated. An exercise test is suggested to be more sensitive than peak expiratory flow and forced expiratory volume in one second, in evaluating the efficacy of treatment with inhaled steroids in asthmatic children with mild disease and near-normal lung function.

Acknowledgements: Astra Draco AB, Lund, Sweden, supplied budesonide, terbutaline and placebo Turbuhaler® inhalers, as well as diaries and case report forms. The authors thank E. Edvardsen, C. Jonasson and B. Stache for their help and assistance, J. Leegaard and J. Grøgaard for their support and J. Ekelund for statistical help.

References

1. Warner JO, Neijens H, Landau LI. Asthma: a follow up statement from an international paediatric asthma consensus group. *Arch Dis Child* 1992; 67: 240–248.
2. Expert Panel Report II. Guidelines for the Diagnosis and Management of Asthma. National Asthma Education and Prevention Program. National Institutes of Health, February 1997.
3. Agertoft L, Pedersen S. Effects of long-term treatment with an inhaled corticosteroid on growth and pulmonary function in asthmatic children. *Respir Med* 1994; 88: 373–381.
4. König P, Shaffer J. The effect of drug therapy on long term outcome of childhood asthma: a possible preview of the international guidelines. *J Allergy Clin Immunol* 1996; 98: 1103–1111.
5. Woodhead M. Guidelines on the management of asthma. *Thorax* 1993; 48: S1–S24.
6. Pedersen S. Safety aspects of corticosteroids in children. *Eur Respir Rev* 1994; 4: 33–43.
7. Clark DJ, Grove A, Cargill RI, Lipworth BJ. Comparative adrenal suppression with inhaled budesonide and fluticasone propionate in adult asthmatic patients. *Thorax* 1996; 51: 262–266.
8. Lönnebo A, Grahén A, Jansson B, Brundin RM, Ling-Andersson A, Eckernäs S-Å. An assessment of the systemic effects of single and repeated doses of inhaled fluticasone propionate and inhaled budesonide in healthy volunteers. *Eur J Clin Pharmacol* 1996; 49: 459–463.
9. Price JF, Weller PH. Comparison of fluticasone propionate and sodium cromoglycate for the treatment of childhood asthma (an open parallel group study). *Respir Med* 1995; 89: 363–368.
10. Volovitz B, Amir J, Malik H, Kauschansky A, Varsano I. Growth and pituitary-adrenal function in children with severe asthma treated with inhaled budesonide. *N Engl J Med* 1993; 329: 1703–1708.

11. Pedersen S, Hansen OR. Budesonide treatment of moderate and severe asthma in children: A dose-response study. *J Allergy Clin Immunol* 1995; 95: 29-33.
12. Sheffer AL, Bousquet J, Busse WW, *et al.* International consensus report on diagnosis and treatment of asthma. *Eur Respir J* 1992; 5: 601-641.
13. Anonymous. Nordic consensus report. In: Nordiska Samarbetsnämnden för Medicinsk Forskning (NOS). Stockholm, Spri, 1992; pp. 19-20.
14. Oseid S, Haaland K. Exercise studies on asthmatic children before and after regular training. In: Erikson B, Furberg B, eds. *Swimming Medicine IV*. Baltimore, MD, University Park Press, 1978; pp. 32-41.
15. Yan K, Salome C, Woolcock AJ. Rapid method for measurement of bronchial responsiveness. *Thorax* 1983; 38: 760-765.
16. Cockcroft DW, Berscheid BA. Slope of the dose-response curve: usefulness in assessing bronchial response to inhaled histamine. *Thorax* 1983; 38: 55-61.
17. Zapletal A, Samanek M, Tuma S, Ruth C, Paul T. Assessment of airway function in children. *Bull Physiopathol Respir* 1972; 8: 535-543.
18. Uwytyed K, Springer C, Avital A, Bar-Yishay E, Godfrey S. Home recording of PEF in young asthmatics does it contribute to management? *Eur Respir J* 1996; 9: 872-879.
19. Waalkens HJ, Gerritsen J, Koëter GH, Krouwels FH, van Aalderen WMC, Knol K. Budesonide and terbutaline or terbutaline alone in children with mild asthma: effects on bronchial hyperresponsiveness and diurnal variation in peak flow. *Thorax* 1991; 46: 499-503.
20. Van Essen-Zandvliet EE, Hughes MD, Waalkens HJ, Duiverman EJ, Pocock SJ, Kerrebijn KF. Effects of 22 months treatment with inhaled corticosteroids and/or beta-2-agonists on lung function, airway responsiveness and symptoms in children with asthma. *Am Rev Respir Dis* 1992; 146: 547-554.
21. Ferguson AC. Persisting airway obstruction in asymptomatic children with asthma with normal peak expiratory flow rates. *J Allergy Clin Immunol* 1988; 82: 19-22.
22. McCarthy TP. The use of a once daily inhaled glucocorticosteroid (budesonide) in the management of childhood asthma. *Br J Clin Res* 1993; 4: 55-61.
23. Moller C, Strømberg L, Oldaeus G, Arweström E, Kjellman M. Administration of budesonide via Turbuhaler® (200 µg and 400 µg) once daily is as effective as when given twice daily in children with asthma. *Eur Respir J* 1996; 9: Suppl. 23, 115s.