



# Nerve growth factor enhances Clara cell proliferation after lung injury

S.S. Sonar\*, D. Schwinge\*, A. Kilic\*, A.Ö. Yildirim\*, M.L. Conrad\*, K. Seidler\*, B. Müller<sup>#</sup>, H. Renz\* and W.A. Nockher\*

**ABSTRACT:** The lung epithelia facilitate wound closure by secretion of various cytokines and growth factors. Nerve growth factor (NGF) has been well described in airway inflammation; however, its likely role in lung repair has not been examined thus far.

To investigate the repair function of NGF, experiments were performed *in vitro* using cultured alveolar epithelial cells and *in vivo* using a naphthalene-induced model of Clara epithelial cell injury.

Both *in vitro* and *in vivo* experiments revealed airway epithelial cell proliferation following injury to be dependent on NGF and the expression of its receptor, tropomyosin-receptor-kinase A. Additionally, NGF also augmented *in vitro* migration of alveolar type II cells. *In vivo*, transgenic mice over-expressing NGF in Clara cells (NGFtg) did not reveal any proliferation or alteration in Clara cell phenotype. However, following Clara cell specific injury, proliferation was increased in NGFtg and impaired upon inhibition of NGF. Furthermore, NGF also promoted the expression of collagen I and fibronectin *in vitro* and *in vivo* during repair, where significantly higher levels were measured in re-epithelialising NGFtg mice.

Our study demonstrates that NGF promotes the proliferation of lung epithelium *in vitro* and the renewal of Clara cells following lung injury *in vivo*.

**KEYWORDS:** Airway epithelium, alveolar type II cells, Clara cells, lung injury and repair, naphthalene, nerve growth factor

The mature healthy airway is a complex, yet fragile structure that is constantly exposed to inhaled environmental agents. Due to this close connection with the outside environment the lung is easily susceptible to damage; thus, injury and epithelial shedding are commonly described events in respiratory infections, asthma, chronic bronchitis and interstitial lung disease [1]. The lung has a remarkable capacity to facilitate repair through the use of specialised compartments with several distinctly functioning cell types. These cell types participate in a series of overlapping events to restore the respiratory epithelium. In the alveolar compartment of the lung, alveolar epithelial type II cells (AECII) are the putative progenitors of the alveolar epithelium and play an important role in the re-epithelialisation of alveolar walls after lung injury [2, 3]. Among the cell types lining the airway, the nonciliated bronchiolar cells (Clara cells) comprise the principal epithelial cell type of many species, including mice. Clara cells, which act as progenitors for goblet and ciliated epithelial cells

are major contributors to the airway repair process [4, 5]. In humans, Clara cells substantially contribute to the proliferation of tracheobronchial epithelium and have been described as an important cell population for the maintenance of the normal epithelium especially in the distal conducting airways [6].

Depending on the wound size and the degree of epithelial denudation, renewal of the damaged airway epithelium involves migration and proliferation of the residual epithelial cells from the adjoining, undamaged areas [7]. Expression of extracellular matrix (ECM) proteins in the provisional matrix are increased, and for Clara cells, factors like fibronectin and collagen I have been described to attract and provide adherence points for migration [8].

The Clara cell is uniquely susceptible to injury due to its capability to transform chemicals like naphthalene (NA) into toxic intermediates. NA is a polycyclic aromatic hydrocarbon commonly found in cigarette smoke and diesel exhaust and

## AFFILIATIONS

\*Dept of Clinical Chemistry and Molecular Diagnostics, University Hospital Marburg, and  
<sup>#</sup>Laboratory of Respiratory Cell Biology, Dept of Internal Medicine, University Hospital Marburg, Germany.

## CORRESPONDENCE

S.S. Sonar  
Dept of Clinical Chemistry and Molecular Diagnostics  
Biomedical Research Center  
Hans Meerweinstrasse  
35043 Marburg  
Germany  
E-mail: sanchaitas@yahoo.co.in

## Received:

Nov 03 2008

Accepted after revision:

Nov 04 2009

First published online:

Jan 14 2010

This article has supplementary material available from [www.erj.ersjournals.com](http://www.erj.ersjournals.com)

is capable of inducing a highly dose-dependent and cell type- and site-selective toxicity [9, 10]. The toxicity of NA is due to the metabolism of this compound by cytochrome P450 monooxygenases to a toxic intermediate that causes Clara cell swelling, vacuolisation, and exfoliation into the lumen of the airways 24 h after injury is initiated [11]. The use of NA in this study enabled us to investigate the ongoing repair processes following Clara cell specific injury *in vivo*.

Following injury, the mediators that drive airway epithelial repair may be derived from many sources. In addition to factors released from activated local immune cells [12], epithelial cells are also likely sources of autocrine and/or paracrine growth factors. One such mediator actively secreted by the airway epithelium is nerve growth factor (NGF), an eminent member of the neurotrophin growth factor family initially described to promote neuronal growth and survival [13]. Recently, the importance of NGF and its high affinity receptor, the tropomyosin-receptor-kinase A (TrkA) has been described in inflammatory diseases related to airway dysfunction [14, 15]. Examples of this include NGF over-expression in the activated allergic bronchial [14], nasal [15] and airway epithelium as well as elevated epithelial NGF mediating eosinophil infiltration [16] and survival [17]. NGF is also expressed in the epithelia of other tissues, especially after activation *via* inflammation or during regenerative conditions [18–21]. Despite a few reports suggesting a role for NGF in wound healing elsewhere [22–24], its involvement in lung epithelial repair processes has not been investigated thus far.

The objective of this study was to examine the role of NGF in the repair of injured airway epithelia and our results highlight its contribution towards the proliferation of Clara airway epithelial cells in the lung following injury.

## METHODS

### Mice

6–8-week-old female C57BL/6 mice were obtained from Harlan Winkelmann (Borchen, Germany). Age-matched transgenic mice overexpressing NGF under the control of the lung-specific Clara-cell secretory protein promoter (NGFtg) were used [25]. All animal experiments were performed according to “The guidelines for the care and use of experimental animals” prepared and published by the Society for Laboratory Animal Sciences [26].

### Wound assay for lung epithelial cells

LA4 murine lung alveolar epithelial cells (American Type Culture Collection, Rockville, IN, USA) seeded at  $1 \times 10^6$  cells·well<sup>-1</sup> and supplemented with F12K Nutrient Mixture containing 15% FCS (Gibco, Carlsbad, CA, USA) were cultured until confluence onto 12-well plates. After overnight (o/n) serum starvation, monolayers were scratched from top to bottom using a 200 µL pipette tip then incubated with either medium, medium containing recombinant human NGF ( $\beta$ -NGF, 100 ng·mL<sup>-1</sup>; R&D systems, Minneapolis, MN, USA), polyclonal blocking antibodies to NGF (anti-NGF, 625 ng·mL<sup>-1</sup>; Peprotech, Rocky Hill, NJ, USA), or immunoglobulin (Ig)G control (625 ng·mL<sup>-1</sup>; Sigma, St. Louis, MO, USA). All groups were incubated in medium containing 1% FCS. Medium with the accompanying supplements was changed every 24 h. Wound size was measured in µm by light microscopy (Olympus, Hamburg, Germany) after

wounding and every 24 h for 72 h using the CellF imaging software program (Soft Imaging System GmbH, Münster, Germany). At least 3–4 wounds of six LA4 cell monolayers were analysed.

### Proliferation of cultured lung epithelial cells

Primary AECII were isolated according to the protocol provided in the supplementary material.  $1 \times 10^5$  AECII or LA4 cells·well<sup>-1</sup> were grown in 96-well plates in triplicates. AECII were grown overnight in medium containing 10% FCS followed by medium containing 0.1, 1 or 10% FCS for 24, 48 and 72 h. Proliferation of AECII cells was further assessed with 0.1% FCS containing medium, NGF, anti-NGF or control IgG for 72 h in concentrations described above for wound assay. For LA4 cells, 20 pmol·µL<sup>-1</sup> of NGF, TrkA or control siRNA (Qiagen, Hilden, Germany) were incubated with cells according to the manufacturer’s instructions. Cell proliferation assays were performed using a BrdU (5-bromo-2'-deoxy-uridine) labelling kit according to the manufacturer’s instructions (Roche, Mannheim, Germany). For further details, including primer sequences, see the supplementary material.

### In vitro migration assay using AECII

AECII were assayed for NGF-induced migration using 24-well modified Boyden chamber (Falcon; BD, Franklin Lakes, NJ, USA) using transwell membranes (8-µm pore size). The upper chamber contained  $10^5$  AECII in 500 µL medium and the lower chamber contained 500 µL of cell-free medium containing NGF (100 ng·mL<sup>-1</sup>). Cells were incubated for 6 h at 37°C with 5% CO<sub>2</sub> after which inserts were removed and cells on the lower surface were counterstained with Diff-Quick (Dade, Behring, Germany) for 5 min. An average of five × 100 fields per insert was counted of five samples per group using light microscopy (Olympus, Hamburg, Germany). Migration is expressed as migrated cells per × 100 field.

### In vivo model of Clara cell injury and denudation using NA

Wildtype (WT) or NGFtg control mice received intraperitoneal (*i.p.*) corn oil (Sigma Aldrich, Munich, Germany), WT or NGFtg mice undergoing Clara cell exfoliation received *i.p.* NA (250 mg·kg<sup>-1</sup>, Sigma Aldrich Munich, Germany) dissolved in corn oil at day 0 (fig. S1a and b in the supplementary material). In additional groups, WT control mice and WT mice treated with NA also received either intranasal anti-NGF (anti-NGF, 2.5S, N 6655; Sigma-Aldrich) or intranasal control IgG (Sigma-Aldrich) at a concentration of 24 µg at day 0, 3, 6 and 9 (fig. S1a and b in the supplementary material). A new set of mice also received BrdU in drinking water at a concentration of 1 mg·mL<sup>-1</sup> (water containing 1% glucose) from day 0 to day 10 (fig. S1a in the supplementary material). On days 1, 10 or 29 after treatment, bronchoalveolar lavage (BAL) was performed twice with 800 µL of ice-cold PBS plus protease inhibitor mix (Roth, Karlsruhe, Germany) and whole lungs were removed for histological, mRNA and protein analysis.

### NGF ELISA

NGF ELISA (Promega, Madison, WI, USA) was performed according to the manufacturer’s instructions with cell free culture supernatants of six LA4 cell monolayers (with and without wound at 6, 12, 24, 48 and 72 h) and BAL Fluid (BALF) of WT and NGFtg animals (control and NA treated at days 10

and 29). Plates were read in a microplate autoreader (Tecan, Salzburg, Austria) at 405 nm.

### Immunoblot analysis

Total protein was extracted from LA4 cell monolayers, AECII and lung tissues of WT and NGFtg control and NA treated mice using the M-PER mammalian protein extraction kit (Thermo Scientific, Pierce, Bonn, Germany). Immunoblotting was performed as previously described [27] with NGF, TrkA (1:500, Santa Cruz Heidelberg, Germany), anti- $\beta$ -actin monoclonal antibody (1:40,000; Sigma) and secondary peroxidase-conjugated antibodies (1:5000; Sigma). Immunoreactive protein bands were visualized with a chemiluminescent substrate (SuperSignal; Thermo Scientific, Pierce). Bands were scanned with the Kodak Image Station CF 450 and analysed using the Kodak 1D 3.5 imaging software. See the supplementary material for further details.

### RNA isolation, real-time PCR and analysis

$5 \times 10^5$  AECII  $\cdot$  mL<sup>-1</sup> were incubated in 24-well plates with either anti-NGF (625 ng  $\cdot$  mL<sup>-1</sup>), control IgG (625 ng  $\cdot$  mL<sup>-1</sup>), or recombinant human NGF (100 ng  $\cdot$  mL<sup>-1</sup>) for 24 h. Total RNA was isolated from AECII or lung tissue using the RNeasy Mini Kit (Qiagen, Hilden, Germany). Quantitative PCR was performed using the QuantiTect SYBR<sup>®</sup> Green PCR Kit (Qiagen). Primers specific for collagen I, and fibronectin were used for real-time PCR. The mRNA copy numbers were compared and expressed relative to mouse ribosomal L32 as the house-keeping gene. See the supplementary material for further details.

### Immunohistochemistry


Immunohistochemistry was performed on sectioned paraffin embedded lung tissues as described previously [28]. Primary antibody concentrations were; anti-Clara cell specific protein (CCSP) (diluted 1:3,000; courtesy of J. Klug, Justis-Liebig-Universität, Giessen, Germany); anti-fibronectin (diluted 1:1,000; Biozol, Eching, Germany), anti-collagen I (diluted 1:500; Biozol), sheep anti-BrdU (diluted 1:10; Biozol). Secondary biotinylated antibodies (Vector Laboratories, Burlingame, CA, USA) were diluted 1:400. Sections were analysed using light microscopy (Olympus Europa GmbH, Hamburg, Germany).

### Morphometric analysis of Clara cells and BrdU-incorporating proliferating cells

Mouse lungs were fixed by airway instillation using 6% phosphate buffered paraformaldehyde at a pressure of 20 cm fluid column. Lung volume was determined by fluid displacement, and systemic uniform random samples of lung tissue were taken and processed according to standard methods [29]. The morphometric analysis for CCSP- and BrdU-positive cells was performed using a modified version of a previously described method [30]. Proximal and distal airways were counted in a blinded manner. CCSP-positive cells containing nuclear profiles and BrdU-positive nuclear profiles were counted. See online supplement for further details.

### Statistical analysis

Statistical significance for normally distributed samples was analysed by using an unpaired t-test. Comparisons between more than two groups were made by one way ANOVA. A p-value




**FIGURE 1.** Production of nerve growth factor (NGF) and expression of TrkA *in vitro* after LA4 cell monolayer wounding. a) NGF production was measured by ELISA in unwounded control (□), and wounded (■) LA4 cells at 6, 12, 24, 48 and 72 h. b) The expression of TrkA was examined by quantifying intensity of bands normalised to  $\beta$ -actin by western blotting in control (□) and LA4 monolayers with wound (■) at 24, 48 and 72 h. c) A representative blot showing expression of TrkA (140 kDa) and  $\beta$ -actin (42 kDa) in wounded and control LA4 monolayers at 24, 48 and 72 h from the same experiment as in (b). Data are presented as mean  $\pm$  sem (n=6) at each time point. \*: p<0.05; \*\*: p<0.01; \*\*\*, p<0.001.

of <0.05 was considered statistically significant. All numerical data are expressed as mean  $\pm$  SEM. Calculations were performed using Graph Pad prism software, version 3.02.


## RESULTS

### NGF production and TrkA expression is increased in wounded lung epithelial cells *in vitro*

The expression of NGF in lung epithelial cells was first examined in an *in vitro* wound assay model using murine LA4 cells. Wounding confluent monolayers of LA4 cells led to an increase in the expression of both NGF and its receptor, TrkA (fig. 1). NGF protein levels in wounded epithelial cells were significantly higher compared with nonwounded LA4 cells at



**FIGURE 2.** Nerve growth factor (NGF) regulates wound closure of LA4 cells *in vitro*. a, b) Confluent monolayers of LA4 cells were wounded and the kinetics of wound closure was monitored in monolayers grown in medium only (medium; □), medium with recombinant NGF (NGF; ○), immunoglobulin (IgG) control (IgG; ●), or with anti-NGF (anti-NGF; ■) at 0, 24, 48, and 72 h. Wound closure was quantified by measuring wound length in μm across the wound. Statistical comparisons for NGF-stimulated and anti-NGF-treated groups were made to the medium only group. c) Cell proliferation was determined by measuring incorporation of BrdU (5-bromo-2'-deoxy-uridine) in DNA of proliferating LA4 cells. The rate of proliferation is represented relative to basal proliferation in the medium control where BrdU incorporation was taken as 100% (.....). TrkA (■) and NGF (■) were blocked by specific siRNA and compared to cells grown in the transfection assay medium (□) and a non-specific siRNA (■). Data are presented as mean ± SEM (n=6) for each time point. \*\*\*: p<0.001. Scale bars=100 μm for all panels.



**FIGURE 3.** Nerve growth factor (NGF) regulates proliferation, migration and extracellular matrix mRNA expression of alveolar epithelial type II cells (AECII). a) NGF and b) TrkA protein expression by AECII. c) AECII proliferation analysed after 72 h incubation with either medium only or medium containing NGF, anti-NGF or control immunoglobulin (IgG). d) AECII migration analysed after 6 h incubation with medium or recombinant NGF. Migration is expressed as migrated cells per  $\times 100$  field of view. Relative mRNA levels of e) fibronectin or f) collagen I in AECII cells incubated for 24 h in either medium only or medium containing NGF, anti-NGF or control IgG. Data are presented as mean  $\pm$  SEM (n=6) copy number target gene per copy L32 (relative mRNA) for each group. \*:  $p < 0.05$ ; \*\*\*:  $p < 0.001$ .

all time points measured (fig. 1a). A progressive increase in NGF after wounding was observed until 72 h, with elevated levels observed as early as 6 h. NGF production also increased with time in the nonwounded cells, correlating with normal cell proliferation. Correspondingly, TrkA was upregulated at 24 and 48 h post wounding and was present in significantly higher amounts compared with nonwounded monolayers (fig. 1b). The increase in TrkA expression coincided with the increase of NGF in the supernatant (fig. 1a). These results indicate that autocrine NGF and TrkA are induced upon lung epithelial cell injury.

#### NGF promotes wound closure and proliferation of epithelial cells *in vitro*

To determine if the observed upregulation of NGF and its receptor has an effect on the closure of wounds, wounded monolayers of LA4 cells were incubated with either blocking antibodies to NGF (anti-NGF) or with exogenous NGF. Inhibition of NGF markedly delayed closure of wounds, whereas exogenous NGF protein significantly accelerated wound closure (fig. 2a and b). Control wounded monolayers incubated with control IgG antibodies or medium only had wounds of similar size (fig. 2a and b). In control monolayers, wound closure was achieved at 72 h, whereas wounds failed to

close in the anti-NGF treated groups even at 72 h, (fig. 2a). In contrast, wounds incubated with NGF closed at an earlier time point of 48 h. Since the proliferative effect of NGF is reported to be mediated by its high-affinity receptor TrkA, the effect of NGF and TrkA on the regulation of lung epithelial cell proliferation was also investigated *in vitro*. siRNA inhibition of NGF or TrkA knocked down the mRNA and protein levels to significantly lower levels compared to the medium control (fig. S2 in the supplementary material). Inhibition of TrkA and NGF with these siRNAs significantly decreased proliferation of LA4 epithelial cells by  $\sim 75$  and 50% of medium control, respectively (fig. 2c). These results indicate that NGF and TrkA are important for the proliferation and repair of wounded lung epithelia *in vitro*.

#### NGF mediates proliferation, migration and ECM expression by AECII

After determining the important role of NGF in a lung epithelial cell line, we were interested to examine the effect of NGF and TrkA in primary AECII. NGF and TrkA were found to be expressed in primary AECII (fig. 3a and b). In order to test the role of NGF on primary AECII cell proliferation, we established an *in vitro* assay. A time and dose response for AECII proliferation *in vitro* at 24, 48 and 72 h

in 0.1%, 1% and 10% FCS concentration (fig. S3c in the supplementary material) revealed 72 h as the optimal time point to examine proliferation. NGF was seen to significantly influence proliferation at 0.1% FCS concentration. In line with our previous *in vitro* results using LA4 cells, NGF increased the proliferative capacity of AECII, whereas anti-NGF significantly downregulated proliferation compared with the medium and IgG control (fig. 3c). We also investigated the role of NGF in mediating migration, which is a critical event in wound repair. NGF was seen to strongly induce migration of AECII upon incubation with exogenous NGF (fig. 3d). Furthermore, the expression of ECM factors, such as collagen I and fibronectin were also investigated *in vitro* in primary cell cultures of AECII. The mRNA expression of fibronectin and collagen I was significantly increased upon incubation with exogenous NGF (fig. 3e and f) and the inhibition of NGF led to a significant decrease in fibronectin mRNA expression (fig. 3e). These results imply that NGF actively promotes proliferation and migration of AECII, which are critical requisites for wound repair.

#### NGF and TrkA are increased during repair of airway epithelia *in vivo*


To examine the role of NGF in airway epithelium repair *in vivo*, we employed a NA induced model of Clara epithelial cell injury in the lung. Expression of NGF and TrkA was determined at 10 and 29 days post injury with NA in the BAL fluid and total lung protein of WT and NGFtg animals (fig. 4). In untreated NGFtg control mice, the baseline level of NGF was significantly higher due to the constitutive over-expression of NGF under the Clara cell specific-promoter compared to WT control animals (fig. 4a). Following injury with NA, WT animals showed a progressive upregulation of NGF which lasted until day 29 (fig. 4a). Conversely, NGF levels in the NGFtg animals declined after NA treatments at all time points compared to untreated NGFtg animals due to the NA induced loss of NGF over-expressing Clara cells. However, the amount of NGF in the NA treated NGFtg animals was significantly higher at all time points compared to NA treated WT animals. The over-expression of NGF in Clara cells had no effect on the total number of Clara cells lining the airways (data not shown) and the denudation of Clara cells 24 h after NA injury was found to be similar in NA treated WT (WT-NA) and NGFtg (NGFtg-NA) animals (fig. S4A in the supplementary material). Additionally, the number of exfoliated cells quantified in the BALF of both groups of mice was comparable (fig. S4b in the supplementary material).

To investigate the role of TrkA as a downstream target of NGF during the repair processes, its levels were quantified concomitantly in lung tissues. TrkA was found to be significantly upregulated at days 10 and 29 post Clara cell injury in the NGFtg animals compared with WT animals (fig. 4b).

These results suggest that NGF and its receptor, TrkA, are upregulated upon airway epithelial damage and their induction is maintained throughout the ongoing repair process.


#### NGF promotes renewal of Clara cells

We next tested the role of NGF in the renewal of Clara cells upon injury. Clara cells comprise a major epithelial cell population lining the murine bronchial airways. Following



**FIGURE 4.** Expression of nerve growth factor (NGF) and TrkA *in vivo* following Clara cell lung injury. a) NGF levels in NGFtg (■) and wildtype (□) animals without naphthalene (NA) treatment (control) and after NA treatment at day 10 and 29 (d10, d29). NGF expression was measured by ELISA in bronchoalveolar lavage. b) The expression of TrkA was quantified and represented as band intensity normalised to the housekeeping gene,  $\beta$ -actin. c) A representative blot showing expression of TrkA (140 kDa) and  $\beta$ -actin (40 kDa) in control NGFtg and wild-type (WT) animals (control) and after NA treatment at day 10 and 29 (d10, d29) from the experiment is shown below the graph. Data are presented as mean  $\pm$  SEM ( $n=6$ ) for each group of mice. \*:  $p<0.05$ , \*\*:  $p<0.01$ , \*\*\*:  $p<0.001$ .

NA treatment, Clara cells were denuded by day 1 (fig. 5a). At day 10, Clara cells in the airways had started reappearing around the airways in all groups treated with NA, as quantified by CCSP staining (fig. 5b). Mice treated with NA or NA and control IgG showed similar Clara cell numbers at day 10, with considerable portions of the airway already lined by Clara cells. In contrast, animals treated with NA and neutralising anti-NGF antibodies, had significantly lower counts of CCSP-positive cells lining the airway at day 10, reappearing only sporadically around the airway epithelium (fig. 5). These results indicate



**FIGURE 5.** Clara cell renewal. a) Intact Clara cells in corn oil treated wild-type (WT) control animals at day 1 and day 10 (WT-control) and after naphthalene (NA) treatment (WT-NA) with anti-nerve growth factor (NGF) (WT-anti-NGF-NA) or control immunoglobulin (IgG) (WT-IgG-NA) antibodies. Clara cells were stained for Clara cell secretory protein (CCSP) as brown-coloured nonciliated columnar cells against the blue counter stain. Scale bars=50  $\mu$ m. b) Quantification of CCSP-positive cells per mm of the basement membrane at day 10 after NA treatment. Data are presented as mean  $\pm$  SEM (n=6) for each group of mice. \*: p<0.05, \*\*: p<0.01, \*\*\*: p<0.001.

that NGF facilitates Clara cell renewal *in vivo* upon NA-induced injury.

#### **NGF induces collagen I and fibronectin mRNA expression upon NA induced Clara cell injury**


Cellular migration and proliferation critically depend on the adhesive interactions between cells and the underlying matrix. Also, considering that Clara cells show preferential adherence to collagen I and fibronectin, we utilised the NA model to examine the effect of NGF on the expression of these matrix molecules (fig. 6). Indeed, collagen I and fibronectin mRNA levels were significantly increased in lung homogenates of NGFtg animals at day 10 after NA exposure compared to untreated NGFtg control animals (fig. 6a and b). Moreover, the application of anti-NGF reduced the expression of collagen I in NA treated WT animals (fig. 6a). Immunohistochemical staining for collagen I and fibronectin corroborated the qPCR data and more intense collagen I and fibronectin staining underlying the epithelial layer was observed in the control and NA treated NGFtg animals compared with their respective controls (fig. 6c and d). The application of anti-NGF revealed relatively reduced staining for collagen I in NA-treated WT animals. These results suggest that NGF actively augments the expression of the ECM factors collagen I and fibronectin during repair processes in the lung.

#### **NGF promotes proliferation of Clara cells in the lung**

After observing that NGF promotes Clara cell renewal as well as the expression of collagen I and fibronectin expression, we investigated the direct effect of NGF on epithelial cell proliferation by administration of BrdU *in vivo* (fig. S1b in the supplementary material). Quantification of BrdU-positive proliferating cells revealed lack of proliferation in the airways of untreated WT and NGFtg control animals, whereas all NA treated groups contained BrdU positive nuclei (fig. 7). Staining serial sections in parallel with BrdU and CCSP revealed proliferating Clara cells (fig. 7a). Injury and cell denudation was therefore shown to induce re-epithelialisation *via* increased Clara cell proliferation in the lung. Upon inhibition of NGF with anti-NGF antibody, significantly lower numbers of proliferating epithelial cells were observed, while application of a control IgG antibody showed no effect. In line with these results, NGFtg mice treated with NA exhibited the highest numbers of proliferating epithelial cells (fig. 7b). Together, these results show strong effects of NGF on Clara cell proliferation during repair of injured lung.

#### **DISCUSSION**

NGF and other members of the neurotrophin family were initially identified as neuronal growth and survival factors but are currently recognised to have biological activities in tissues




**FIGURE 6.** Modulation of collagen I and fibronectin expression by nerve growth factor (NGF). Relative mRNA levels of a) collagen I and b) fibronectin are shown. Data are expressed as copy number of target gene per copy L32 (relative mRNA). mRNA was analysed in wild-type (WT; □), NGFtg (▨), WT animals treated with immunoglobulin (Ig)G (■) or anti-NGF (▩). Immunohistochemical staining of c) collagen I and d) fibronectin in corn oil-treated WT (WT-control) and NGFtg (NGFtg-control) animals and NA-treated WT (WT-NA), with anti-NGF (WT-anti-NGF-NA) or with control IgG (WT-IgG-NA) and NA-treated NGFtg (NGFtg-NA) animals analysed at day 10. Scale bars=50  $\mu$ m. Data are presented as mean  $\pm$  SEM (n=6) for each group of mice. \*:  $p < 0.05$ ; \*\*\*:  $p < 0.001$ .

outside the nervous system [31]. Various visceral epithelia are known to produce neurotrophins which were initially speculated to regulate tissue innervation [32] and autocrine functions of neurotrophins are increasingly being described. For example, NGF production is increased during inflammatory processes that are accompanied by epithelium damage in asthma [17, 33], infections [34] and sarcoidosis [35].

Previous reports have suggested the involvement of NGF in epithelial wound healing in other tissues. Topical application of NGF has been shown to accelerate wound closure and

re-epithelialisation in surgical wounds of the skin [36] and the cornea [23]. Moreover, NGF is strongly produced at the site of the wounded skin and is increased in newly formed epithelial cells at the edge of the wound but not in the surrounding epidermal keratinocytes [37]. In the cornea, an increase in NGF was observed after epithelial injury and corneal epithelial cell expression of NGF and its receptor TrkA was observed [38]. When epithelial surfaces are injured, the normal response is to upregulate receptors for growth factors that drive proliferation and repair [39]. The purpose of the present study was to determine the function of NGF in facilitating proliferation of





**FIGURE 7.** Analysis of proliferation by BrdU (5-bromo-2'-deoxy-uridine) incorporation in bronchial epithelium. a) Serial sections of wild-type (WT) control (WT-control), naphthalene (NA)-treated WT (WT-NA) with control immunoglobulin (Ig)G (WT-IgG-NA) or with anti-nerve growth factor (NGF) (WT-anti-NGF-NA) and NGFtg control (NGFtg-control) and NA treated NGFtg (NGFtg-NA) animals stained for Clara cell secretory protein (CCSP) and BrdU are shown. The CCSP-positive Clara cells are stained brown against the blue counter stain and the BrdU-positive cells exhibit a dark brown nucleus against the nuclear fast red counter stain. All animals treated with NA showed positive nuclear staining for BrdU. Scale bars=50  $\mu$ m. b) Quantification of BrdU positive nuclear profiles per mm of the basement membrane at day 10 after NA treatment. Data are presented as mean  $\pm$  SEM (n=6) for each group of mice. \*: p<0.05.

lung epithelial cells, especially Clara cells that comprise the predominant airway epithelial cell population lining murine airways.

The contribution of epithelial NGF in wound repair was first investigated *in vitro* using both a clonal murine lung alveolar type II epithelial cell line (LA4) and primary murine AECII. *In vitro* experiments with wounded LA4 cell monolayers revealed increased protein expression of NGF and its receptor TrkA upon injury. Furthermore, wound closure was accelerated with NGF supplementation and suppressed upon NGF inhibition. Investigation of primary AECII cells, which also express NGF and TrkA, revealed a chemotactic response of AECII to NGF. AECII cells also responded to NGF by expressing increased collagen I and fibronectin mRNA. Most importantly, in both cell types, NGF was seen to strongly mediate proliferation. We then investigated the pro-proliferative function of NGF *in vivo* by employing a NA model of Clara cell injury.

We investigated the functional effects of NGF on airway epithelium repair in the lung *in vivo* by utilising a Clara cell-specific injury model. The complete renewal of Clara cells after NA mediated injury takes 1–3 months. The NA model of epithelial injury was investigated for a 29-day period to assess the role of NGF in lung epithelia repair. For this, we included NGFtg animals that overexpress NGF constitutively under the CCSP promoter in Clara cells. The NGFtg animals express around 155-fold higher NGF levels compared with WT mice. The presence of increased NGF levels under normal conditions caused no alteration in the total numbers of Clara cells or any change in cell phenotype. Furthermore, WT and NGFtg animals revealed similar levels of injury at day 1 demonstrated by comparable numbers of exfoliated Clara cells in both mouse strains (fig. S4 in the supplementary material). The NGFtg animals lost most of the NGF-overexpressing Clara cells upon exfoliation; however, the remaining cells produced significantly higher levels of NGF compared with WT animals, which

may be associated with the accelerated epithelial repair seen in the NGFtg animals. Following injury of Clara cells, lung NGF levels were increased at day 10 and remained significantly elevated until day 29 post injury in both WT and NGFtg mice, emphasising the importance of NGF in the continuing lung repair processes. This is supported by *in vitro* findings with wounded lung epithelial cells, where NGF synthesis was observed to be upregulated as early as 6 h after injury and increased progressively with time. Interestingly, the lack of proliferation observed in uninjured lungs of NGFtg animals indicates that the pro-proliferative function of NGF is limited to regenerative situations. Therefore, Clara cell renewal was, at least in part, dependent on the pro-proliferative function of NGF post injury, where BrdU incorporation *in vivo* was enhanced in NGFtg animals and could be inhibited by anti-NGF antibodies.

The overexpression of NGF in the lungs of NGFtg mice has been previously reported to modulate lung innervation and regulate airway hyperresponsiveness [25]. NGF has been suggested as a potential candidate for airway remodelling in asthma and also reported to induce smooth muscle cell proliferation [40, 41], migration and differentiation of fibroblasts [42]. In this study, NGF also appeared to modulate collagen I and fibronectin expression, supported by the observations *in vitro* in AECII and *in vivo* in the NA model of Clara cell injury. Collagen I and fibronectin mRNA expression were significantly increased upon incubation of primary AECII with exogenous NGF and similar results were seen *in vivo* under higher NGF levels in lungs of transgenic animals after injury. Based on these findings we suggest that NGF could potentially modulate ECM expression in other processes involving injury and repair, such as airway remodelling in diseases like asthma, where NGF has been previously implicated. Future investigation of the role of NGF in remodelling and fibrosis may shed further light on mechanisms that turn reparative healing processes to pathological events.

In summary, our findings provide evidence that NGF orchestrates proliferative events during repair of the lung epithelia, promoting the renewal of Clara cells *in vivo*. Therefore, NGF may be considered as an important autocrine epithelial growth factor governing repair in the bronchial airway. While neurotrophins were originally described as molecules that maintain neuronal outgrowth and survival, there is now growing evidence that they belong to a class of multifunctional signalling molecules, which can also mediate repair and proliferation of lung epithelia.

#### SUPPORT STATEMENT

This work was supported, in part, by grants from the Deutsche Forschungsgemeinschaft (Transregio 22, Project A9), and Stiftung für Pathobiochemie und Molekulare Diagnostik, DGKL.

#### STATEMENT OF INTEREST

None declared.

#### ACKNOWLEDGEMENTS

We thank S. Achenbach, E. Kotte for their excellent technical assistance and T. Rausch for performing the CC10 immunohistochemistry.

#### REFERENCES

- 1 Knight DA, Holgate ST. The airway epithelium: structural and functional properties in health and disease. *Respirology* 2003; 8: 432–446.
- 2 Mason RJ. Biology of alveolar type II cells. *Respirology* 2006; 11: S12–S15.
- 3 Lee TS, Liu YJ, Tang GJ, *et al.* Wood smoke extract promotes both apoptosis and proliferation in rat alveolar epithelial type II cells: the role of oxidative stress and heme oxygenase-1. *Crit Care Med* 2008; 36: 2597–2606.
- 4 Giangreco A, Reynolds SD, Stripp BR. Terminal bronchioles harbor a unique airway stem cell population that localizes to the bronchoalveolar duct junction. *Am J Pathol* 2002; 161: 173–182.
- 5 Reynolds SD, Giangreco A, Power JH, *et al.* Neuroepithelial bodies of pulmonary airways serve as a reservoir of progenitor cells capable of epithelial regeneration. *Am J Pathol* 2000; 156: 269–278.
- 6 Boers JE, Ambergen AW, Thunnissen FB. Number and proliferation of clara cells in normal human airway epithelium. *Am J Respir Crit Care Med* 1999; 159: 1585–1591.
- 7 Zahm JM, Chevillard M, Puchelle E. Wound repair of human surface respiratory epithelium. *Am J Respir Cell Mol Biol* 1991; 5: 242–248.
- 8 Atkinson JJ, Adair-Kirk TL, Kelley DG, *et al.* Clara cell adhesion and migration to extracellular matrix. *Respir Res* 2008; 9: 1.
- 9 Buckpitt AR, Castagnoli N Jr, Nelson SD, *et al.* Stereoselectivity of naphthalene epoxidation by mouse, rat, and hamster pulmonary, hepatic, and renal microsomal enzymes. *Drug Metab Dispos* 1987; 15: 491–498.
- 10 Plopper CG, Suverkropp C, Morin D, *et al.* Relationship of cytochrome P-450 activity to Clara cell cytotoxicity. I. Histopathologic comparison of the respiratory tract of mice, rats and hamsters after parenteral administration of naphthalene. *J Pharmacol Exp Ther* 1992; 261: 353–363.
- 11 Van Winkle LS, Buckpitt AR, Nishio SJ, *et al.* Cellular response in naphthalene-induced Clara cell injury and bronchiolar epithelial repair in mice. *Am J Physiol* 1995; 269: L800–L818.
- 12 Takizawa H, Beckmann JD, Shoji S, *et al.* Pulmonary macrophages can stimulate cell growth of bovine bronchial epithelial cells. *Am J Respir Cell Mol Biol* 1990; 2: 245–255.
- 13 Lewin GR, Barde YA. Physiology of the neurotrophins. *Annu Rev Neurosci* 1996; 19: 289–317.
- 14 Freund-Michel V, Frossard N. The nerve growth factor and its receptors in airway inflammatory diseases. *Pharmacol Ther* 2008; 117: 52–76.
- 15 Raap U, Fokkens W, Bruder M, *et al.* Modulation of neurotrophin and neurotrophin receptor expression in nasal mucosa after nasal allergen provocation in allergic rhinitis. *Allergy* 2008; 63: 468–475.
- 16 Path G, Braun A, Meents N, *et al.* Augmentation of allergic early-phase reaction by nerve growth factor. *Am J Respir Crit Care Med* 2002; 166: 818–826.
- 17 Hahn C, Islamian AP, Renz H, *et al.* Airway epithelial cells produce neurotrophins and promote the survival of eosinophils during allergic airway inflammation. *J Allergy Clin Immunol* 2006; 117: 787–794.
- 18 Stanzel RD, Lourensens S, Blennerhassett MG. Inflammation causes expression of NGF in epithelial cells of the rat colon. *Exp Neurol* 2008; 211: 203–213.
- 19 Marinova TS, Markova M, Aloe L. NGF overexpression and distribution in cortical thymic epithelial cells of mice with experimentally-induced leukemia. *In Vivo* 2006; 20: 259–264.
- 20 Hayashi K, Storesund T, Schreurs O, *et al.* Nerve growth factor  $\beta$ /pro-nerve growth factor and their receptors in normal human oral mucosa. *Eur J Oral Sci* 2007; 115: 344–354.
- 21 Gigliozzi A, Alpini G, Baroni GS, *et al.* Nerve growth factor modulates the proliferative capacity of the intrahepatic biliary epithelium in experimental cholestasis. *Gastroenterology* 2004; 127: 1198–1209.

- 22 Muangman P, Muffley LA, Anthony JP, *et al.* Nerve growth factor accelerates wound healing in diabetic mice. *Wound Repair Regen* 2004; 12: 44–52.
- 23 Cellini M, Bendo E, Bravetti GO, *et al.* The use of nerve growth factor in surgical wound healing of the cornea. *Ophthalmic Res* 2006; 38: 177–181.
- 24 Nithya M, Suguna L, Rose C. The effect of nerve growth factor on the early responses during the process of wound healing. *Biochim Biophys Acta* 2003; 1620: 25–31.
- 25 Hoyle GW, Graham RM, Finkelstein JB, *et al.* Hyperinnervation of the airways in transgenic mice overexpressing nerve growth factor. *Am J Respir Cell Mol Biol* 1998; 18: 149–157.
- 26 Society for Laboratory Animal Sciences. The guidelines for the care and use of experimental animals. Biberach an der Riss, GV-SOLAS, 1988.
- 27 Yildirim AO, Bulau P, Zakrzewicz D, *et al.* Increased protein arginine methylation in chronic hypoxia: role of protein arginine methyltransferases. *Am J Respir Cell Mol Biol* 2006; 35: 436–443.
- 28 Sel S, Wegmann M, Dicke T, *et al.* Effective prevention and therapy of experimental allergic asthma using a GATA-3-specific DNase. *J Allergy Clin Immunol* 2008; 121: 910–916.
- 29 Weibel ER, Hsia CC, Ochs M. How much is there really? Why stereology is essential in lung morphometry. *J Appl Physiol* 2007; 102: 459–467.
- 30 Stevens TP, McBride JT, Peake JL, *et al.* Cell proliferation contributes to PNEC hyperplasia after acute airway injury. *Am J Physiol* 1997; 272: L486–L493.
- 31 Nockher WA, Renz H. Neurotrophins in allergic diseases: from neuronal growth factors to intercellular signaling molecules. *J Allergy Clin Immunol* 2006; 117: 583–589.
- 32 Lommatzsch M, Braun A, Mannsfeldt A, *et al.* Abundant production of brain-derived neurotrophic factor by adult visceral epithelia. Implications for paracrine and target-derived Neurotrophic functions. *Am J Pathol* 1999; 155: 1183–1193.
- 33 Frossard N, Naline E, Olgart HC, *et al.* Nerve growth factor is released by IL-1 $\beta$  and induces hyperresponsiveness of the human isolated bronchus. *Eur Respir J* 2005; 26: 15–20.
- 34 Tortorolo L, Langer A, Polidori G, *et al.* Neurotrophin overexpression in lower airways of infants with respiratory syncytial virus infection. *Am J Respir Crit Care Med* 2005; 172: 233–237.
- 35 Ricci A, Graziano P, Bronzetti E, *et al.* Increased pulmonary neurotrophin protein expression in idiopathic interstitial pneumonias. *Sarcoidosis Vasc Diffuse Lung Dis* 2007; 24: 13–23.
- 36 Matsuda H, Koyama H, Sato H, *et al.* Role of nerve growth factor in cutaneous wound healing: accelerating effects in normal and healing-impaired diabetic mice. *J Exp Med* 1998; 187: 297–306.
- 37 Matsuda H, Koyama H, Sato H, *et al.* Role of nerve growth factor in cutaneous wound healing: accelerating effects in normal and healing-impaired diabetic mice. *J Exp Med* 1998; 187: 297–306.
- 38 Lambiase A, Manni L, Bonini S, *et al.* Nerve growth factor promotes corneal healing: structural, biochemical, and molecular analyses of rat and human corneas. *Invest Ophthalmol Vis Sci* 2000; 41: 1063–1069.
- 39 Holgate ST, Lackie P, Wilson S, *et al.* Bronchial epithelium as a key regulator of airway allergen sensitization and remodeling in asthma. *Am J Respir Crit Care Med* 2000; 162: S113–S117.
- 40 Dagnell C, Kemi C, Klominek J, *et al.* Effects of neurotrophins on human bronchial smooth muscle cell migration and matrix metalloproteinase-9 secretion. *Transl Res* 2007; 150: 303–310.
- 41 Freund-Michel V, Bertrand C, Frossard N. TrkA signalling pathways in human airway smooth muscle cell proliferation. *Cell Signal* 2006; 18: 621–627.
- 42 Micera A, Vigneti E, Pickholtz D, *et al.* Nerve growth factor displays stimulatory effects on human skin and lung fibroblasts, demonstrating a direct role for this factor in tissue repair. *Proc Natl Acad Sci USA* 2001; 98: 6162–6167.