

All comers

BEYER-WESTENDORF *et al.* 2013 [19][#]

5.2 (3.5–7.7)

100

Survivors

RIBEIRO *et al.* 1999 [45][#]

5.1 (2.0–12.5)

24.5

OTERO *et al.* 2011 [41][#]

1.3 (0.7–2.4)

42.0

CHOI *et al.* 2014 [21]

6.1 (3.7–9.8)

33.5

Subtotal ($I^2 = 81.9\%$)

3.85 (0.16–7.54)

100

Survivors without major comorbidities

DENTALI *et al.* 2009 [23]

8.8 (4.5–16.4)

11.1

GOLPE *et al.* 2010 [9][#]

11.1 (6.5–18.4)

11.0

VANNI *et al.* 2010 [49]

3.4 (1.9–6.1)

14.4

GARY *et al.* 2012 [26][#]

3.0 (1.0–8.5)

13.4

KORKMAZ *et al.* 2012 [37]

6.5 (4.2–9.9)

13.9

ABUL *et al.* 2014 [15]

7.9 (4.9–12.4)

13.2

TOSUN *et al.* 2014 [48][#]

40.2 (31.6–49.4)

8.1

PESAVVENTO *et al.* 2015 [43]

1.4 (0.8–2.6)

14.9

Subtotal ($I^2 = 93.0\%$)

8.52 (4.68–12.35)

100

