



# Chronic breathlessness: re-thinking the symptom

*To the Editor:*

Is chronic breathlessness a symptom or a syndrome? Although hotly debated [1–5], we would like to suggest an alternative viewpoint. Here, we speculate that the argument over chronic breathlessness being considered as either a symptom or syndrome both holds us within a reductionist framework, and somewhat misses the point.

The approach of traditional medicine has been to identify and target the pathophysiology thought to underlie symptoms, and thus treat a disease. However, more modern neuroscientific approaches have taken a significant leap forward within our understanding of perceptual systems. First examined within the conventional, exteroceptive senses (vision, touch, *etc.*), the “Bayesian brain hypothesis” outlines how perception occurs as a result of a delicate balance between incoming sensory information and the brain’s predictions about the world around us, based on learned experiences (priors) (figure 1). This hypothesis has recently been applied with vigour toward a more abstract set of perceptions, which consider the monitoring of our internal sensations (termed interoception [6–8]), such as breathing [9, 10].

Thus, this framework emboldens us for a more comprehensive appreciation of the important point raised by CALVERLEY [5]; namely, understanding what illness means to the patient and how it affects their lives. Each individual brings their own set of prior expectations, interoceptive abilities and bodily awareness [11, 12] and, thus, breathlessness will be both vastly quantitatively and qualitatively different between individuals. The breadth of these differences cannot hope to be enumerated into a narrow, homogenous set of symptoms that fall within a customary definition of a “syndrome”, but that does not need to be what JOHNSON *et al.* [1] were trying to achieve. The point, instead, is to view breathlessness as something that might not directly correspond to airway pathophysiology, and which may need to be treated both in parallel and independently of the lungs.

Furthermore, this theoretical view of breathlessness does not simply stop at lung disease. Whilst descriptive differences are often apparent between individuals who experience breathlessness in health, lung disease and other breathlessness-associated diseases, there is unlikely to be a hard qualitative boundary that exists between these (sometimes transient) health statuses. The wealth of experiences and prior expectations brought to the table by even healthy individuals will evoke a spectrum of breathlessness perceptions, and may even alter an individual’s propensity towards symptom severity and debilitation, should they ever become associated with disease. Therefore, attempts to quantify even the intensity of breathlessness as a more purely physiological perception determined by neural respiratory drive, whilst appealing within a traditional reductionist framework, appears to be a significant oversimplification that moves us further from understanding breathlessness within an individual.

So where do we go from here? How do we tackle these seemingly infinite degrees of freedom and dimensions of breathlessness within an individual? Whilst phrenology was popular within the Victorian era, where different parts of the brain were assigned different functions, we must now update our methodology to equip us to test these current theories. Modern neuroimaging techniques allow us to envision cortical activity that may flow from areas such as primary motor/sensory cortices and posterior insula, relevant for primary signal transduction of afferent inputs, but we must tread carefully in labelling this “conscious perception”. Instead, perception (and indeed breathlessness) is likely to be an evolving



@ERSpublications

**Chronic breathlessness, symptom or syndrome: are we asking the right question?**

<http://ow.ly/qJSX30hRpFq>

**Cite this article as:** Faull OK, Marlow L, Finnegan SL, *et al.* Chronic breathlessness: re-thinking the symptom. *Eur Respir J* 2018; 51: 1702238 [<https://doi.org/10.1183/13993003.02238-2017>].




FIGURE 1 Schematic of the Bayesian brain hypothesis. Both prior expectations and incoming sensory information contribute to the resulting perception, where each is a distribution of possible values. Thus, both the precision and position of priors and sensory inputs can influence perception.

function embedded within dynamic brain networks, where transduced sensory inputs are continuously compared to the brain's model of the world. Thus, linear increases in activity within one brain area is not likely to produce a corresponding linear change in breathlessness, and we need to equip ourselves with appropriate computational strategies [13, 14] to tackle these more difficult, more multi-dimensional brain network models.

Therefore, the suggestion from CAMPBELL and HOWELL [15] that “a respiratory physiologist offering a unitary explanation for breathlessness should arouse the same suspicions as a tattooed archbishop offering a free ticket to heaven”, appears to have been a foreseen and apt word of caution to our modern neuroscientific selves. Indeed, to understand breathlessness we need to delve into the brain more deeply, requiring an understanding of the networks that generate perception, and not just correlated activity in single brain areas. In turn, this may lead us to identify important treatable traits relating to breathlessness, complementing periphery-based therapies across a host of diseases and disorders, and hopefully better address the holistic effect of an illness on the life of an individual.

**Olivia K. Faull, Lucy Marlow, Sarah L. Finnegan and Kyle T.S. Pattinson**

Breathe Oxford Research Group, Nuffield Dept of Clinical Neurosciences, University of Oxford, Oxford, UK.

Correspondence: Kyle T.S. Pattinson, Breathe Oxford Research Group, Nuffield Dept of Clinical Neurosciences, University of Oxford, John Radcliffe Hospital, Oxford, OX3 9DU, UK. E-mail: kyle.pattinson@nda.ox.ac.uk

Received: Oct 31 2017 | Accepted: Nov 02 2017

Conflict of interest: Disclosures can be found alongside this article at [erj.ersjournals.com](http://erj.ersjournals.com)

## References

- 1 Johnson MJ, Yorke J, Hansen-Flaschen J, *et al.* Towards an expert consensus to delineate a clinical syndrome of chronic breathlessness. *Eur Respir J* 2017; 49: 1602277.
- 2 Johnson MJ, Yorke J, Hansen-Flaschen J, *et al.* Breathlessness despite optimal pathophysiological treatment: on the relevance of being chronic. *Eur Respir J* 2017; 50: 1701297.
- 3 Calverley PMA. Breathlessness despite optimal pathophysiological treatment: on the relevance of being chronic. *Eur Respir J* 2017; 50: 1701376.
- 4 Morélot-Panzini C, Adler D, Aguilaniu B, *et al.* Breathlessness despite optimal pathophysiological treatment: on the relevance of being chronic. *Eur Respir J* 2017; 50: 1701159.
- 5 Calverley PMA. Chronic breathlessness: symptom or syndrome? *Eur Respir J* 2017; 49: 1700366.
- 6 Seth AK. Interoceptive inference, emotion, and the embodied self. *Trends Cogn Sci* 2013; 17: 565–573.
- 7 Stephan KE, Manjaly ZM, Mathys CD, *et al.* Allostatic self-efficacy: a metacognitive theory of dyshomeostasis-induced fatigue and depression. *Front Hum Neurosci* 2016; 10: 49–27.
- 8 Van den Bergh O, Witthöft M, Petersen S, *et al.* Symptoms and the body: taking the inferential leap. *Neurosci Biobehav Rev* 2017; 74: 185–203.
- 9 Herigstad M, Faull OK, Hayen A, *et al.* Treating breathlessness via the brain: changes in brain activity over a course of pulmonary rehabilitation. *Eur Respir J* 2017; 50: 1701029.
- 10 Faull OK, Hayen A, Pattinson KTS. Breathlessness and the body: neuroimaging clues for the inferential leap. *Cortex* 2017; 95: 211–221.
- 11 Garfinkel SN, Seth AK, Barrett AB, *et al.* Knowing your own heart: distinguishing interoceptive accuracy from interoceptive awareness. *Biol Psychol* 2015; 104: 65–74.
- 12 Garfinkel SN, Manassei MF, Hamilton-Fletcher G, *et al.* Interoceptive dimensions across cardiac and respiratory axes. *Philos Trans R Soc Lond B Biol Sci* 2016; 371: 20160014.
- 13 Stephan KE, Friston KJ. Analyzing effective connectivity with functional magnetic resonance imaging. *WIREs Cogn Sci* 2010; 1: 446–459.

- 14 Petzschner FH, Weber LAE, Gard T, *et al.* Computational psychosomatics and computational psychiatry: toward a joint framework for differential diagnosis. *Biol Psychiatry* 2017; 1–10.
- 15 Campbell EJM, Howell JBL. The sensation of dyspnoea. *Br Med J* 1963; 2: 868.

Copyright ©ERS 2018

*From the authors:*

We agree with much presented in O.K. Faull and colleagues' response to our proposed chronic breathlessness syndrome and thank them for their views supporting chronic breathlessness as more than just a symptom [1]. In particular, we agree that naming and defining chronic breathlessness as a syndrome does not aim, or need, to constrain a recognition or understanding of quantitative and qualitative between-individual differences. We agree that our intention is to ensure patients and their families living with the daily disability affecting all domains of life, irrespective of cause, may legitimately bring this experience to the attention of health and social professionals, receive a systematic assessment and subsequently benefit from individually tailored evidence-based management.

There are two issues for comment. Firstly, O.K. Faull and colleagues appear to equate breathlessness with primarily a sensory, effective and cognitive perception. However, the foremost burden of chronic breathlessness is not sensory, but rather the functional confinement that breathlessness imposes on an individual. In chronic breathlessness, affective and cognitive perceptions relate more to loss of function, socialisation and fulfilment, and to fear of suffocation and death. Chronic breathlessness syndrome highlights that the impact of this condition reaches well beyond the unpleasant perceptions of, and emotional response to, a symptom into the realm of functional impairment and disability. This concept is illustrated well in recently published practical clinical assessment and management approaches to chronic breathlessness, the Breathing, Thinking, Functioning and Breathing SPACE models [2, 3].

Secondly, the use of the term reductionist is unhelpful as implied by O.K. Faull and colleagues, that is, a syndrome attempts the impossible; a complex idea cannot be completely understood in terms of its components and an attempt to do so is foolish. However, reductionist may also be used in the sense of analysing "complex things into less complex constituents" [4]. "Breathlessness persists despite optimal treatment and causes disability" [1] for many, yet remains almost totally invisible to most health and social care professionals. Therefore chronic breathlessness needs an identifiable profile in order to bring patient experience into view, help them access evidence-based interventions and stimulate research into the complexity of breathlessness in order to identify future therapeutic targets. Already, the work of O.K. Faull and colleagues has increased our understanding that perception of chronic breathlessness (experienced by patients) compared with that in healthy volunteer models is different; but much more research is needed [5]. If reducing the complex to recognisable components in clinical practice helps clinicians to routinely assess patients, helps patients have their concerns heard, helps carers to seek support and understanding, and helps services to develop and research to grow, then the reduction of chronic breathlessness to a syndrome will have made a major difference to the lives of many.

**Miriam J. Johnson<sup>1</sup>, Janelle Yorke<sup>2</sup>, John Hansen-Flaschen<sup>3</sup>, Robert Lansing<sup>4</sup>, Magnus Ekström<sup>5</sup> and David C. Currow<sup>1,6</sup>**

<sup>1</sup>Wolfson Palliative Care Research Centre, Hull York Medical School, University of Hull, Hull, UK. <sup>2</sup>School of Nursing, Midwifery and Social Work, University of Manchester, Manchester, UK. <sup>3</sup>Perelman School of Medicine, Hospital of the University of Pennsylvania, Philadelphia, PA, USA. <sup>4</sup>Beth Israel Hospital Harvard Medical School, Boston, MA, USA. <sup>5</sup>Dept of Clinical Sciences, Division of Respiratory Medicine & Allergology, Lund University, Lund, Sweden. <sup>6</sup>Faculty of Health, University of Technology Sydney, Sydney, Australia.

Correspondence: Miriam J. Johnson, Allam Medical Building, Hull York Medical School, University of Hull, Hull, HU6 7RX, UK. E-mail: miriam.johnson@hyms.ac.uk

Received: Nov 20 2017 | Accepted: Nov 21 2017

Conflict of interest: Disclosures can be found alongside this article at [erj.ersjournals.com](http://erj.ersjournals.com)



@ERSpublications

**Chronic breathlessness reaches beyond symptom perception into the realm of functional impairment and disability** <http://ow.ly/IBRd30h1wmS>

**Cite this article as:** Johnson MJ, Yorke J, Hansen-Flaschen J, *et al.* Chronic breathlessness: re-thinking the symptom. *Eur Respir J* 2018; 51: 1702326 [<https://doi.org/10.1183/13993003.02326-2017>].

## References

- 1 Johnson MJ, Yorke J, Hansen-Flaschen J, *et al.* Towards an expert consensus to delineate a clinical syndrome of chronic breathlessness. *Eur Respir J* 2017; 49: 1602277.
- 2 Spathis A, Booth S, Moffat C, *et al.* The Breathing, Thinking, Functioning clinical model: a proposal to facilitate evidence-based breathlessness management in chronic respiratory disease. *NPJ Prim Care Respir Med* 2017; 27: 27.
- 3 Hopkinson NS, Baxter N. Breathing SPACE – a practical approach to the breathless patient. *NPJ Prim Care Respir Med* 2017; 27: 5.
- 4 Collins English Dictionary – Complete and Unabridged 2012 Digital Edition. New York, Harper Collins, 2012.
- 5 Herigstad M, Hayen A, Evans E, *et al.* Dyspnea-related cues engage the prefrontal cortex: evidence from functional brain imaging in COPD. *Chest* 2015; 148: 953–961.

Copyright ©ERS 2018



### To the Editor:

We wholly applaud the move by JOHNSON *et al.* [1] to improve awareness of breathlessness and to raise its profile as a subject for focussed clinical research. We consider their research and the ensuing proposal to recognise breathlessness *via* a new medical term, “chronic breathlessness syndrome”, as important and justified. We share their goal, which is to direct attention to this neglected, undertreated and under-researched symptom.

There are two important caveats to be made in response to this article, however. First, there is a need to involve those who live with chronic breathlessness and are thus “experts by experience” in discussions about the framework proposed here, rather than bringing them into the conversation once consensus has been achieved. Second, further medicalisation of breathlessness *via* the term “syndrome” may not be the best way forward. Research into patients’ experience of breathlessness shows that the ways in which breathlessness is spoken about (medicalised and otherwise) not only reflect their experiences but also helps to shape how breathlessness is lived [2–6].

O.K. Faull and colleagues, in their response to JOHNSON *et al.* [1], comment on the individuality of responses to breathlessness that rely on prior experiences and bodily awareness (interoception). Context and culture play an important role in shaping the understanding and perception of breathlessness [2–4, 6]. For example, among African American communities across the USA, the last words of Eric Garner, “I can’t breathe”, as he suffocated in a tussle with police officers, have become a slogan for the Black Lives Matter movement and a metaphor for the lives of those living under other kinds of oppression [2]. BAŞOĞLU [7] suggests that asphyxiation is the most traumatic form of torture and that persistent breathlessness because of an underlying medical condition may be even worse due to the duration of the suffering involved.

Sufferers of respiratory illness vary in relation to the intensity, affect, ideation and meaning they attribute to their breathlessness [3–6]. It affects every aspect of the life of a breathless person in ways that description of it as a medical symptom cannot capture in full [8]. There is a need to legitimise a range of attitudes towards breathlessness in order for them to inform the clinical encounter. Collecting such experiences under the umbrella term “syndrome” may not be sufficient to enable full expression of the variability and multiple meanings of the experience of breathlessness, and may carry unexpected cognitive and affective “baggage” that detracts from its utility as a proxy for experience.

In view of the highly contextualised experience of breathlessness, it is critical to think about whose views are part of the debate. Discussions with experts by experience and first-person reports of experiences of breathlessness [3, 5] have revealed how powerful language and context are in determining how people with breathlessness think about and experience their problem [4, 6], and how this influences what they might do. Words such as “pulmonary” and “rehabilitation”, for example, may negatively impact upon the uptake of one of the most effective interventions for breathlessness [6].



@ERSpublications

**People with breathlessness should be involved in renaming their condition as their experience can be influenced by clinical language** <http://ow.ly/ZKNj30hzUUM>

**Cite this article as:** Macnaughton J, Oxley R, Rose A, *et al.* Chronic breathlessness: re-thinking the symptom. *Eur Respir J* 2018; 51: 1702331 [<https://doi.org/10.1183/13993003.02331-2017>].

There is a further stage necessary in the research process of JOHNSON *et al.* [1] in order to validate the claim made in the paper that “a recognised syndrome would [...] give permission for patients to discuss their ongoing breathlessness with their clinicians”. As JOHNSON *et al.* [1] suggest, patients and their families need to be involved in the discussion, but they should be able to critique the framework suggested by the paper, rather than be presented with it as a *fait accompli*. Otherwise there is a danger that the words “chronic” and “syndrome” will drive people with breathlessness further underground, in part because they have not been involved in the process of describing their own condition [9]. We encourage JOHNSON *et al.* [1] to take this research on to its next logical stage, that of developing a truly consensual terminology that considers the critical role language, metaphor and meaning play in both living with and treating breathlessness. This could be done using the Delphi technique within a more participatory paradigm [10]. Such an approach offers the chance of empowering patients and caregivers in ways that would result in real changes to both their experience and treatment.

**Jane Macnaughton<sup>1,2,3</sup>, Rebecca Oxley<sup>1,2,3</sup>, Arthur Rose<sup>1,3,4</sup>, Andrew Russell<sup>3,5</sup>, James W. Dodd<sup>6,7,8</sup> and Havi Carel<sup>8,9</sup>**

<sup>1</sup>Centre for Medical Humanities, Durham University, Durham, UK. <sup>2</sup>Dept of Anthropology, Durham University, Durham, UK. <sup>3</sup>Life of Breath Project, Durham, UK. <sup>4</sup>Dept of English Studies, Durham University, Durham, UK. <sup>5</sup>Dept of Anthropology, Durham University, Durham, UK. <sup>6</sup>Academic Respiratory Unit, Southmead Hospital, Bristol, UK. <sup>7</sup>Respiratory Medicine, University of Bristol, Bristol, UK. <sup>8</sup>Life of Breath Project, Bristol, UK. <sup>9</sup>Dept of Philosophy, University of Bristol, Bristol, UK.

Correspondence: Jane Macnaughton, Centre for Medical Humanities, Durham University, Caedmon Hall, Leazes Road, Durham, DH1 1SZ, UK. E-mail: jane.macnaughton@durham.ac.uk

Received: Nov 12 2017 | Accepted: Nov 23 2017

Conflict of interest: Disclosures can be found alongside this article at [erj.ersjournals.com](http://erj.ersjournals.com)

## References

- 1 Johnson MJ, Yorke J, Hansen-Flaschen J, *et al.* Towards an expert consensus to delineate a clinical syndrome of chronic breathlessness. *Eur Respir J* 2017; 49: 1602277.
- 2 Rose A. Tim Winton's pneumatic materialism. *Interventions* 2017; in press.
- 3 Carel H. *Phenomenology of Illness*. Oxford, Oxford University Press, 2016.
- 4 Oxley R, Macnaughton J. Inspiring change: humanities and social science insights into the experience and management of breathlessness. *Curr Opin Support Palliat Care* 2016; 10: 256–261.
- 5 Carel H. *Illness: the cry of the flesh*. 2nd Edn. Durham, Acumen, 2013.
- 6 Oxley R. Lost in Translation? Exploring the Language of Breathlessness. Life of Breath blog. <https://lifeofbreath.org/2017/09/lost-in-translation-exploring-the-language-of-breathlessness/> Date last accessed: November 6, 2017. Date last updated: September 1, 2017.
- 7 Başoğlu M. Effective management of breathlessness: a review of potential human rights issues. *Eur Respir J* 2017; 49: 1602099.
- 8 Hayen A, Herigstad M, Pattinson KTS. Understanding dyspnea as a complex individual experience. *Maturitas* 2013; 76: 45–50.
- 9 Kidd IJ, Carel H. Epistemic injustice and illness. *J Appl Philos* 2016; 34: 172–190.
- 10 Kezar A, Maxey D. The Delphi technique: an untapped approach of participatory research. *Int J Soc Res Methodol* 2016; 19: 143–160.

Copyright ©ERS 2018

This article is open access and distributed under the terms of the Creative Commons Attribution Licence 4.0