Version 1.0 (12.11.18)

### Appendix 1 - Details of search strategies

## MEDLINE (Ovid)

- 1. exp Pneumonia/
- 2. exp Community-Acquired Infections/
- 3. "community-acquired pneumonia.tw.
- 4. exp "Outcome Assessment (Health Care)"/
- 5. exp "Recovery of Function"/
- 6. exp "Activities of Daily Living"/
- 7. exp Symptom Assessment
- 8. exp Treatment Outcome/
- 9. exp Patient Readmission/
- 10. symptom recovery.tw.
- 11. symptomatic recovery.tw.
- 12. functional recovery.tw.
- 13. 1 and 2
- 14. 3 or 13
- 15. OR/ 4-12
- 16. 14 and 15

Version 1.0 (12.11.18)

#### EMBASE (OVID)

- 1. exp pneumonia/
- 2. exp community acquired infection/
- 3. community acquired pneumonia\*.tw.
- 4. community-acquired pneumonia\*.tw.
- 5. 1 and 2
- 6. OR/ 3-5
- 7. exp outcome assessment/
- 8. recovery of function\*.mp. or exp convalescence/
- 9. exp daily life activity/
- 10. exp symptom assessment/
- 11. exp treatment outcome/
- 12. exp hospital readmission/
- 13. symptom recovery\*.tw.
- 14. symptomatic recovery\*.tw.
- 15. OR/ 7-14
- 16. 6 and 15

Version 1.0 (12.11.18)

#### AMED (OVID)

- 1. exp Pneumonia/
- 2. community-acquired infection\*.mp.
- 3. community-acquired pneumonia\*.mp.
- 4. community acquired pneumonia\*.mp.
- 5. OR/ 1-4
- 6. exp "recovery of function"/
- 7. exp "Activities of daily living"/
- 8. exp Symptoms/
- 9. exp Treatment outcome/
- 10. exp Rehabilitation/
- 11. patient readmission\*.mp.
- 12. functional recovery\*.mp.
- 13. OR/ 6-12
- 14. 5 and 13

#### Version 1.0 (12.11.18)

# Appendix 2; Online supplementary material; Table S1; Risk of bias in included studies

First Author	Primary Outcome	Selection Domain (max 3)	Outcome Domain (max 3)	Comments
Adamuz 2011	Additional healthcare utilisation within 30days of discharge	3	2	Outcome measure not clearly defined or justified
Adamuz 2015	Additional healthcare utilisation within 30days of discharge	3	3	
Bruns 2010	Rate of radiographic resolution	2	2	20.1% loss to follow-up
Carratala 2005	Successful outcome' - composite of 7 predefined criteria	3	3	Low severity CAP (PSI classes II and III only)
Daniel 2018	Healthcare re-consultation within 4 weeks of discharge	2	3	Working age adults, low severity CAP
El Moussaoui 2006	Pneumonia related symptoms during follow-up	2	2	Low severity CAP
Fine 1999	Not defined - process of care and outcome of outpatients and hospitalised patients with CAP	2	2	
Labarere 2007	30 day mortality rate	2	3	Low severity CAP
Marrie 2000	Proportion of patients with symptom resolution at 6 weeks	2	2	No details regarding loss to follow-up
Metlay 1998	Proportion of patients with symptom resolution at 4 weeks	2	2	
Nickler 2016	Decline in QoL from hospital admission (baseline) to day 30 and after 6 years.	2	2	Not representative sample excluded elderly, dementia, significant co-morbid disease
Sharma 2006	Decline in ADL score by at least 1 point, comparing baseline prehospitalization ADL to 1 and 6 months post discharge.	2	2	Not representative sample, from cohort study of delirium in elderly, no details on loss to follow-up
Uranaga 2016	Clinical success rate at day 10 and day 30 since admission	3	3	
Wootoon 2017	Time for symptomatic recovery to baseline	2	2	Excluded patients with cognitive impairment, no loss to follow-up information
Wyrwich 2015	Time for symptom resolution	0	2	