

**Electronic Annex 6: Interval between drug administration and adverse event occurrence, according to the treatment outcome at the study data collection.**

	Still on treatment	Other outcomes	Total
<b>Median (IQR) interval between first drug administration and AEs, days</b>			
<b>Linezolid</b>	71 (32-139)	115 (52-191)	93.5 (40-169)
<b>Bedaquiline</b>	71 (28-160)	94 (30-184)	90 (30-177)
<b>Capreomycin</b>	57 (32-62)	75 (34-176)	62 (32.5-150.0)
<b>Pyrazinamide</b>	61 (30-105)	69 (33-178)	62 (31-116)
<b>Ethionamide/prothionamide</b>	162.5 (48.5-235.5)	159.5 (25-305)	159.5 (30.5-281.5)
<b>Amikacin</b>	81.5 (46.5-94.5)	61 (45-109)	80 (45-106)
<b>Cycloserine/terizidone</b>	105 (55-291)	93.5 (43-153)	105 (44-215)
<b>Para-aminosalicilic acid</b>	114 (32.5-206.0)	258 (70-629)	216 (93-603)
<b>Delamanid</b>	121.5 (59-181)	76 (22.5-100.5)	77 (24-179)
<b>Levofloxacin</b>	57 (53-103)	17 (7-160)	57 (12-160)
<b>Clofazimine</b>	132 (130-132)	97 (83-161)	131 (83-161)
<b>Moxifloxacin</b>	70.5 (22.5-118.5)	119.5 (52-254)	88 (34-170)
<b>Clarithromycin</b>	-	-	-
<b>Mean (SD) total duration, days</b>	<b>92 (34.3)</b>	<b>102.9 (60.1)</b>	<b>79.5 (42.4)</b>

Legend: IQR: interquartile range; AE: adverse event; PAS: Para-aminosalicylic acid; SD: standard deviation