



SHAREABLE PDF

Impact of the SARS-CoV-2 epidemic on tuberculosis treatment outcome in Northern Italy

Paola Magro^{1,2,6}, Beatrice Formenti^{3,6}, Valentina Marchese^{2,4},
Maurizio Gulletta⁵, Lina Rachele Tomasoni⁵, Silvio Caligaris⁵,
Francesco Castelli^{1,2} and Alberto Matteelli^{1,2}

Affiliations: ¹Dept of Infectious and Tropical Diseases, University of Brescia, ASST Spedali Civili, Brescia, Italy. ²WHO Collaborating Centre for TB/HIV and TB elimination, Brescia, Italy. ³Dept of Infectious Diseases, ASST Spedali Civili of Brescia, Brescia, Italy. ⁴Specialist Consultation Service, ASST Spedali Civili of Brescia, Brescia, Italy. ⁵Unit of Tropical Diseases, ASST Spedali Civili, Brescia, Brescia, Italy. ⁶These authors contributed equally.

Correspondence: Beatrice Formenti, Dept of Infectious Diseases, ASST Spedali Civili of Brescia, 25125 Brescia, Italy. E-mail: beatriceformenti1992@gmail.com



@ERSpublications

It is important to recognise the impact of COVID-19 on TB care to mitigate its consequences. TB services have an urgent need to identify novel strategies to ensure the continuum of TB care at the time of the coronavirus pandemic. <https://bit.ly/2OIKnwL>

Cite this article as: Magro P, Formenti B, Marchese V, *et al.* Impact of the SARS-CoV-2 epidemic on tuberculosis treatment outcome in Northern Italy. *Eur Respir J* 2020; 56: 2002665 [<https://doi.org/10.1183/13993003.02665-2020>].

This single-page version can be shared freely online.

To the Editor:

We read with interest the letters by TADOLINI *et al.* [1] and STOCHINO *et al.* [2] in a recent issue of the *European Respiratory Journal*, which summarise the first available evidence on tuberculosis (TB) and coronavirus disease 2019 (COVID-19) [3].