





Oral iron supplementation with ferric maltol in patients with pulmonary hypertension

Karen M. Olsson^{1,2}, Jan Fuge ^{1,2}, Torben Brod¹, Jan C. Kamp ¹, Jan Schmitto³, Tibor Kempf⁴, Johann Bauersachs ⁴ and Marius M. Hoeper ^{1,2}

Affiliations: ¹Dept of Respiratory Medicine, Hannover Medical School, Hannover, Germany. ²German Centre of Lung Research (DZL), Germany. ³Dept of Cardiothoracic, Transplantation and Vascular Surgery, Hannover Medical School, Hannover, Germany. ⁴Dept of Cardiology and Angiology, Hannover Medical School, Hannover, Germany.

Correspondence: Karen M. Olsson, Dept of Respiratory Medicine, Hannover Medical School, 30623 Hannover, Germany. E-mail: olsson.karen@mh-hannover.de

@ERSpublications

Oral ferric maltol was well tolerated and effective in patients with pulmonary hypertension and iron deficiency anaemia. Restoration of iron stores corrected anaemia and resulted in improved right ventricular function and exercise tolerance https://bit.ly/2zMwvxc

Cite this article as: Olsson KM, Fuge J, Brod T, *et al.* Oral iron supplementation with ferric maltol in patients with pulmonary hypertension. *Eur Respir J* 2020; 56: 2000616 [https://doi.org/10.1183/13993003.00616-2020].

This single-page version can be shared freely online.

To the Editor:

In patients with pulmonary hypertension and iron deficiency anaemia, oral iron supplementation with ferric maltol was well tolerated and restored iron and haemoglobin.

Copyright ©ERS 2020.. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.