







Performance of prediction models for COVID-19: the Caudine Forks of the external validation

Glen P. Martin¹, Matthew Sperrin¹ and Giovanni Sotgiu ©²

Affiliations: ¹Division of Informatics, Imaging and Data Science, Faculty of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK. ²Clinical Epidemiology and Medical Statistics Unit, Dept of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy.

Correspondence: Giovanni Sotgiu, Clinical Epidemiology and Medical Statistics Unit, Dept of Medical, Surgical and Experimental Sciences, University of Sassari, via Padre Manzella, 4 Sassari 07100, Italy. E-mail: gsotgiu@uniss.it

@ERSpublications

Existing evidence suggests that none of the COVID-19 prediction models can be supported for clinical use. Here we discuss "what next" in COVID-19 prediction. https://bit.ly/2SMtoLV

Cite this article as: Martin GP, Sperrin M, Sotgiu G. Performance of prediction models for COVID-19: the Caudine Forks of the external validation. *Eur Respir J* 2020; 56: 2003728 [https://doi.org/10.1183/13993003.03728-2020].

This single-page version can be shared freely online.

Healthcare systems worldwide have observed significant changes to meet demands due to the coronavirus disease 2019 (COVID-19) pandemic. The uncertainty surrounding optimal treatment, the rapid public health urgency and clinical emergencies have caused a chaotic disruption of the cases and their related contacts at inpatient and outpatient settings. Developing more tailored healthcare plans based on the currently available scientific evidence, could help improve clinical efficacy, treatment outcomes, prognosis, and health efficiency.

Copyright ©ERS 2020. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.