



Prevalence of pulmonary embolism in patients with COVID-19 at the time of hospital admission and role for pre-test probability scores and home treatment

Mitja Jevnikar^{1,2,3}, Olivier Sanchez^{4,5,6,7}, Marc Humbert ^{1,2,3} and Florence Parent^{1,2,3}

¹Université Paris-Saclay, Faculty of Medicine, Le Kremlin-Bicêtre, France. ²INSERM UMR_S 999, Le Kremlin-Bicêtre, France. ³AP-HP, Service de Pneumologie et soins intensifs respiratoires, Hôpital Bicêtre, Le Kremlin-Bicêtre, France. ⁴AP-HP, Service de Pneumologie et Soins Intensifs, Hôpital Européen Georges Pompidou, Paris, France. ⁵INSERM UMR-S 1140, Paris, France. ⁶INNOVTE, St-Etienne, France. ⁷Université Paris Descartes, Faculty of Medicine, Paris, France.

Florence Parent (florence.parent@aphp.fr)



Shareable abstract (@ERSpublications)

There is a high prevalence of pulmonary embolism in patients with COVID-19 at the time of hospital admission despite a low level of clinical suspicion <https://bit.ly/3ungLrQ>

Cite this article as: Jevnikar M, Sanchez O, Humbert M, *et al.* Prevalence of pulmonary embolism in patients with COVID-19 at the time of hospital admission and role for pre-test probability scores and home treatment. *Eur Respir J* 2021; 58: 2101033 [DOI: 10.1183/13993003.01033-2021].

This single-page version can be shared freely online.

Copyright ©The authors 2021

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

Received: 9 April 2021
Accepted: 28 April 2021

Reply to A. Porfidia and co-workers:

