



Hyperventilation as one of the mechanisms of persistent dyspnoea in SARS-CoV-2 survivors

Justina Motiejunaite ^{1,2}, Pauline Balagny^{1,3}, Florence Arnoult¹, Laurence Mangin^{1,4}, Catherine Bancal¹, Emmanuelle Vidal-Petiot^{1,2,5}, Martin Flamant^{1,2,5}, Guillaume Jondeau^{2,6,7}, Alain Cohen-Solal^{2,8,9}, Marie-Pia d'Ortho^{1,2,10} and Justine Fria-Masson ^{1,2,10}

¹Service de Physiologie Clinique-Explorations Fonctionnelles, Assistance Publique Hôpitaux de Paris, Hôpital Bichat-Claude Bernard, Paris, France. ²Université de Paris, Paris, France. ³INSERM, Population-based Epidemiological Cohorts Unit, UMS 011, Paris, France. ⁴Laboratoire Matière et Système Complexes UMR 7057, CNRS, Paris, France. ⁵INSERM, U1149, Paris, France. ⁶Service de Cardiologie, Assistance Publique Hôpitaux de Paris, Hôpital Bichat-Claude Bernard, Paris, France. ⁷INSERM, U1148, Paris, France. ⁸Service de Cardiologie, Assistance Publique Hôpitaux de Paris, Hôpital Lariboisière, Paris, France. ⁹INSERM UMR-S 942, Paris, France. ¹⁰INSERM, UMR 1141 NeuroDiderot, Paris, France.

Justina Motiejunaite (justina.motiejunaite@aphp.fr)



Shareable abstract (@ERSpublications)

Inadequate exercise hyperventilation should not be overlooked while exploring the causes of exertional dyspnoea in SARS-CoV-2 survivors <https://bit.ly/3AxOiDh>

Cite this article as: Motiejunaite J, Balagny P, Arnoult F, *et al.* Hyperventilation as one of the mechanisms of persistent dyspnoea in SARS-CoV-2 survivors. *Eur Respir J* 2021; 58: 2101578 [DOI: 10.1183/13993003.01578-2021].

This single-page version can be shared freely online.

Copyright ©The authors 2021.

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

This article has an editorial commentary:
<https://doi.org/10.1183/13993003.01763-2021>

Received: 4 June 2021
Accepted: 28 June 2021

To the Editor:

There are increasing reports of persistent dyspnoea several months after the onset of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection [1]. In most cases, functional disability seems out of proportion compared to residual pulmonary function impairment [2]. To date, knowledge about the functional limitations following a SARS-CoV-2 infection remains limited.

