



Early View

Review

Predictors of progression in systemic sclerosis patients with interstitial lung disease

Oliver Distler, Shervin Assassi, Vincent Cottin, Maurizio Cutolo, Sonye K. Danoff, Christopher P. Denton, Jörg H.W. Distler, Anna-Maria Hoffmann-Vold, Sindhu R. Johnson, Ulf Müller-Ladner, Vanessa Smith, Elizabeth R. Volkmann, Toby M. Maher

Please cite this article as: Distler O, Assassi S, Cottin V, *et al.* Predictors of progression in systemic sclerosis patients with interstitial lung disease. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.02026-2019>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Predictors of progression in systemic sclerosis patients with interstitial lung disease

Authors: Oliver Distler¹, Shervin Assassi², Vincent Cottin³, Maurizio Cutolo⁴, Sonye K. Danoff⁵, Christopher P. Denton⁶, Jörg H.W. Distler⁷, Anna-Maria Hoffmann-Vold⁸, Sindhu R. Johnson⁹, Ulf Müller-Ladner¹⁰, Vanessa Smith^{11,12}, Elizabeth R. Volkmann¹³, Toby M. Maher^{14,15}

Affiliations: ¹Department of Rheumatology, University Hospital Zurich, Zurich, Switzerland; ²Department of Rheumatology and Clinical Immunogenetics, McGovern Medical School, University of Texas, Houston, Texas, USA; ³National Reference Center for Rare Pulmonary Diseases, Louis Pradel Hospital, Hospices Civils de Lyon, Claude Bernard University Lyon 1, UMR754, Lyon, France (ORCID: 0000-0002-5591-0955); ⁴Research Laboratory, Clinical Division of Rheumatology, Department of Internal Medicine DIMI, University of Genoa, IRCSS Polyclinic Hospital San Martino, Genoa, Italy; ⁵Division of Pulmonary and Critical Care Medicine, Johns Hopkins Medicine, Baltimore, Maryland, USA; ⁶Royal Free Hospital, London, UK; ⁷Department of Internal Medicine 3, University of Erlangen-Nuremberg, Erlangen, Germany; ⁸Department of Rheumatology, Oslo University Hospital, Oslo, Norway; ⁹Toronto Scleroderma Program, Department of Medicine, Toronto Western and Mount Sinai Hospitals, University of Toronto, Toronto, Canada; ¹⁰Department of Rheumatology and Clinical Immunology, Justus-Liebig University Giessen, Campus Kerckhoff, Bad Nauheim, Germany; ¹¹Department of Rheumatology, Ghent University Hospital, Ghent, Belgium; ¹²Department of Internal Medicine, Ghent University, Unit for Molecular Immunology and Inflammation, VIB Inflammation Research Center (IRC), Ghent, Belgium; ¹³Department of Medicine, Division of Rheumatology, University of California, David Geffen School of Medicine, Los

Angeles, California, USA; ¹⁴National Heart and Lung Institute, Imperial College, London, UK; ¹⁵NIHR Respiratory Clinical Research Facility, Royal Brompton Hospital, London, UK.

Corresponding author: Dr Oliver Distler, University Hospital Zurich, Department of Rheumatology, Gloriastr. 25, Zürich 8091, Switzerland. E-mail: Oliver.Distler@usz.ch

Take home message: Lung function tests and chest imaging help predict who has SSc-associated ILD and whether it will progress. In the absence of standardised methods for doctors, we recommend a strategy that combines both lung function tests and chest imaging.

Abstract

Systemic sclerosis (SSc) is a systemic autoimmune disease affecting multiple organ systems, including the lungs. Interstitial lung disease (ILD) is the leading cause of death in SSc.

There are no valid biomarkers to predict the occurrence of SSc-ILD, although auto-antibodies against anti-topoisomerase I and several inflammatory markers are candidate biomarkers that need further evaluation. Chest auscultation, presence of shortness of breath and pulmonary function testing are important diagnostic tools, but lack sensitivity to detect early ILD. Baseline screening with high-resolution computed tomography (HRCT) is therefore necessary to confirm an SSc-ILD diagnosis. Once diagnosed with SSc-ILD, patients' clinical courses are variable and difficult to predict, though certain patient characteristics and biomarkers are associated with disease progression. It is important to monitor patients with SSc-ILD for signs of disease progression, though there is no consensus about which diagnostic tools to use or how often monitoring should occur. In this article, we review methods used to define and predict disease progression in SSc-ILD. There is no valid definition of SSc-ILD disease progression, but we suggest that either a decline in forced vital capacity (FVC) from baseline of $\geq 10\%$, or an FVC decline of 5–9% in association with a decline in diffusing capacity of carbon monoxide of $\geq 15\%$ represents progression. An increase in the radiographic extent of ILD on HRCT imaging would also signify progression. A time period of 1–2 years is generally used for this definition, but a decline over a longer time period may also reflect clinically relevant disease progression.

Lay summary

Patients with systemic sclerosis can develop serious lung fibrosis. Predicting who will develop this and whether the disease will get worse helps patients receive the best treatments. Currently, there are no standard methods for doctors to follow to identify whether patients are getting worse. We suggest an approach that combines lung function testing, changes in shortness of breath and chest imaging to help monitor the course of the disease.

Introduction

Systemic sclerosis (SSc) is a systemic autoimmune disease that is characterised by endothelial dysfunction, resulting in small-vessel vasculopathy, immune dysregulation, fibroblast dysfunction and subsequent fibrosis; however, its detailed pathogenesis remains unclear [1]. Due to the heterogeneity of the disease, SSc represents a major clinical challenge for both physicians and patients [2]. In addition to disfiguring skin involvement, SSc can affect multiple organ systems, including the lungs [3]. The clinical course is variable, but organ manifestations frequently occur early in the disease [4]. SSc is associated with a 250% increase in mortality risk compared with healthy controls and, overall, this has not changed significantly in the last 40 years [5]. Mortality is primarily due to pulmonary complications: in the largest observational study conducted to date, the leading cause of death was interstitial lung disease (ILD, 17%), with pulmonary arterial hypertension (PAH) accounting for 15% of deaths [6, 7]. Furthermore, in a Norwegian cohort study, mortality in SSc correlated with the extent of lung fibrosis [8].

Between 1972 and 2002, the proportion of deaths due to scleroderma renal crisis declined from 42% to 6% of SSc-related deaths, likely due to identification of risk factors, prevention, the use of angiotensin-converting enzyme inhibitors to treat this dimension of the disease and a greater awareness of milder SSc cases less likely to develop renal crisis. In contrast, over the same period, the proportion of SSc deaths due to ILD increased from 6% to 33% [9], and in recent interventional studies, respiratory failure due to ILD has been reported to account for 43% of deaths [10]. This increase in the proportion of deaths due to ILD may reflect decreases in other causes of deaths. The overall 10-year survival improved from 54% to 66% from 1972 to 2002 [9].

Patients with SSc are routinely divided into limited cutaneous (lcSSc) and diffuse cutaneous (dcSSc) subsets based on the extent of skin fibrosis [11, 12]. Pulmonary involvement can occur in both subsets of the disease, and it can affect all aspects of the respiratory tract, including the parenchyma, vasculature and musculature [8, 13]. ILD is an early complication in SSc, and in some patients (approximately 4%) the first clinical symptom of SSc is directly related to ILD [2, 14]. Most patients who develop severe restrictive lung disease do so in the first 5 years following the onset of SSc symptoms [2].

Despite the established relationship between SSc-associated ILD (SSc-ILD) and morbidity and mortality, there is still no consensus on screening for ILD, nor on monitoring for disease progression. This issue is further complicated by the lack of validated biomarkers for SSc-ILD and an absence of clinical recommendations to inform the method and timing of investigations to diagnose patients with SSc-ILD and identify those at risk of progression [15].

In this perspective piece, we summarise the current understanding of disease pathogenesis and risk factors for the presence of ILD in patients with SSc, discuss screening, early detection of ILD and risk factors for progression, and propose guidance on monitoring disease progression and the implications of this for treating SSc-ILD.

Pathogenesis of SSc-ILD

SSc may affect multiple organs and systems, with most irreversible damage due to progressive fibrosis [2, 16]. Based on research to date, it is likely that there is a complex interplay between innate and adaptive immunity, endothelial dysfunction, small vessel abnormalities, inflammation and fibrosis [17]. The key pathways

implicated in SSc-ILD are summarised in Figure 1. The most common initiator of the pathogenesis of SSc-ILD is injury to alveolar epithelial and vascular endothelial cells, in which inflammatory pathways lead to pro-fibrotic stimuli that induce the differentiation of lung fibroblasts to the myofibroblast phenotype [6, 18]. Fibrosis results from an abnormal healing process coupled with an aberrant expansion of mesenchymal cells that originate from multiple sources including: bone marrow, the transdifferentiation of pericytes, epithelial cells and endothelial cells, and the differentiation from resident fibroblasts to myofibroblasts [16, 18]. Transforming growth factor-beta is thought to play a central role in the development and progression of fibrosis in SSc-ILD [6, 16] and platelet-derived growth factor, Wnt- β -catenin and hedgehog signalling also drive fibrosis [18, 19]. SSc fibroblasts have been shown to be more resistant to the fibrotic-inhibitory potential of cytokines, such as interferon gamma [20]. Furthermore, patients with SSc-ILD can have increased levels of pro-inflammatory cytokines such as interleukin (IL)-8, IL-6, tumour necrosis factor-alpha and macrophage inflammatory protein-1alpha in bronchoalveolar lavage (BAL) fluid and serum [21-23]. It has also been hypothesised that B-cells may be involved, as patients with SSc-ILD more frequently present with anti-topoisomerase I antibodies and antifibroblast antibodies [24]. The functional role of antifibroblast antibodies has not been clarified, but they might directly affect extracellular matrix remodelling by binding to fibroblasts to increase the collagen breakdown and matrix metalloproteinase-1 production [25]. The increased release of these cytokines, growth factors and bioactive proteins contribute to a self-perpetuating cycle of stimulation of the fibrotic process [16, 18]. A greater understanding of the underlying mechanisms may enable the identification of biomarkers that could be used to screen at-risk patients and facilitate the development of targeted and personalised

therapy. For instance, higher-circulating mixed M1 (classically activated)/M2 (alternatively activated) monocyte/macrophage cell percentages are associated with ILD, suggesting a need for research on their possible role as a biomarker or pathogenic element for SSc lung involvement [26].

Risk factors for presence of ILD

Estimates of the prevalence of ILD in SSc vary depending on how ILD is defined; using high-resolution computed tomography (HRCT), prevalence is estimated at 47–84%. In the Canadian Scleroderma Research Group registry, 65% of patients had evidence of ILD on HRCT scans, 26% had ILD identified by physical examination and 22% identified by chest radiography [27]. In a Norwegian cohort, 50% of patients had ILD on HRCT [8]. Relying on pulmonary function testing (forced vital capacity [FVC] % predicted or a decrease in at least one pulmonary function test [PFT]) results in a lower estimate of prevalence of ILD [28]. At autopsy, evidence of interstitial fibrosis was found in 74% of patients with SSc [29].

A number of patient characteristics are associated with a higher risk of developing ILD in SSc (Table 1). In the EUSTAR database, ILD was more common in dcSSc (53.4%) than in lcSSc (34.7%) [30]. Sex and ethnicity also appear to be important determinants of ILD development. Despite the strong female predominance reported in SSc (female to male ratio 4.7:1), men with SSc are more likely to develop ILD than women with SSc (relative risk 1.24; 95% confidence interval 1.01–1.52) [31, 32]. A large multi-ethnic observational study conducted in the setting of a universal healthcare system found that in comparison to white patients of European descent, patients of Afro-Caribbean descent with SSc more frequently have ILD (31% versus 53%; $p=0.007$) [33].

Risk factors for the development of SSc-ILD also include anti-topoisomerase I positivity, lower FVC and lower diffusing capacity of the lung for carbon monoxide (DL_{CO}) [34], although FVC and DL_{CO} changes may be caused by ILD and not present before ILD onset. Multivariate analyses suggest that the risk of developing ILD is more closely associated with anti-topoisomerase I positivity than to dcSSc versus lcSSc [30].

A number of studies also support an association of gastroesophageal reflux with ILD in SSc, as reviewed by Christmann RB et al. [35].

Diagnosis and screening of ILD

Diagnosis of ILD

The most common initial symptoms of SSc-ILD are exertional dyspnoea and dry cough, although early SSc-ILD can be relatively asymptomatic [6]. As lung involvement can affect all aspects of the respiratory tract, when a patient with SSc presents with symptoms of dyspnoea, the differential diagnosis can be quite broad and may include PAH, deconditioning, chronic anaemia [13], and left heart involvement with reduced or preserved ejection fraction.

SSc-ILD is diagnosed using HRCT to detect parenchymal lung disease [36].

Typically, a pattern suggestive of nonspecific interstitial pneumonia (NSIP) appears on HRCT as bilateral ground-glass opacities, reticulation and traction bronchiectasis most prominent in the lower lobes (Figure 2A) [36]. Generally, HRCT appearance can differentiate NSIP and usual interstitial pneumonia (UIP) patterns [37]. Although lung biopsy is very rarely indicated to diagnose SSc-ILD, if performed, interstitial fibrosis with temporal homogeneity and a modest inflammatory cell infiltrate (i.e. fibrotic NSIP) is typically visualised (Figure 2B). In a small number of cases, cellular

NSIP or UIP are observed [13]. Although computed tomography (CT) scans are necessary for diagnosis, they expose patients to radiation [38]. Reduced-dose HRCT protocols may reliably detect SSc-ILD with a much lower radiation exposure than whole-chest HRCT [38, 39]. While these reduced-dose protocols have become routine in some centres, further validation in larger cohorts is required.

PFTs in patients with SSc-ILD generally show a restrictive ventilatory defect with a decreased FVC and/or total lung capacity, a preserved forced expiratory volume in 1 second (FEV_1), and a normal or increased FEV_1/FVC ratio, as well as a decreased single-breath DL_{CO} [6]. It is important to note that patients with early SSc-ILD may have preserved lung volumes, despite clear evidence of structural lung disease on HRCT [28, 40], and decreases in DL_{CO} . Two studies found that over 60% of patients with SSc and ILD diagnosed using HRCT had normal spirometry [28, 40]. Therefore, while PFTs are an important part of the diagnostic work-up for SSc-ILD, they are not sensitive enough to reliably detect early ILD [28, 40, 41]. Performing PFTs regularly (at least once a year) in the first 5 years following SSc diagnosis may be useful for detecting changes in lung function indicative of ILD.

At present, there is no widely accepted consensus on when SSc-ILD becomes clinically significant and when treatment is indicated.

Challenges in diagnosis

There are many challenges associated with the diagnosis of SSc-ILD. One potential barrier to diagnosis is a lack of awareness within primary care of SSc, which can lead to late referrals [42]. There is also variability in the degree of involvement of clinical specialties in establishing a diagnosis of SSc-ILD, and a lack of formally designated SSc specialist centres [42].

Early detection of ILD

There is a need for early systematic screening of patients to identify pulmonary complications, though there is a lack of consensus regarding which tests should be used during screening and follow-up [2, 42]. In one study, screening for ILD at diagnosis of SSc using HRCT identified 35% of subjects with no ILD, 51% of subjects with ILD of limited extent (1–20% fibrosis), and 13% of subjects with ILD involving >20% of the lungs [43].

Subclinical interstitial lung abnormalities (ILAs) detectable on CT scans also offer a possible approach to detect ILD early [44, 45]. ILAs have been studied in rheumatoid arthritis-associated ILD, where they are associated with increased cough and dyspnoea and decreased PFT results [46]; however, there is little evidence characterising them in SSc-ILD. One study in early SSc-ILD demonstrated ground-glass opacities that later progressed to NSIP pattern [47].

Future directions in the diagnosis and screening of SSc-ILD

In future, lung ultrasound may be integrated with other approaches in the screening of ILD [48-51]. Preclinical studies of positron emission tomography scanning suggest that targeted nuclear imaging of components of ILD disease processes (such as integrin alpha-v-beta-3 and somatostatin receptor 2) could be integrated with other approaches in the screening and evaluation of ILD [52, 53]. However, this method is limited by the high levels of radiation exposure associated with the technique.

Another potential approach to help identify patients with ILD is nailfold videocapillaroscopy (NVC). NVC is a safe and validated tool, and NVC patterns are associated with different SSc clinical complications [54, 55]. Recently, NVC was

tested in 287 patients with SSc to evaluate whether it could improve the detection of patients with high risk for cardiopulmonary involvement. In this analysis, NVC pattern was associated with the presence of ILD and heart/lung involvement, independent of specific anti-extractable nuclear antigen antibodies [56]. This might indicate that microangiopathy is an important cause of organ involvement. NVC may in future be incorporated, together with other parameters, in high-performance algorithms in the early detection of lung involvement in SSc.

However, despite identified risk factors for the development of ILD in patients with SSc, and newer techniques that may contribute to identify subpopulations with increased risk of developing ILD, currently none of the available techniques other than CT can detect ILD with appropriate sensitivity and specificity. Chest CT, therefore, remains the gold standard to screen for and diagnose SSc-ILD.

Risk factors for progression of ILD

Once ILD has been diagnosed in patients with SSc, it is important to identify which patients are likely to progress, because the natural history of SSc-ILD is variable. Some patients experience accelerated loss of lung function, and others progress slowly or exhibit stable disease [57, 58].

At present, there is no biomarker that predicts the course of SSc-ILD across multiple clinical cohorts [59]. Initial reports that BAL cellularity could predict mortality, benefit of immunosuppressive therapy, progression or progression-free survival [60, 61] have not been substantiated in studies examining larger cohorts [62]. However, several strategies are available to help identify patients at risk for the development and progression of SSc-ILD, and these should be considered as a first step in the management of patients with SSc (Table 2).

In clinical practice, the severity of SSc-ILD may be staged based on the “Goh criteria” whereby patients have “limited” or “extensive” disease based on HRCT clearly <20% or clearly >20%, respectively, with the use of an FVC threshold of 70% in indeterminate cases [41]. Baseline HRCT extent has been associated with decline in FVC % predicted [63] and, in a more recent study, extensive disease (>20%) was associated with a threefold increase in deterioration or death [64].

An evidence-based prediction model – SPAR – that combines peripheral capillary oxygen saturation (SpO₂) and arthritis history has recently been developed to predict progression of mild ILD [65]. In 215 patients with SSc and mild ILD, lower SpO₂ after a 6-minute walk test and the presence of arthritis at any point were independent predictors of ILD progression (defined as a relative decrease in FVC ≥15%, or FVC ≥10% combined with DL_{CO} ≥15% at 1-year follow-up) [65]. By combining both predictors, the derived SPAR model was able to increase the prediction rate from 25.5% to 91.7% and, ultimately, may enable risk stratification of patients with early, mild SSc-ILD [65].

A number of other risk factors for SSc-ILD disease progression have been identified. A large SSc-ILD cohort study demonstrated that FVC (hazard ratio [HR] 0.97; p=0.0009), DL_{CO} (HR 0.97; p=0.0004) and increasing age (HR 1.03; p=0.04) were significantly associated with mortality after adjusting for sex, body mass index, smoking status, serology, disease subtype and immunosuppressive use [66]. A recent analysis showed that a decline in FVC or DL_{CO} over 2 years was a better predictor of mortality than baseline FVC and DL_{CO} [10].

Auto-antibodies against the DNA-modifying enzyme topoisomerase I are frequently present in patients with dcSSc and may be associated with the faster progression of SSc-ILD [34, 67, 68]. Owing to the pathogenic role of the chemokine receptors

CXCR3 and CXCR4 in fibrosis, auto-antibodies against them have been investigated in patients with SSc [69]. Anti-CXCR3 and anti-CXCR4 antibody levels have been shown to be different among SSc subgroups compared with healthy subjects, and highest in diffuse SSc patients. Patients with SSc-ILD exhibited higher antibody levels than those without ILD; however, patients with deterioration of lung function (FVC % predicted <70%) showed lower anti-CXCR3/4 antibody levels compared with those with stable disease [69]. However, these biomarkers are not routinely available.

Several inflammatory markers have been investigated. Elevated serum chemokine ligand 18 is predictive of FVC decline [70, 71] and is a risk factor for death or progression of SSc-ILD [71-73]. Baseline C-reactive protein levels predict long-term progression of SSc-ILD, and higher levels are also associated with shorter survival [74] and a decline in PFTs [75]. Krebs von den Lungen-6 (KL-6) and surfactant protein-D (SP-D) are proteins predominantly expressed in the lung, and are associated with lung damage, fibrosis and inflammation [71, 76]. Serum KL-6 level is inversely correlated with DL_{CO} % predicted, and elevated levels are associated with end-stage lung disease and faster FVC decline [71, 76-78], even in the presence of treatment with immunosuppression [79]. Further research is required to validate these biomarkers.

Monitoring disease progression

The variable rates of disease progression in SSc-ILD mean that it is important that patients are appropriately monitored after a diagnosis of ILD in SSc, although there is currently no established definition for disease progression. Serial measurements of symptoms (e.g. appearance or aggravation of dyspnoea, cough and/or fatigue),

physical examination and PFTs are important to assess disease progression at regular clinic visits [6, 21], and recent studies highlight the impact of change in PFT on mortality [10, 80].

Patient-reported outcomes may also be important for measuring disease progression; there is an association between lower FVC and worsening scores on St. George's Respiratory Questionnaire (SGRQ), a measure of respiratory-related health status, in patients with ILD [81]. It should be noted that this study largely evaluated patients with idiopathic pulmonary fibrosis, hypersensitivity pneumonitis and nonspecific interstitial pneumonia; SGRQ scores in SSc may be affected by comorbidities, including musculoskeletal problems.

Outstanding questions remain regarding the use of repeated HRCT, including how often it should be performed routinely (for instance, every year or every other year), and whether PFT or other tools should be used for monitoring, with HRCT only used if necessary (e.g. unexpected worsening or suspicion of infection) [38, 82]. In addition, there is some debate as to whether and how often CT scans should be repeated in patients with SSc who do not have an ILD when first examined.

SSc-ILD expert panel

An international panel of SSc-ILD experts from Europe and North America was assembled to discuss developments in the field. Topics of interest were selected by panel chairs and discussed in groups. The selected group of pulmonologists and rheumatologists identified predictors of progression of SSc-ILD as an area of unmet need. We reviewed evidence for the clinical course of SSc-ILD, examined the challenges involved in diagnosis, and provided guidance on the staging and risk stratification of patients with SSc-ILD.

We support the definition of progression of disease in individual patients as a sustained change in lung function characterised by an FVC decline from baseline of $\geq 10\%$, or an FVC decline of 5–9% in association with a DL_{CO} decline of $\geq 15\%$ (Figure 3) [80]. If serial CTs show an increase in the extent of ILD on HRCT imaging this would also be taken into account. This definition is one of the most commonly used definitions for progression, although it has not been formally validated in SSc-ILD. Researchers usually use this definition with a time period of 1 year, but there is no agreement on how disease progression should be defined over a longer time period. It should be acknowledged that this decline in FVC and/or DL_{CO} observed over a longer time period still demonstrates potentially clinically relevant disease progression, albeit at a slower rate.

Without intervention, the natural annual loss of lung function is 15–25 mL (inferred from regression equations in [83]). Diurnal variability in lung function is 8–9% and measurement error in clinical trials is about 1% [84]. In the worst-case scenario, this amounts to 10%. Thus, for a relative 5–9% loss in FVC, we agreed that it should be accompanied by a decline in DL_{CO} to confirm disease progression.

Implications for therapy

In the recently published SENSICIS[®] trial, patients with SSc-ILD treated with nintedanib had a lower rate of annual FVC decline than those receiving placebo (treatment difference 41 mL) [85]. Based on these data, nintedanib has recently been approved by the US Food and Drug Administration for slowing the rate of decline in pulmonary function in patients with SSc-ILD [86]. Guidelines issued by the European League Against Rheumatism and the EUSTAR group in 2017 recommend consideration of tailored cyclophosphamide (CYC) treatment in patients with SSc-

ILD, and the use of haematopoietic stem cell transplant in selected patients with rapidly progressing SSc [87]. Following the SLS II trial [10], a Delphi consensus treatment algorithm advocated for first line treatment of SSc-ILD with mycophenolate mofetil (MMF). Second-line treatments included CYC or rituximab as induction therapy, followed by MMF as maintenance therapy [88].

However, there is currently no consensus regarding treatment initiation or escalation [89]. Most SSc experts draw on their clinical experience and consider the risk factors for SSc-ILD progression to inform their decision on when to initiate therapy on a case-by-case basis [90]. Better identification of patients at greatest risk of progression will help to identify those who can benefit the most from early treatment [6]. There is a clear need for new treatments. As our understanding of the pathogenesis of SSc-ILD evolves, there is the hope that specific therapies will be introduced that treat the disease in the organ systems affected, improving patient well-being, function and survival [91-94]. Targeting the self-perpetuating fibrosis may provide therapeutic options in the future [16], although it remains unknown whether aggressive treatment of the disease early is beneficial in terms of modifying disease progression and preventing irreversible lung damage. In the near future, it is hoped that molecular detection of disease will inform immunomodulatory therapy in SSc patients [95].

Conclusions

By increasing awareness of frequency, progression and mortality associated with SSc-ILD among physicians, there is the potential to improve patient outcomes. Better identification of which patients are at risk of progression will also be beneficial. ILD is an early complication of SSc, and there is a need for early and systematic

screening of patients to potentially enable treatment prior to deterioration of lung function in subjects at high risk of disease progression. Pulmonary function testing remains an important diagnostic tool, but may need to be performed regularly following SSc diagnosis to detect changes suggesting ILD. Baseline screening with HRCT, as part of a wider clinical work-up that includes lung function testing, is necessary to confirm an SSc-ILD diagnosis. We favour a definition of disease progression as a sustained change in lung function characterised by an FVC decline from baseline of $\geq 10\%$, or an FVC decline of 5–9% in association with a DL_{CO} decline of $\geq 15\%$. This decline in FVC or DL_{CO} should be considered as disease progression even if it is only observed beyond the 12-month period. Patient-reported outcome tools focused on breathlessness and imaging may provide valuable supporting information. In the future, it is hoped that new imaging techniques and diagnostic and predictive biomarkers will play an increasingly important role in defining SSc-ILD progression.

Acknowledgements

The authors meet criteria for authorship as recommended by the International Committee of Medical Journal Editors. They take full responsibility for the scope, direction, content of, and editorial decisions relating to, the manuscript, were involved at all stages of development and have approved the submitted manuscript. The authors received no compensation related to the development of the manuscript. Boehringer Ingelheim International GmbH supported the meeting of the international panel of experts.

Dr. Johnson has been awarded a Canadian Institutes of Health Research New Investigator Award; and is supported by the Oscar and Eleanor Markovitz Fund for Scleroderma Research of the Arthritis Research Foundation.

Prof. Vanessa Smith is a Senior Clinical Investigator of the Research Foundation - Flanders (Belgium) (FWO) [1.8.029.15N]. The FWO was not involved in study design, collection, analysis and interpretation of data, writing of the report, nor in the decision to submit the article for publication.

TMM is supported by a National Institute for Health Research Clinician Scientist Fellowship (NIHR ref: CS-2013-13-017) and is a British Lung Foundation Chair in Respiratory Research (C17-3).

Funding sources

Writing, editorial support and formatting assistance was provided by Tom Priddle, DPhil and Claire Scofield, MRes, of Nucleus Global, which was contracted and funded by Boehringer Ingelheim International GmbH (BI). Boehringer Ingelheim International GmbH was given the opportunity to review the manuscript for medical and scientific accuracy as well as intellectual property considerations.

References

1. Varga J, Trojanowska M, Kuwana M. Pathogenesis of systemic sclerosis: recent insights of molecular and cellular mechanisms and therapeutic opportunities. *Journal of Scleroderma and Related Disorders* 2017; 2(3): 137-152.
2. Denton CP, Khanna D. Systemic sclerosis. *Lancet (London, England)* 2017; 390(10103): 1685.
3. Caron M, Hoa S, Hudson M, Schwartzman K, Steele R. Pulmonary function tests as outcomes for systemic sclerosis interstitial lung disease. *EurRespirRev* 2018; 27(148): pii 170102.
4. Jaeger VK, Wirz EG, Allanore Y, Rossbach P, Riemekasten G, Hachulla E, Distler O, Airo P, Carreira PE, Balbir Gurman A, Tikly M, Vettori S, Damjanov N, Muller-Ladner U, Distler JH, Li M, Walker UA, co-authors E. Incidences and risk factors of organ manifestations in the early course of systemic sclerosis: a longitudinal EUSTAR study. *PLoS One* 2016; 11(10): e0163894.
5. Elhai M, Meune C, Avouac J, Kahan A, Allanore Y. Trends in mortality in patients with systemic sclerosis over 40 years: a systematic review and meta-analysis of cohort studies. *Rheumatology (Oxford)* 2012; 51(6): 1017.
6. Volkmann ER, Tashkin DP. Treatment of systemic sclerosis-related interstitial lung disease: a review of existing and emerging therapies. *Annals of the American Thoracic Society* 2016; 13(11): 2045-2056.
7. Elhai M, Meune C, Boubaya M, Avouac J, Hachulla E, Balbir-Gurman A, Riemekasten G, Airo P, Joven B, Vettori S, Cozzi F, Ullman S, Czirjak L, Tikly M, Muller-Ladner U, Caramaschi P, Distler O, Iannone F, Ananieva LP, Hesselstrand R, Becvar R, Gabrielli A, Damjanov N, Salvador MJ, Riccieri V, Mihai C, Szucs G, Walker UA, Hunzelmann N, Martinovic D, Smith V, Muller CS, Montecucco CM, Opris D, Ingegnoli F, Vlachoyiannopoulos PG, Stamenkovic B, Rosato E, Heitmann S, Distler JHW, Zenone T, Seidel M, Vacca A, Langhe E, Novak S, Cutolo M, Mouthon L, Henes J, Chizzolini C, Muhlen CAV, Solanki K, Rednic S, Stamp L, Anic B, Santamaria VO, De Santis M, Yavuz S, Sifuentes-Giraldo WA, Chatelus E, Stork J, Laar JV, Loyo E, Garcia de la Pena Lefebvre P, Eyerich K, Cosentino V, Alegre-Sancho JJ, Kowal-Bielecka O, Rey G, Matucci-Cerinic M, Allanore Y, group E. Mapping and predicting mortality from systemic sclerosis. *AnnRheumDis* 2017; 76(11): 1897-1905.
8. Hoffmann-Vold A-M, Fretheim H, Halse A-K, Seip M, Bitter H, Wallenius M, Garen T, Salberg A, Brunborg C, Midtvedt Ø, Lund MB, Aaløkken TM, Molberg Ø. Tracking impact of interstitial lung disease in systemic sclerosis in a complete nationwide cohort. *American Journal of Respiratory and Critical Care Medicine* 2019; 200(10): 1258-1266.
9. Steen VD, Medsger TA. Changes in causes of death in systemic sclerosis, 1972-2002. *AnnRheumDis* 2007; 66(7): 940-944.
10. Volkmann ER, Tashkin DP, Sim M, Li N, Goldmuntz E, Keyes-Elstein L, Pinckney A, Furst DE, Clements PJ, Khanna D, Steen V, Schraufnagel DE, Arami S, Hsu V, Roth MD, Elashoff RM, Sullivan KM, groups SlsSls. Short-term progression of interstitial lung disease in systemic sclerosis predicts long-term survival in two independent clinical trial cohorts. *AnnRheumDis* 2019; 78(1): 122-130.
11. Fan MH, Feghali-Bostwick CA, Silver RM. Update on scleroderma-associated interstitial lung disease. *CurrOpinRheumatol* 2014; 26(6): 630-636.
12. LeRoy EC. Scleroderma (systemic sclerosis): classification, subsets and pathogenesis. *J Rheumatol* 1988; 15: 202.

13. Culp Silver K, Silver RM. Management of systemic-sclerosis-associated interstitial lung disease. *RheumDisClinNorth Am* 2015; 41(3): 439-457.
14. Rubio-Rivas M, Corbella X, Pestana-Fernandez M, Tolosa-Vilella C, Guillen-Del Castillo A, Colunga-Arguelles D, Trapiella-Martinez L, Iniesta-Arandia N, Castillo-Palma MJ, Saez-Comet L, Egurbide-Arberas MV, Ortego-Centeno N, Freire M, Vargas-Hitos JA, Rios-Blanco JJ, Todoli-Parra JA, Rodriguez-Carballeira M, Marin-Ballve A, Segovia-Alonso P, Pla-Salas X, Madronero-Vuelta AB, Ruiz-Munoz M, Fonollosa-Pla V, Simeon-Aznar CP, Rescle investigators AD SG, Callejas Moraga E, Calvo E, Carbonell C, Castillo MJ, Chamorro AJ, Colunga D, Egurbide MV, Espinosa G, Fonollosa V, Garcia Hernandez FJ, Gonzalez Leon R, Guillen Del Castillo A, Iniesta N, Lorenzo R, Madronero AB, Mari B, Marin A, Perez Conesa M, Pestana M, Pla X, Rios Blanco JJ, Rodriguez Carballeira M, Rubio Rivas M, Ruiz Munoz M, Saez Comet L, Segovia P, Simeon CP, Soto A, Tari E, Todoli JA, Tolosa C, Trapiella L, Vargas Hitos JA, Verdejo G. First clinical symptom as a prognostic factor in systemic sclerosis: results of a retrospective nationwide cohort study. *ClinRheumatol* 2018; 37(4): 999-1009.
15. Kennedy B, Branagan P, Moloney F, Haroon M, O'Connell OJ, O'Connor TM, O'Regan K, Harney S, Henry MT. Biomarkers to identify ILD and predict lung function decline in scleroderma lung disease or idiopathic pulmonary fibrosis. *Sarcoidosis VascDiffuse Lung Dis* 2015; 32(3): 228.
16. Bhattacharyya S, Wei J, Varga J. Understanding fibrosis in systemic sclerosis: shifting paradigms, emerging opportunities. *NatRevRheumatol* 2011; 8(1): 42.
17. Cappelli S, Bellando Randone S, Camiciottoli G, De Paulis A, Guiducci S, Matucci-Cerinic M. Interstitial lung disease in systemic sclerosis: where do we stand? *EurRespirRev* 2015; 24(137): 411.
18. Akter T, Silver RM, Bogatkevich GS. Recent advances in understanding the pathogenesis of scleroderma-interstitial lung disease. *CurrRheumatolRep* 2014; 16(4): 411.
19. Distler JHW, Györfi A-H, Ramanujam M, Whitfield ML, Königshoff M, Lafyatis R. Shared and distinct mechanisms of fibrosis. *Nature Reviews Rheumatology* 2019; 15(12): 705-730.
20. Chizzolini C, Rezzonico R, Ribbens C, Burger D, Wollheim FA, Dayer JM. Inhibition of type I collagen production by dermal fibroblasts upon contact with activated T cells: different sensitivity to inhibition between systemic sclerosis and control fibroblasts. *Arthritis Rheum* 1998; 41(11): 2039-2047.
21. Solomon JJ, Olson AL, Fischer A, Bull T, Brown KK, Raghu G. Scleroderma lung disease. *EurRespirRev* 2013; 22(127): 6.
22. De Lauretis A, Sestini P, Pantelidis P, Hoyles R, Hansell DM, Goh NS, Zappala CJ, Visca D, Maher TM, Denton CP, Ong VH, Abraham DJ, Kelleher P, Hector L, Wells AU, Renzoni EA. Serum interleukin 6 is predictive of early functional decline and mortality in interstitial lung disease associated with systemic sclerosis. *JRheumatol* 2013; 40(4): 435.
23. Kania G, Rudnik M, Distler O. Involvement of the myeloid cell compartment in fibrogenesis and systemic sclerosis. *NatRevRheumatol* 2019; 15(5): 288.
24. Terrier B, Tamby MC, Camoin L, Guilpain P, Bérezné A, Tamas N, Broussard C, Hotellier F, Humbert M, Simonneau G, Guillevin L, Mouthon L. Antifibroblast antibodies from systemic sclerosis patients bind to α -enolase and are associated with interstitial lung disease. *AnnRheumDis* 2010; 69(2): 428.
25. Fineschi S, Cozzi F, Burger D, Dayer JM, Meroni PL, Chizzolini C. Anti-fibroblast antibodies detected by cell-based ELISA in systemic sclerosis enhance the

- collagenolytic activity and matrix metalloproteinase-1 production in dermal fibroblasts. *Rheumatology* 2007; 46(12): 1779-1785.
26. Trombetta AC, Soldano S, Contini P, Tomatis V, Ruaro B, Paolino S, Brizzolara R, Montagna P, Sulli A, Pizzorni C, Smith V, Cutolo M. A circulating cell population showing both M1 and M2 monocyte/macrophage surface markers characterizes systemic sclerosis patients with lung involvement. *RespirRes* 2018; 19(1): 186.
 27. Steele R, Hudson M, Lo E, Baron M, Canadian Scleroderma Research G. Clinical decision rule to predict the presence of interstitial lung disease in systemic sclerosis. *Arthritis CareRes(Hoboken)* 2012; 64(4): 519-524.
 28. Suliman YA, Dobrota R, Huscher D, Nguyen-Kim TD, Maurer B, Jordan S, Speich R, Frauenfelder T, Distler O. Brief Report: Pulmonary function tests: high rate of false-negative results in the early detection and screening of scleroderma-related interstitial lung disease. *Arthritis Rheumatol* 2015; 67(12): 3256.
 29. D'Angelo WA, Fries JF, Masi AT, Shulman LE. Pathologic observations in systemic sclerosis (scleroderma): A study of fifty-eight autopsy cases and fifty-eight matched controls. *American Journal of Medicine* 1969; 46(3): 428-440.
 30. Walker UA, Tyndall A, Czirjak L, Denton C, Farge-Bancel D, Kowal-Bielecka O, Muller-Ladner U, Bocelli-Tyndall C, Matucci-Cerinic M. Clinical risk assessment of organ manifestations in systemic sclerosis: a report from the EULAR Scleroderma Trials And Research group database. *AnnRheumDis* 2007; 66(6): 754-763.
 31. Peoples C, Medsger TA, Jr., Lucas M, Rosario BL, Feghali-Bostwick CA. Gender differences in systemic sclerosis: relationship to clinical features, serologic status and outcomes. *Journal of scleroderma and related disorders* 2016; 1(2): 177-240.
 32. Hussein H, Lee P, Chau C, Johnson SR. The effect of male sex on survival in systemic sclerosis. *JRheumatol* 2014; 41(11): 2193.
 33. Al-Sheikh H, Ahmad Z, Johnson SR. Ethnic Variations in Systemic Sclerosis Disease Manifestations, Internal Organ Involvement, and Mortality. *JRheumatol* 2019.
 34. Nihtyanova SI, Schreiber BE, Ong VH, Rosenberg D, Moinzadeh P, Coghlan JG, Wells AU, Denton CP. Prediction of pulmonary complications and long-term survival in systemic sclerosis. *Arthritis Rheumatol* 2014; 66(6): 1625.
 35. Christmann RB, Wells AU, Capelozzi VL, Silver RM. Gastroesophageal reflux incites interstitial lung disease in systemic sclerosis: clinical, radiologic, histopathologic, and treatment evidence. *SeminArthritis Rheum* 2010; 40(3): 241-249.
 36. Herzog EL, Mathur A, Tager AM, Feghali-Bostwick C, Schneider F, Varga J. Review: interstitial lung disease associated with systemic sclerosis and idiopathic pulmonary fibrosis: how similar and distinct? *Arthritis Rheumatol* 2014; 66(8): 1967.
 37. Desai SR, Veeraraghavan S, Hansell DM, Nikolakopoulou A, Goh NSL, Nicholson AG, Colby TV, Denton CP, Black CM, Du Bois RM. CT features of lung disease in patients with systemic sclerosis: comparison with idiopathic pulmonary fibrosis and nonspecific interstitial pneumonia. *Radiology* 2004; 232(2): 560-567.
 38. Frauenfelder T, Winklehner A, Nguyen TD, Dobrota R, Baummueller S, Maurer B, Distler O. Screening for interstitial lung disease in systemic sclerosis: performance of high-resolution CT with limited number of slices: a prospective study. *AnnRheumDis* 2014; 73(12): 2069-2073.
 39. Nguyen-Kim TDL, Maurer B, Suliman YA, Morsbach F, Distler O, Frauenfelder T. The impact of slice-reduced computed tomography on histogram-

- based densitometry assessment of lung fibrosis in patients with systemic sclerosis. *JThoracDis* 2018: 10(4): 2142-2152.
40. Molberg O, Hoffmann-Vold AM. Interstitial lung disease in systemic sclerosis: progress in screening and early diagnosis. *CurrOpinRheumatol* 2016: 28(6): 613-618.
41. Goh NS, Desai SR, Veeraraghavan S, Hansell DM, Copley SJ, Maher TM, Corte TJ, Sander CR, Ratoff J, Devaraj A, Bozovic G, Denton CP, Black CM, du Bois RM, Wells AU. Interstitial lung disease in systemic sclerosis: a simple staging system. *American Journal of Respiratory and Critical Care Medicine* 2008: 177(11): 1248-1254.
42. Distler O, Allanore Y, Denton CP, Matucci-Cerinic M, Pope JE, Hinzmann B, Davies S, de Oliveira Pena J, Khanna D. Factors influencing early referral, early diagnosis and management in patients with diffuse cutaneous systemic sclerosis. *Rheumatology (Oxford)* 2018: 57(5): 813.
43. Hoffmann-Vold AM, Aalokken TM, Lund MB, Garen T, Midtvedt O, Brunborg C, Gran JT, Molberg O. Predictive value of serial high-resolution computed tomography analyses and concurrent lung function tests in systemic sclerosis. *Arthritis Rheumatol* 2015: 67(8): 2205.
44. Wells AU, Kokosi MA. Subclinical interstitial lung abnormalities: toward the early detection of idiopathic pulmonary fibrosis? *American Journal of Respiratory and Critical Care Medicine* 2016: 194(12): 1445-1446.
45. Doyle TJ, Hunninghake GM, Rosas IO. Subclinical interstitial lung disease: why you should care. *American journal of respiratory and critical care medicine* 2012: 185(11): 1147-1153.
46. Doyle TJ, Dellaripa PF, Batra K, Frits ML, Iannaccone CK, Hatabu H, Nishino M, Weinblatt ME, Ascherman DP, Washko GR, Hunninghake GM, Choi AMK, Shadick NA, Rosas IO. Functional impact of a spectrum of interstitial lung abnormalities in rheumatoid arthritis. *Chest* 2014: 146(1): 41-50.
47. Launay D, Remy-Jardin M, Michon-Pasturel U, Mastora I, Hachulla E, Lambert M, Delannoy V, Queyrel V, Duhamel A, Matran R, De Groote P, Hatron PY. High resolution computed tomography in fibrosing alveolitis associated with systemic sclerosis. *JRheumatol* 2006: 33(9): 1789.
48. Tardella M, Di Carlo M, Carotti M, Filippucci E, Grassi W, Salaffi F. Ultrasound B-lines in the evaluation of interstitial lung disease in patients with systemic sclerosis: Cut-off point definition for the presence of significant pulmonary fibrosis. *Medicine* 2018: 97(18): e0566-e0566.
49. Gigante A, Rossi Fanelli F, Lucci S, Barilaro G, Quarta S, Barbano B, Giovannetti A, Amoroso A, Rosato E. Lung ultrasound in systemic sclerosis: correlation with high-resolution computed tomography, pulmonary function tests and clinical variables of disease. *InternEmergency Med* 2016: 11(2): 213-217.
50. Song G, Bae SC, Lee YH. Diagnostic accuracy of lung ultrasound for interstitial lung disease in patients with connective tissue diseases: a meta-analysis. *ClinExpRheumatol* 2016: 34(1): 11-16.
51. López Martínez R, Hassan R, Lubertino L, Cosentino V, Barth M, Saenz C, Binda M, Montoya S, Kerzberg E. Lung ultrasound for detecting interstitial lung disease in patients with systemic sclerosis. *ClinExpRheumatol* 2014: 32(2): S29.
52. Schniering J, Guo L, Brunner M, Schibli R, Ye S, Distler O, Behe M, Maurer B. Evaluation of (99m)Tc-rhAnnexin V-128 SPECT/CT as a diagnostic tool for early stages of interstitial lung disease associated with systemic sclerosis. *Arthritis ResTher* 2018: 20(1): 183.

53. Schniering J, Benesova M, Brunner M, Haller S, Cohrs S, Frauenfelder T, Vrugt B, Feghali-Bostwick CA, Schibli R, Distler O, Mueller C, Maurer B. Visualisation of interstitial lung disease by molecular imaging of integrin alphavbeta3 and somatostatin receptor 2. *AnnRheumDis* 2019; 78(2): 218-227.
54. Cutolo M, Herrick AL, Distler O, Becker MO, Beltran E, Carpentier P, Ferri C, Inanc M, Vlachoyiannopoulos P, Chadha-Boreham H, Cottreel E, Pfister T, Rosenberg D, Torres JV, Smith V, Investigators CAPS. Nailfold Videocapillaroscopic Features and Other Clinical Risk Factors for Digital Ulcers in Systemic Sclerosis: A Multicenter, Prospective Cohort Study. *Arthritis Rheumatol* 2016; 68(10): 2527.
55. Smith V, Thevissen K, Trombetta AC, Pizzorni C, Ruaro B, Piette Y, Paolino S, De Keyser F, Sulli A, Melsens K, Cutolo M, Diseases ESGoMiR. Nailfold Capillaroscopy and Clinical Applications in Systemic Sclerosis. *Microcirculation (New York, NY: 1994)* 2016; 23(5): 364.
56. Markusse IM, Meijs J, de Boer B, Bakker JA, Schippers HPC, Schouffoer AA, Ajmone Marsan N, Kroft LJM, Ninaber MK, Huizinga TWJ, de Vries-Bouwstra JK. Predicting cardiopulmonary involvement in patients with systemic sclerosis: complementary value of nailfold videocapillaroscopy patterns and disease-specific autoantibodies. *Rheumatology (Oxford)* 2017; 56(7): 1081.
57. Mouthon L, Berezne A, Guillevin L, Valeyre D. Therapeutic options for systemic sclerosis related interstitial lung diseases. *Respir Med* 2010; 104 Suppl 1: S59.
58. Benan M, Hande I, Gul O. The natural course of progressive systemic sclerosis patients with interstitial lung involvement. *ClinRheumatol* 2007; 26(3): 349-354.
59. Lee CG, Herzog EL, Ahangari F, Zhou Y, Gulati M, Lee CM, Peng X, Feghali-Bostwick C, Jimenez SA, Varga J, Elias JA. Chitinase 1 is a biomarker for and therapeutic target in scleroderma-associated interstitial lung disease that augments TGF-beta1 signaling. *JImmunol* 2012; 189(5): 2635.
60. Silver RM, Miller KS, Kinsella MB, Smith EA, Schabel SI. Evaluation and management of scleroderma lung disease using bronchoalveolar lavage. *AmJMed* 1990; 88(5): 470.
61. White B, Moore WC, Wigley FM, Xiao HQ, Wise RA. Cyclophosphamide is associated with pulmonary function and survival benefit in patients with scleroderma and alveolitis. *AnnInternMed* 2000; 132(12): 947-954.
62. Goh NS, Veeraraghavan S, Desai SR, Cramer D, Hansell DM, Denton CP, Black CM, du Bois RM, Wells AU. Bronchoalveolar lavage cellular profiles in patients with systemic sclerosis-associated interstitial lung disease are not predictive of disease progression. *Arthritis Rheum* 2007; 56(6): 2005-2012.
63. Khanna D, Tseng CH, Farmani N, Steen V, Furst DE, Clements PJ, Roth MD, Goldin J, Elashoff R, Seibold JR, Saggarr R, Tashkin DP. Clinical course of lung physiology in patients with scleroderma and interstitial lung disease: analysis of the Scleroderma Lung Study Placebo Group. *Arthritis Rheum* 2011; 63(10): 3078-3085.
64. Moore OA, Goh N, Corte T, Rouse H, Hennessy O, Thakkar V, Byron J, Sahhar J, Roddy J, Gabbay E, Youssef P, Nash P, Zochling J, Proudman SM, Stevens W, Nikpour M. Extent of disease on high-resolution computed tomography lung is a predictor of decline and mortality in systemic sclerosis-related interstitial lung disease. *Rheumatology (Oxford)* 2013; 52(1): 155.
65. Wu W, Jordan S, Becker MO, Dobrota R, Maurer B, Fretheim H, Ye S, Siegert E, Allanore Y, Hoffmann-Vold AM, Distler O. Prediction of progression of interstitial

- lung disease in patients with systemic sclerosis: the SPAR model. *AnnRheumDis* 2018; 77(9): 1326-1332.
66. Ahmed SS, Johnson SR, Meaney C, Chau C, Marras TK. Lung function and survival in systemic sclerosis interstitial lung disease. *JRheumatol* 2014; 41(11): 2326.
67. Giacomelli R, Afeltra A, Alunno A, Bartoloni-Bocci E, Berardicurti O, Bombardieri M, Bortoluzzi A, Caporali R, Caso F, Cervera R, Chimenti MS, Cipriani P, Coloma E, Conti F, D'Angelo S, De Vita S, Di Bartolomeo S, Distler O, Doria A, Feist E, Fisher BA, Gerosa M, Gilio M, Guggino G, Liakouli V, Margiotta DPE, Meroni P, Moroncini G, Perosa F, Prete M, Priori R, Rebuffi C, Ruscitti P, Scarpa R, Shoefeld Y, Todoerti M, Ursini F, Valesini G, Vettori S, Vitali C, Tzioufas AG. Guidelines for biomarkers in autoimmune rheumatic diseases - evidence based analysis. *AutoimmunRev* 2019; 18(1): 93-106.
68. Assassi S, Sharif R, Lasky RE, McNearney TA, Estrada-Y-Martin RM, Draeger H, Nair DK, Fritzler MJ, Reveille JD, Arnett FC, Mayes MD, Group GS. Predictors of interstitial lung disease in early systemic sclerosis: a prospective longitudinal study of the GENISOS cohort. *Arthritis ResTher* 2010; 12(5): R166.
69. Weigold F, Gunther J, Pfeiffenberger M, Cabral-Marques O, Siegert E, Dragun D, Philippe A, Regensburger AK, Recke A, Yu X, Petersen F, Catar R, Biesen R, Hiepe F, Burmester GR, Heidecke H, Riemekasten G. Antibodies against chemokine receptors CXCR3 and CXCR4 predict progressive deterioration of lung function in patients with systemic sclerosis. *Arthritis ResTher* 2018; 20(1): 52.
70. Tiev KP, Hua-Huy T, Kettaneh A, Gain M, Duong-Quy S, Toledano C, Cabane J, Dinh-Xuan AT. Serum CC chemokine ligand-18 predicts lung disease worsening in systemic sclerosis. *EurRespirJ* 2011; 38(6): 1355.
71. Elhai M, Hoffmann-Vold AM, Avouac J, Pezet S, Cauvet A, Leblond A, Fretheim H, Garen T, Kuwana M, Molberg Ø. Performance of candidate serum biomarkers for systemic sclerosis-interstitial lung disease. *Arthritis & Rheumatology* 2019.
72. Schupp J, Becker M, Gunther J, Muller-Quernheim J, Riemekasten G, Prasse A. Serum CCL18 is predictive for lung disease progression and mortality in systemic sclerosis. *EurRespirJ* 2014; 43(5): 1530.
73. Hoffmann-Vold A-M, Tennøe AH, Garen T, Midtvedt Ø, Abraitte A, Aaløkken TM, Lund MB, Brunborg C, Aukrust P, Ueland T, Molberg Ø. High level of chemokine CCL18 is associated with pulmonary function deterioration, lung fibrosis progression, and reduced survival in systemic sclerosis. *CHEST* 2016; 150(2): 299-306.
74. Liu X, Mayes MD, Pedroza C, Draeger HT, Gonzalez EB, Harper BE, Reveille JD, Assassi S. Does C-reactive protein predict the long-term progression of interstitial lung disease and survival in patients with early systemic sclerosis? *Arthritis CareRes(Hoboken)* 2013; 65(8): 1375-1380.
75. Ross L, Stevens W, Rabusa C, Wilson M, Ferdowsi N, Walker J, Sahhar J, Ngian GS, Zochling J, Roddy J, Tymms K, Major G, Strickland G, Proudman SM, Nikpour M. The role of inflammatory markers in assessment of disease activity in systemic sclerosis. *ClinExpRheumatol* 2018; 36 Suppl 113(4): 126-134.
76. Yamakawa H, Hagiwara E, Kitamura H, Yamanaka Y, Ikeda S, Sekine A, Baba T, Okudela K, Iwasawa T, Takemura T, Kuwano K, Ogura T. Serum KL-6 and surfactant protein-D as monitoring and predictive markers of interstitial lung disease in patients with systemic sclerosis and mixed connective tissue disease. *JThoracDis* 2017; 9(2): 362-371.

77. Kuwana M, Shirai Y, Takeuchi T. Elevated serum Krebs von den Lungen-6 in early disease predicts subsequent deterioration of pulmonary function in patients with systemic sclerosis and interstitial lung disease. *JRheumatol* 2016; 43(10): 1825.
78. Salazar GA, Kuwana M, Wu M, Estrada-Y-Martin RM, Ying J, Charles J, Mayes MD, Assassi S. KL-6 but not CCL-18 is a predictor of early progression in systemic sclerosis-related interstitial lung disease. *JRheumatol* 2018; 45(8): 1153.
79. Volkmann ER, Tashkin DP, Kuwana M, Li N, Roth MD, Charles J, Hant FN, Bogatkevich GS, Akter T, Kim G, Goldin J, Khanna D, Clements PJ, Furst DE, Elashoff RM, Silver RM, Assassi S. Progression of interstitial lung disease in systemic sclerosis: the importance of pneumoproteins Krebs von den Lungen 6 and CCL18. *Arthritis & Rheumatology* 2019; 71(12): 2059-2067.
80. Goh NS, Hoyles RK, Denton CP, Hansell DM, Renzoni EA, Maher TM, Nicholson AG, Wells AU. Short-term pulmonary function trends are predictive of mortality in interstitial lung disease associated with systemic sclerosis. *Arthritis Rheumatol* 2017; 69(8): 1670.
81. Berry CE, Drummond MB, Han MK, Li D, Fuller C, Limper AH, Martinez FJ, Schwarz MI, Sciurba FC, Wise RA. Relationship between lung function impairment and health-related quality of life in COPD and interstitial lung disease. *Chest* 2012; 142(3): 704-711.
82. Wells AU. High-resolution computed tomography and scleroderma lung disease. *Rheumatology (Oxford)* 2008; 47 Suppl 5: v59.
83. Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, Enright PL, Hankinson JL, Ip MS, Zheng J, Stocks J, E. R. S. Global Lung Function Initiative. Multi-ethnic reference values for spirometry for the 3-95-yr age range: the global lung function 2012 equations. *European Respiratory Journal* 2012; 40(6): 1324-1343.
84. Langhammer A, Johannessen A, Holmen TL, Melbye H, Stanojevic S, Lund MB, Melsom MN, Bakke P, Quanjer PH. Global Lung Function Initiative 2012 reference equations for spirometry in the Norwegian population. *European Respiratory Journal* 2016; 48(6): 1602-1611.
85. Distler O, Highland KB, Gahlemann M, Azuma A, Fischer A, Mayes MD, Raghu G, Sauter W, Girard M, Alves M, Clerisme-Beaty E, Stowasser S, Tetzlaff K, Kuwana M, Maher TM. Nintedanib for Systemic Sclerosis–Associated Interstitial Lung Disease. *N Engl J Med* 2019.
86. Boehringer Ingelheim GmbH. OFEV® (nintedanib) capsules, for oral use. 2016 [cited 2019 November 11]; Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/205832s001lbl.pdf
87. Kowal-Bielecka O, Franssen J, Avouac J, Becker M, Kulak A, Allanore Y, Distler O, Clements P, Cutolo M, Czirjak L, Damjanov N, Del Galdo F, Denton CP, Distler JHW, Foeldvari I, Figelstone K, Frerix M, Furst DE, Guiducci S, Hunzelmann N, Khanna D, Matucci-Cerinic M, Herrick AL, van den Hoogen F, van Laar JM, Riemekasten G, Silver R, Smith V, Sulli A, Tarner I, Tyndall A, Welling J, Wigley F, Valentini G, Walker UA, Zulian F, Muller-Ladner U, Coauthors E. Update of EULAR recommendations for the treatment of systemic sclerosis. *AnnRheumDis* 2017; 76(8): 1327–1339.
88. Fernandez-Codina A, Walker KM, Pope JE, Scleroderma Algorithm G. Treatment algorithms for systemic sclerosis according to experts. *Arthritis Rheumatol* 2018; 70(11): 1820.
89. Johnson SR, Devakandan K. Guidelines and recommendations towards evidence-based management of systemic sclerosis. 2019; 5: 115.

90. Cottin V, Brown KK. Interstitial lung disease associated with systemic sclerosis (SSc-ILD). *RespirRes* 2019; 20(1): 13.
91. Zeisberg M, Kalluri R. Cellular mechanisms of tissue fibrosis. 1. Common and organ-specific mechanisms associated with tissue fibrosis. *Am J Physiol Cell Physiol* 2013; 304(3): C216-C225.
92. Distler JH, Feghali-Bostwick C, Soare A, Asano Y, Distler O, Abraham DJ. Review: frontiers of antifibrotic therapy in systemic sclerosis. *Arthritis Rheumatol* 2017; 69(2): 257.
93. Ho YY, Lagares D, Tager AM, Kapoor M. Fibrosis – a lethal component of systemic sclerosis. *NatRevRheumatol* 2014; 10(7): 390.
94. Khanna D, Distler JHW, Sandner P, Distler O. Emerging strategies for treatment of systemic sclerosis. *Journal of Scleroderma and Related Disorders* 2016; 1(2): 186-193.
95. Johnson SR, Hinchcliff M, Asano Y. Controversies: molecular vs. clinical systemic sclerosis classification. *Journal of Scleroderma and Related Disorders* 2016; 1(3): 277-285.
96. Wu M, Baron M, Pedroza C, Salazar GA, Ying J, Charles J, Agarwal SK, Hudson M, Pope J, Zhou X, Reveille JD, Fritzler MJ, Mayes MD, Assassi S. CCL2 in the circulation predicts long-term progression of interstitial lung disease in patients with early systemic sclerosis: data from two independent cohorts. *Arthritis Rheumatol* 2017; 69(9): 1871.
97. van Bon L, Affandi AJ, Broen J, Christmann RB, Marijnissen RJ, Stawski L, Farina GA, Stifano G, Mathes AL, Cossu M, York M, Collins C, Wenink M, Huijbens R, Hesselstrand R, Saxne T, DiMarzio M, Wuttge D, Agarwal SK, Reveille JD, Assassi S, Mayes M, Deng Y, Drenth JP, de Graaf J, den Heijer M, Kallenberg CG, Bijl M, Loof A, van den Berg WB, Joosten LA, Smith V, de Keyser F, Scorza R, Lunardi C, van Riel PL, Vonk M, van Heerde W, Meller S, Homey B, Beretta L, Roest M, Trojanowska M, Lafyatis R, Radstake TR. Proteome-wide analysis and CXCL4 as a biomarker in systemic sclerosis. *NEnglJMed* 2014; 370(5): 433.
98. Volkmann ER, Tashkin DP, Roth MD, Clements PJ, Khanna D, Furst DE, Mayes M, Charles J, Tseng C-H, Elashoff RM, Assassi S. Changes in plasma CXCL4 levels are associated with improvements in lung function in patients receiving immunosuppressive therapy for systemic sclerosis-related interstitial lung disease. *Arthritis ResTher* 2016; 18(1): 305.
99. Steen VD, Conte C, Owens GR, Medsger TA, Jr. Severe restrictive lung disease in systemic sclerosis. *Arthritis Rheum* 1994; 37(9): 1283-1289.
100. Morgan C, Knight C, Lunt M, Black CM, Silman AJ. Predictors of end stage lung disease in a cohort of patients with scleroderma. *AnnRheumDis* 2003; 62(2): 146-150.
101. Plastiras SC, Karadimitrakis SP, Ziakas PD, Vlachoyiannopoulos PG, Moutsopoulos HM, Tzelepis GE. Scleroderma lung: initial forced vital capacity as predictor of pulmonary function decline. *Arthritis Rheum* 2006; 55(4): 598-602.
102. Ariani A, Silva M, Seletti V, Bravi E, Saracco M, Parisi S, De Gennaro F, Idolazzi L, Caramaschi P, Benini C, Bodini FC, Scire CA, Carrara G, Lumetti F, Alfieri V, Bonati E, Lucchini G, Aiello M, Santilli D, Mozzani F, Imberti D, Michieletti E, Arrigoni E, Delsante G, Pellerito R, Fusaro E, Chetta A, Sverzellati N. Quantitative chest computed tomography is associated with two prediction models of mortality in interstitial lung disease related to systemic sclerosis. *Rheumatology (Oxford)* 2017; 56(6): 922.

Tables

Table 1. Clinical and biological factors associated with the presence of ILD in SSc

Characteristics predictive of ILD in SSc			
Demographic	Disease status	Biomarkers	Pulmonary function tests
Male Afro-Caribbean ethnicity Peoples et al. 2016 [31]; Hussein et al. 2014 [32]; Al-Sheikh et al. 2019 [33]	Diffuse cutaneous sclerosis Walker et al. 2007 [30]	Anti-topoisomerase I positivity Walker et al. 2007 [30]	Lower FVC Lower DL_{CO} Nihtyanova et al. 2014 [34]; Peoples et al. 2016 [31]

Abbreviations: DL_{CO}, diffusing capacity of the lungs for carbon monoxide; FVC, forced vital capacity.

Table 2. Clinical and biological factors associated with progression of SSc-ILD^a

Characteristics predictive of disease progression				
Demographic	Disease status	Biomarkers	Pulmonary function tests	Imaging
<p>African American ethnicity</p> <p>Advanced age</p> <p>Assassi et al., 2010 [68]; Al Sheikh et al. 2019 [33]; Ahmed et al., 2014 [66]</p>	<p>Diffuse cutaneous sclerosis</p> <p>Nihtyanova et al., 2014 [34]</p>	<p>IL-6</p> <p>CRP</p> <p>CCL2</p> <p>CCL18</p> <p>CXCL4</p> <p>KL-6</p> <p>SP-D</p> <p>Anti-topoisomerase I</p>	<p>Low baseline FVC</p> <p>Low baseline DL_{CO}</p> <p>Steen et al., 1994 [99]; Morgan et al., 2003 [100]; Plastiras et al., 2006 [101]; Nihtyanova et al., 2014 [34]; Ahmed et al., 2014 [66]; Morgan et al., 2003 [100]; Nihtyanova et al., 2014</p>	<p>Extent of ILD on HRCT</p> <p>Goh et al., 2008 [41]; Khanna et al., 2011 [63]; Moore et al., 2013 [64]; Ariani et al., 2017 [79, 98, 102]</p>

		Anti-CXCR3 Anti-CXCR4 De Lauretis et al., 2013 [22]; Liu et al., 2013 [74]; Ross et al., 2018 [75]; Wu et al., 2017 [96]; Tiev et al., 2011 [70]; Schupp et al., 2014 [72]; Elhai et al., 2019 [71]; van Bon et al., 2014 [97]; Kuwana et al. 2016 [77]; Salazar et al., 2018 [78]; Yamakawa et al., 2017 [76]; Assassi et al., 2010 [68];	[34]	
--	--	--	------	--

		Nihtyanova et al., 2014 [34]; Weigold et al., 2018 [69]; Volkmann et al. 2019 [79]; Volkmann et al. 2016 [98]		
--	--	---	--	--

^aWorsening of pulmonary function.

Abbreviations: CCL, chemokine ligand; CI, confidence interval; CRP, C-reactive protein; DL_{CO}, diffusing capacity of the lung for carbon monoxide; FVC, forced vital capacity; HR, hazard ratio; HRCT, high-resolution computed tomography; IL, interleukin; ILD, interstitial lung disease; KL-6, Krebs von den Lungen-6; NR, not reported; OR, odds ratio; SP-D, surfactant protein D.

Figure legends

Figure 1. Schematic of the key pathways implicated in systemic SSc-ILD [16, 18, 36]

Abbreviations: TGF β , transforming growth factor beta.

Figure 2. (A) HRCT from a patient with SSc-ILD (disease extent >20%). (B) Lung histology from a patient with SSc showing NSIP

Abbreviations: HRCT, high-resolution computed tomography; ILD, interstitial lung disease; NSIP, nonspecific interstitial pneumonia; SSc, systemic sclerosis.

Figure 3. Proposed definition of disease progression

Abbreviations: DLco, diffusing capacity of the lungs for carbon monoxide; FVC, forced vital capacity.





