



## Early View

Research letter

### **Five-year survival after an acute episode of decompensated pulmonary arterial hypertension in the modern management era of right heart failure**

Laurent Savale, Constance Vuillard, Jérémie Pichon, Athénaïs Boucly, Anne Roche, Mitja Jevnikar, Nathan Ebstein, Xavier Jaïs, Jérôme Le Pavec, David Montani, Olaf Mercier, Olivier Sitbon, Elie Fadel, Marc Humbert

Please cite this article as: Savale L, Vuillard C, Pichon Jérémie, *et al.* Five-year survival after an acute episode of decompensated pulmonary arterial hypertension in the modern management era of right heart failure. *Eur Respir J* 2021; in press (<https://doi.org/10.1183/13993003.00466-2021>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

**Five-year survival after an acute episode of decompensated pulmonary arterial hypertension in the modern management era of right heart failure**

Laurent Savale<sup>1,2,3</sup>, Constance Vuillard<sup>1,2,3</sup>, Jérémie Pichon<sup>1,2,3</sup>, Athénaïs Boucly<sup>1,2,3</sup>, Anne Roche<sup>1,2,3</sup>, Mitja Jevnikar<sup>1,2,3</sup>, Nathan Ebstein<sup>1,2,3</sup>, Xavier Jaïs<sup>1,2,3</sup>, Jérôme Le Pavec<sup>1,2,4</sup>, David Montani<sup>1,2,3</sup>, Olaf Mercier<sup>1,2,4</sup>, Olivier Sitbon<sup>1,2,3</sup>, Elie Fadel<sup>1,2,4</sup>, Marc Humbert<sup>1,2,3</sup>

1. Université Paris-Saclay, Faculty of Medicine, Le Kremlin-Bicêtre, France

2. INSERM UMR\_S 999, Le Kremlin-Bicêtre, France

3. AP-HP, Service de Pneumologie et soins intensifs respiratoires, Hôpital Bicêtre, Le Kremlin-Bicêtre, France

4. Service de Chirurgie Thoracique, Vasculaire et Transplantation Cardio-pulmonaire, Hôpital Marie-Lannelongue, Le Plessis-Robinson, France

**Correspondance :**

Laurent Savale, MD , PhD

Service de pneumologie, Le Kremlin-Bicêtre, France

78 rue du général Leclerc, 94270 Le Kremlin Bicêtre, France.

Tél: +33 145 21 79 74

Fax: +33 145 21 79 71

Email: [laurent.savale@aphp.fr](mailto:laurent.savale@aphp.fr)

Acute decompensated pulmonary arterial hypertension (PAH) is characterized by rapid worsening of clinical signs of right heart failure (RHF) with subsequent congestion and systemic circulatory insufficiency that can lead to multisystem organ failure [1–3]. Short-term outcomes of acute decompensated RHF is very poor and remains the first cause of mortality in PAH [4, 5]. Intensive care of acute decompensated PAH is based on treatment of triggering factors, careful fluid management, and strategies to improve cardiac function and reduce right ventricular afterload [1]. However, this medical strategy is not always sufficient to restore a long-lasting balance between the afterload imposed on the right ventricle and its capacity for compensation. In case of refractory RHF despite maximal medical treatment, the use of mechanical support should now be considered in selected candidates for lung transplantation, or less commonly as a bridge to recovery in patients with a treatable cause of right-sided heart failure [1]. Veno-arterial extracorporeal membrane oxygenation (ECMO) is currently the most widely used strategy to support the right ventricle in PAH patients. This strategy, combined with changes in organ allocation rules to prioritize patients with a short-term life-threatening condition should contribute to improve survival of eligible patients with end-stage PAH [6]. However, long-term survival of patients admitted in ICU for severe acute RHF management has not been studied extensively in the modern management era of mechanical support and high-priority lung transplantation.

In the present report, we describe the management and 5-year outcomes of consecutive PAH patients with acute RHF referred to the intensive care unit (ICU) of the French reference center of pulmonary hypertension between February 2015 and February 2016. All patients were enrolled in the French PAH Registry and followed-up during a 5-year period. This retrospective study complied with the Declaration of Helsinki. Although French law does not require informed consent for retrospective data collection, the data collected were anonymized and complied with the requirements of the ‘*Commission Nationale Informatique*

*et Libertés*’, the organization dedicated to privacy, information technology and civil rights in France. The committee approved the methods used to collect and analyse the data on 24 May 2003 (approval number 842063).

Patients were divided into three age grades corresponding to tertiles *i.e* < 50 year-old group, 50-65 year-old group and >65 year-old group. Continuous variables were expressed as mean  $\pm$  standard deviation (SD). Categorical variables were expressed as number of patients and relative frequencies (percent). Overall survival time was defined from the date of ICU admission to death. Veno-arterial ECMO- and transplant-free survival was defined from the date of ICU admission to death, ECMO, or transplantation. Surviving patients were censored at the date of last clinical contact. The Kaplan-Meier method was used to estimate the proportion of patients surviving at each time point up to 5 years post-ICU admission. Survival curves according to the three age grades were compared with log-rank test. Univariable analysis based on the Cox proportional hazards model was used to examine the relationship between survival and selected variables at baseline and day 3.

Seventy patients (60% females, n=42) were enrolled in the study, with a mean age of  $57\pm 16$  years old at the time of ICU admission (n=24, 34%, in the less than 50 year-old age group, n=22, 32%, in the 50-65 year-old group, and n=24, 34%, in the more than 65 year-old age group). According to the last classification [7], 21 patients had idiopathic PAH (28%), 2 had heritable PAH (3%), 5 had PAH induced by drug (7%), 29 had PAH associated with another condition (41%) (connective tissue disease, n=15, HIV infection, n=2, portal hypertension, n=6, congenital heart disease, n=6) and 12 had PAH with overt features of venous involvement (pulmonary veno-occlusive disease) (17%). PAH diagnosis was revealed by the episode of acute decompensation in 20 patients (28%). At time of admission, 15 patients were treated with a monotherapy (21%), 21 with a dual combination (30%) and 14

with a triple therapy (20%). A parenteral prostacyclin analog was included in the treatment regimen at admission in 16 patients (23%). A triggering factor was identified in 19 cases (27%) (infection, n=9; arrhythmia, n=4; postoperative, n=2; hemorrhage, n=3; treatment disruption, n=1). The value of the simplified acute physiology score at admission (SAPS II) was  $22\pm 9$ .

All were treated with intravenous diuretics and the use of inotropic support (dobutamine in all cases) and vasopressors (norepinephrine in all cases) was required in 46 (66%) and 23 (33%) patients, respectively. At time of admission, 5 patients were already listed for lung (LT) or heart-lung transplantation (HLT). In other patients less than 65 years old, eligibility for LT or HLT was systematically assessed, considering contraindications to transplantation defined in international recommendations [8]. Proportion of patients who were potentially eligible for LT or HLT was 71% and 27% in the less than 50 year-old age group and the 50-65 year-old group, respectively. Finally, 11 patients (16%) were transplanted with high priority organ allocation (national priority access to lung transplantation) including 3 patients aged 50-65 years. A venous-arterial ECMO was implanted in four awake patients under local anesthesia as bridge to transplantation. One of them died in ICU before transplantation and three were transplanted 1 to 9 days after circulatory assistance initiation (**Figure 1A**). After adjusting for age, creatinine at admission (HR 0.993, p=0.03), brain natriuretic peptide (BNP) plasma levels at day 3 (HR 0.999, p=0.03) and mean arterial pressure (HR 1.044, p=0.009) at day 3 were associated with higher risk of ECMO implantation, urgent lung transplantation or death in ICU. Death in ICU was higher in patients with a triggering factor identified at time of admission (37% vs 14%, p=0.03).

Patients were not lost of follow-up. The rate of death at 3 months, 1 year and 5 years after ICU admission were 31%, 41% and 66% respectively (**Figure 1B**). The overall survival at 5 years was significantly higher in the less than 50 year-old patient's group in comparison with

the 50-65 year-old and more than 65-year-old patient's groups ( $p=0.0004$ ) (**Figure 1C**): survival at 5 years in less than 50 year-old patients was 70%, while it was 23% and 12% in the 50-65 year-old and more than 65-year old patient's groups, respectively. By contrast, lung transplantation and ECMO-free survival was similar in the three age grades (**Figure 1D**).

Our results show that acute decompensation of PAH remains a devastating life-threatening condition in the current management era. However, our data suggest recent improvements of short and long-term outcomes in patients less than 50 year-old, at least in part thanks to novel organ allocation rules, implementation of high-priority lung transplantation and use of ECMO as bridge to transplantation in eligible patients with refractory RHF. While event-free survival without ECMO and/or lung transplantation was similar in all age groups, the overall survival of less than 50 year-old patients was better than in older patients in our contemporary cohort (70% versus 23% and 12% in the 50-65 year-old and more than 65-year old patient's groups, respectively). In a previously reported cohort of patients admitted in the same ICU for acute PAH decompensation before the modern era of RHF management, the short-term survival at 3 months was similarly poor in all age groups (50%) [4]. At the time of our previous study, the time on the waiting list for lung transplantation was exceedingly long for a life-threatening condition and the use of ECMO was not yet generalized to the management of refractory RHF. These observations underscore that patient with acute decompensated PAH should be preferably managed in expert centers that are able to access all medical, interventional and surgical options including medical therapies, ECMO and lung transplantation, as emphasized by the 6<sup>th</sup> world symposium on PH [1]. Most health systems have updated their lung allocation pathways with the implementation of scores and/or rules aiming to shorten the waiting time for the most advanced patients with an urgent need of high priority access to transplantation [6, 8, 9]. The possibility to offer high-

priority lung transplantation to PAH patients with refractory RHF has led to a major reduction of the waiting time and, by inference, reduced the risks of pre- and post-operative complications. The youngest patients have benefited more from this novel management approach. By contrast, refractory RHF in patients above the age of 50 remains a dramatic life-threatening event in the evolution of the disease. Frequent comorbidities in these patients limit the access to an urgent transplantation. These results underline that fully evaluation for lung transplantation must be systematically considered early in the course of PAH in patients who do not reach a low risk profile with medical therapy, as proposed by the last 6<sup>th</sup> world symposium [1, 10]. The long-term outcomes of our cohort highlight that transplantation of outpatients in stable patients remains the best option.

The main limitation of the present study is its monocentric nature. However, it should be highlighted that the French PAH reference centre hosts a dedicated pulmonary vascular disease ICU working in close partnership with a nationwide network of 25 centres. We have followed country-specific lung allocation rules (High Priority Allocation Program) which may not apply to other institutions [6]. However, several countries have developed and refined lung allocation scores in order to prioritise the most severe patients at the time of listing and therefore reduce the risk of death on the waiting list [9, 11].




In conclusion, our data suggest that the implementation of novel medical, instrumental and surgical management of RHF translate to improvements in long-term survival of our youngest patients with acute decompensated PAH. Last, our data argue for management of such patients in expert centres with multidisciplinary teams able to offer all modern treatment options.

**Figure 1. Outcomes after an acute episode of decompensated PAH.**

**A. Characteristics of patients who needed veno-arterial ECMO and/or urgent transplantation (n=12)**

age	sex	Type of PAH	PAH therapy at admission	On list at admission	Triggering factor	Inotropic support	Vasopressor	V-A ECMO	Time in ICU before LT/HLT	Outcomes at 5 years
46y	F	Idiopathic	Triple therapy*	No	No	Yes	Yes	Yes	23 days	Alive
55y	M	Idiopathic	Triple therapy*	No	No	Yes	Yes	No	11 days	Death
20y	M	Drug-induced	None	No	No	Yes	No	No	21 days	Alive
49y	F	SSc	Oral dual therapy	Yes	Arrhythmia	Yes	Yes	Yes	2 days	Death
43y	M	CHD	Triple therapy*	Yes	No	Yes	Yes	No	4 days	Alive
28y	F	HIV	Triple therapy*	No	No	Yes	No	No	4 days	Alive
57y	F	SSc	Oral dual therapy	Yes	No	Yes	Yes	No	15 days	Death
56y	F	Drug-induced	None	No	No	Yes	Yes	No	14 days	Alive
17y	F	Heritable	Triple therapy*	Yes	No	Yes	No	No	9 days	Alive
24y	F	Heritable PVOD	Monotherapy	Yes	No	Yes	No	No	9 days	Alive
28y	F	Idiopathic	Dual therapy*	No	No	Yes	Yes	Yes	7 days	Alive
50y	M	Idiopathic	Triple therapy*	No	No	Yes	Yes	Yes	-	Death before LT





(A) Characteristics of patients who needed veno-arterial ECMO and/or urgent transplantation.

(B) Short- and long-term outcomes of patients admitted in ICU for acute decompensated

PAH. (C) Overall survival. (D) ECMO- and transplant-free survival.

Abbreviations: CHD: congenital heart disease; HIV: human immunodeficiency virus; LT: lung transplantation; PVOD: pulmonary veno-occlusive disease; SSc: systemic scleroderma; V-A ECMO: veno-arterial extracorporeal membrane oxygenation.

\* including a parenteral prostacyclin analog

## References

1. Hoeper MM, Benza RL, Corris P, de Perrot M, Fadel E, Keogh AM, Kühn C, Savale L, Klepetko W. Intensive care, right ventricular support and lung transplantation in patients with pulmonary hypertension. *Eur. Respir. J.* 2018; .
2. Savale L, Weatherald J, Jaïs X, Vuillard C, Boucly A, Jevnikar M, Montani D, Mercier O, Simonneau G, Fadel E, Sitbon O, Humbert M. Acute decompensated pulmonary hypertension. *Eur Respir Rev* 2017; 26.
3. Harjola V-P, Mebazaa A, Čelutkienė J, Bettex D, Bueno H, Chioncel O, Crespo-Leiro MG, Falk V, Filippatos G, Gibbs S, Leite-Moreira A, Lassus J, Masip J, Mueller C, Mullens W, Naeije R, Nordegraaf AV, Parissis J, Riley JP, Ristic A, Rosano G, Rudiger A, Ruschitzka F, Seferovic P, Sztrymf B, Vieillard-Baron A, Yilmaz MB, Konstantinides S. Contemporary management of acute right ventricular failure: a statement from the Heart Failure Association and the Working Group on Pulmonary Circulation and Right Ventricular Function of the European Society of Cardiology. *Eur. J. Heart Fail.* 2016; 18: 226–241.
4. Sztrymf B, Souza R, Bertoletti L, Jaïs X, Sitbon O, Price LC, Simonneau G, Humbert M. Prognostic factors of acute heart failure in patients with pulmonary arterial hypertension. *Eur. Respir. J.* 2010; 35: 1286–1293.
5. Haddad F, Peterson T, Fuh E, Kudelko KT, de Jesus Perez V, Skhiri M, Vagelos R, Schnittger I, Denault AY, Rosenthal DN, Doyle RL, Zamanian RT. Characteristics and outcome after hospitalization for acute right heart failure in patients with pulmonary arterial hypertension. *Circ Heart Fail* 2011; 4: 692–699.
6. Savale L, Le Pavec J, Mercier O, Mussot S, Jaïs X, Fabre D, O’Connell C, Montani D, Stephan F, Sitbon O, Simonneau G, Dartevelle P, Humbert M, Fadel E. Impact of High-Priority Allocation on Lung and Heart-Lung Transplantation for Pulmonary Hypertension. *Ann. Thorac. Surg.* 2017; 104: 404–411.
7. Simonneau G, Montani D, Celermajer DS, Denton CP, Gatzoulis MA, Krowka M, Williams PG, Souza R. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur. Respir. J.* 2018; .
8. Weill D, Benden C, Corris PA, Dark JH, Davis RD, Keshavjee S, Lederer DJ, Mulligan MJ, Patterson GA, Singer LG, Snell GI, Verleden GM, Zamora MR, Glanville AR. A consensus document for the selection of lung transplant candidates: 2014--an update from the Pulmonary Transplantation Council of the International Society for Heart and Lung Transplantation. *J. Heart Lung Transplant.* 2015; 34: 1–15.
9. Gottlieb J, Smits J, Schramm R, Langer F, Buhl R, Witt C, Strueber M, Reichenspurner H. Lung Transplantation in Germany Since the Introduction of the Lung Allocation Score. *Dtsch Arztebl Int* 2017; 114: 179–185.
10. Galiè N, Channick RN, Frantz RP, Grünig E, Jing ZC, Moiseeva O, Preston IR, Pulido T, Safdar Z, Tamura Y, McLaughlin VV. Risk stratification and medical therapy of pulmonary arterial hypertension. *Eur. Respir. J.* 2018; .

11. Schaffer JM, Singh SK, Joyce DL, Reitz BA, Robbins RC, Zamanian RT, Mallidi HR. Transplantation for idiopathic pulmonary arterial hypertension: improvement in the lung allocation score era. *Circulation* 2013; 127: 2503–2513.