



Early View

Research letter

Effect of listening to breathing recordings on self-reported breathlessness: a public experiment

Natasha Lovell, Simon N Etkind, Joanna M Davies, Wendy Prentice, Irene J Higginson, Katherine E Sleeman

Please cite this article as: Lovell N, Etkind SN, Davies JM, *et al.* Effect of listening to breathing recordings on self-reported breathlessness: a public experiment. *Eur Respir J* 2023; in press (<https://doi.org/10.1183/13993003.01439-2022>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©The authors 2023. For reproduction rights and permissions contact permissions@ersnet.org

Effect of listening to breathing recordings on self-reported breathlessness: a public experiment

Author list

Natasha Lovell^{#1}, Simon N Etkind^{#1,2}, Joanna M Davies¹, Wendy Prentice³, Irene J Higginson¹, Katherine E Sleeman¹

1. Cicely Saunders Institute of Palliative Care, Policy and Rehabilitation, King's College London, Bessemer Road, London, SE5 9PJ

2. Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

3. King's College Hospital, Denmark Hill, London, SE5 9RS

Contributed equally

Corresponding author

Natasha.lovell@nbt.nhs.uk

Take home message

In this public experiment - a collaboration between the Cicely Saunders Institute and Science Gallery London – we found that listening to audio recordings of breathlessness resulted in a noticeable increase in self-reported breathlessness.

To the Editor.

Breathlessness is a common and persistent symptom for those living with chronic lung conditions and advanced disease(1). It can be distressing for those who experience it, and often results in anxiety, physical inactivity, and a poorer quality-of-life(2, 3). Additionally, it impacts significantly on those who are close including friends and family, with a considerable care burden(4).

Breathlessness can occur physiologically e.g. during exercise, but can also be experienced in many scenarios, including respiratory pathology or psychological distress. Complex interactions between neural networks in the brain are now thought to underpin the perception of breathlessness, and a growing body of research has identified consistent areas of the brain that are associated with breathlessness including the insula, cingulate and sensory cortices, the amygdala, and the periaqueductal gray matter(5-7). A recent body of research proposes that the perception of breathlessness is influenced by priors generated based on expectations learned from past experiences, which determine the importance assigned to sensory inputs(8, 9). These priors are compared against incoming sensory signal to generate the symptom experience. This could explain why simply observing breathlessness in others can elicit breathlessness and negative affect in the absence of physiological changes(10). Additionally a placebo effect has been demonstrated, where expectation of breathlessness resulted in increased neural processing and subsequent intensity of symptom experience(11).

It is not currently known (i) whether breathlessness is induced when people are exposed to audio recordings of breathlessness, or (ii) whether people listening to breathlessness precipitated by different scenarios will experience breathlessness to different degrees. We aimed to explore these questions in healthy volunteers.

Our hypotheses were:

- 1) Listening to brief audio recordings of breathlessness precipitated by different scenarios will induce breathlessness in healthy volunteers.
- 2) The degree of breathlessness that is induced will be affected by the scenario of breathlessness that is being listened to.

To collect preliminary data to test these hypotheses, we conducted a live public experiment in collaboration with Science Gallery, King's College London. We ran the experiment on three dates between 12th October and 16th November 2019 as part of the 'On Edge' exhibition at Science Gallery, King's College London. Participants were members of the public aged 16 and above who attended

the exhibition and opted to participate. Data were collected via an online interface using a touchscreen tablet. Written consent was obtained prior to taking part. Participants listened to audio recordings of breathlessness, which we defined as breathing patterns elicited by one of 4 scenarios: 1. A healthy person post-exercise, 2. An anxious person, 3. A person with chronic lung disease and 4. A person approaching the end of life. They were asked to correctly identify the breathlessness scenario, and were asked to report how breathless they felt at baseline and after each recording using the Numerical Rating Scale (NRS) 'now' which is a validated measure of how breathless a person is feeling at the current time on a scale of 0-10 (high scores represent worse breathlessness) (12). We estimated that a minimum sample size of 200 participants would be sufficient based on experience from Herzog et al's 2018 experiment(10).

Recordings were sourced from consenting patients and using an online audio bank. They were edited to create a 6 second looped clip. The order that recordings were presented to participants was randomised as follows: for each participant, an automated computer program allocated each of the four recordings a random position (between 1 & 4) in the experimental sequence.

To test hypothesis 1) we compared mean (sd) NRS scores at baseline to scores after listening to any of the breathlessness recordings. To test hypothesis 2) we compared mean (sd) NRS scores at baseline, to scores after each breathlessness scenario. We used a one-way ANOVA with repeated measures and post-hoc pair-wise comparisons, and applied a Bonferroni adjustment for multiple comparisons. Results were confirmed using a Friedman test. Due to randomisation of the recording order, we did not adjust for order of recordings during analysis. Ethical approval was obtained from King's College London (Reference HR-18/19-12676).

268 people participated in the live experiment. 63% were female (n=169) and most were aged 18-35 (67%, n=180). 87% (n=232) reported a previous experience of anxiety, and 70.5% (n=189) reported a previous experience of breathlessness. The breathlessness scenario was correctly identified as anxiety by 69% of participants, exercise by 59%, chronic lung disease by 39%, and approaching the end of life by 33%.

Hypothesis 1) Participants' self-reported breathlessness was greater after listening to any of the recordings than at baseline. NRS was mean (SD) 1.29(1.82) at baseline, 1.42(1.85) after recording 1, 1.50(1.82) after recording 2, 1.51(1.84) after recording 3, and 1.56(1.98) after recording 4.

Hypothesis 2) Self-reported breathlessness increased after listening to any of the four recordings, and the increase was largest after listening to the anxiety scenario (table 1). The increase in NRS

score from baseline was statistically significant at $p < 0.05$ after listening to the recordings of anxiety, end of life or exercise, but there was no statistically significant change after listening to the recording of lung disease (table 1).

Figure 1: NRS scores prior to listening to any recordings of breathlessness, and after listening to 4 randomly ordered recordings of breathlessness

In this public experiment, listening to audio recordings of breathlessness increased participants' self-reported breathlessness. Listening to the audio recording of breathlessness caused by anxiety increased self-reported breathlessness most.

The sound of breathlessness caused by anxiety or exercise were correctly identified more often than the breathlessness recordings precipitated by the other scenarios. It may be that these sounds are more familiar than the sound of someone approaching the end-of-life or of chronic lung disease, which participants may have been less likely to have experienced prior to taking part in the experiment. We were unable to explore the relationship between prior experience of breathlessness/anxiety and the effect of the breathlessness recordings on NRS score in this preliminary dataset, and this is an important question for future research.

These findings, whilst preliminary, raise the question of how exposure to audible breathlessness may affect informal carers and health professionals involved in the care of those living with chronic breathlessness. Findings from Herzog et al indicate that observation of breathlessness can induce vicarious breathlessness(10), and our findings suggest this may be the case even when the breathless person is not visible. The impact for caregivers who may be exposed to constant breathlessness could be important and therefore education and training for caregivers to better understand breathlessness is essential(13). It may also be interesting to further consider the effects of listening to breathlessness outside of the healthcare setting, for example exercise induced.

This study demonstrates that in the right circumstances a public experiment can enable efficient recruitment; this approach could be usefully applied to appropriate research questions in future. For this study we were able to exceed the planned sample size in three afternoons. We were able to conduct a rigorous study, with randomisation of recordings to eliminate the impact of recording order on the findings.

The limitations of this methodology were that it required a very brief experimental protocol and meant we were unable to collect sufficient data on prior experience of breathlessness/anxiety to investigate the impact of past experience on induced breathlessness. We were also unable to apply a control group or washout period between recordings. This has limited the inferences that can be drawn from our data and we cannot rule out that the findings from hypothesis 1), which suggest a dose-response effect of listening to breathlessness recordings, relate to the lack of washout period rather than true experience. Further, participants were mostly aged 18 – 35 and were attending an exhibition focused on anxiety. Therefore, results are unlikely to be representative of the general population or of those caring for people with chronic breathlessness who are likely to have regular exposure to the sound of breathlessness. Further work with more extensive data collection in more targeted populations could help to understand the clinical implications of our findings, for example how high levels of exposure to breathlessness among carers of people living with chronic breathlessness or anxiety could impact their own experience.

There were no clinically significant changes in participants' breathlessness based on a threshold of 0.5 on the NRS (14), however we used very short recordings to keep the total study duration down and this may have limited the impact on participants. A future study with longer exposure time might identify greater effects on breathlessness and would more closely reflect real life experience of those with breathlessness.

In conclusion this public experiment generated initial data showing that listening to breathlessness recordings increased participants' self-reported breathlessness. The cause of breathlessness appeared to mediate this effect. Further investigation is needed to determine mechanisms behind these findings, and the impact of prior experience of breathlessness. The clinical implications deserve further consideration.

Acknowledgements

We would like to thank the individuals who contributed recordings of their breathing, and St Christopher's Hospice for their role in identifying potentially suitable individuals. We would like to thank Science Gallery London for their support in setting up and running the experiment during the 'Sound of Anxiety' season and for facilitating the development of the online interface. Thanks to individuals from the Cicely Saunders Institute, King's College London (Anna Johnston, Anna E Bone, Cheng-Pei Lin, Lisa Brighton) who supported data collection –Thanks to Hamid Benalia for assisting with capturing and editing the recordings.

This study/project is supported by the National Institute for Health and Care Research (NIHR) Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

References

1. Moens K, Higginson IJ, Harding R. Are there differences in the prevalence of palliative care-related problems in people living with advanced cancer and eight non-cancer conditions? A systematic review. *J Pain Symptom Manage*. 2014;48(4):660-77.
2. Gysels MH, Higginson IJ. The lived experience of breathlessness and its implications for care: a qualitative comparison in cancer, COPD, heart failure and MND. *BMC palliative care*. 2011;10:15.
3. Currow DC, Dal Grande E, Ferreira D, et al. Chronic breathlessness associated with poorer physical and mental health-related quality of life (SF-12) across all adult age groups. *Thorax*. 2017;72(12):1151-3.
4. Malik FA, Gysels M, Higginson IJ. Living with breathlessness: a survey of caregivers of breathless patients with lung cancer or heart failure. *Palliat Med*. 2013;27(7):647-56.
5. Stoeckel MC, Esser RW, Gamer M, et al. Brain responses during the anticipation of dyspnea. *Neural plasticity*. 2016;2016.
6. Stoeckel MC, Esser RW, Gamer M, et al. Amygdala response to anticipation of dyspnea is modulated by 5-HTTLPR genotype. *Psychophysiology*. 2015;52(7):973-6.
7. Faull OK, Pattinson KT. The cortical connectivity of the periaqueductal gray and the conditioned response to the threat of breathlessness. *Elife*. 2017;6:e21749.
8. Faull OK, Subramanian HH, Ezra M, et al. The midbrain periaqueductal gray as an integrative and interoceptive neural structure for breathing. *Neuroscience & Biobehavioral Reviews*. 2019;98:135-44.
9. Marlow LL, Faull OK, Finnegan SL, et al. Breathlessness and the brain: the role of expectation. *Current opinion in supportive and palliative care*. 2019;13(3):200-10.
10. Herzog M, Sucec J, Van Diest I, et al. Observing dyspnoea in others elicits dyspnoea, negative affect and brain responses. *European Respiratory Journal*. 2018;51(4):1702682.
11. Vlemincx E, Sprenger C, Büchel C. Expectation and dyspnoea: the neurobiological basis of respiratory nocebo effects. *European Respiratory Journal*. 2021;58(3):2003008
12. Gift AG, Narsavage G. Validity of the numeric rating scale as a measure of dyspnea. *Am J Crit Care*. 1998;7(3):200-4.
13. Johnson MJ, Gozal D. Vicarious breathlessness: an inferential perceptual learned transposition process that may not be inconsequential to either patient or caregiver. *European Respiratory Journal*. 2018;51(4):1800306.
14. Ekström M, Johnson MJ, Huang C, et al. Minimal clinically important differences in average, best, worst and current intensity and unpleasantness of chronic breathlessness. *Eur Respir J*. 2020;56(2).

