

Pulmonary hypertension in lymphangioleiomyomatosis: characteristics in 20 patients

ELECTRONIC REPOSITORY

Authors: Vincent COTTIN, MD PhD (1), Sergio HARARI, MD (2), Marc HUMBERT, MD PhD (3), Hervé MAL, MD PhD (4), Peter DORFMÜLLER, MD PhD (5), Xavier JAIS, MD (3), Martine REYNAUD-GAUBERT, MD PhD (6), Grégoire PREVOT, MD (7), Romain LAZOR, MD (1,8), Camille TAILLÉ, MD (4), Jacques LACRONIQUE (9), Sabrina ZEGHMAR, MSc (1), Gérald SIMONNEAU, MD (3), Jean-François CORDIER, MD (1), and the Groupe d'Etudes et de Recherche sur les Maladies “Orphelines” Pulmonaires (GERM“O”P)

Table E1. Correlations between haemodynamics and pulmonary function

	<i>Mean PAP</i>	<i>Cardiac index</i>	<i>PVR</i>
FEV ₁	0.526 (0.021)	-0.351 (0.140)	0.593 (0.007)
FVC	0.309 (0.212)	-0.485 (0.042)	0.516 (0.028)
DLco	0.286 (0.266)	-3.346 (0.174)	0.657 (0.004)
Kco	0.530 (0.076)	-0.673 (0.016)	0.834 (< 0.001)
PaO ₂ at rest	0.286 (0.301)	-0.180 (0.301)	0.419 (0.094)
SpO ₂ (%) at end of 6-min walk test	-0.314 (0.219)	0.163 (0.533)	0.027 (0.918)

Data are presented as Pearson's correlation coefficient (p values). PAP: pulmonary arterial pressure; PVR: pulmonary vascular resistance; FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 s; DLco: single-breath diffusing capacity of the lungs for carbon monoxide; Kco: single breath transfer factor of the lungs for carbon monoxide; PaO₂: arterial oxygen tension; SpO₂: peripheral oxygen saturation. P<0.05 values are indicated in bold.