| | Included (n=724) | Excluded | P value |
|--|------------------|---------------------|---------|
| Age at onset, yr | 36.3±15.7 | 42.2±16.8 (n=640) | 0.001 |
| Body mass index (kg·m ^{-2}) | 23.81±3.28 | 24.07±3.40 (n=636) | 0.158 |
| Smoking (pack-years) | 5.8±9.8 | 6.9±11.6 (n=510) | 0.046 |
| Atopy, % positive | 44.8 | 52.6 (n=315) | 0.479 |
| PostBD FEV1 (%) | 83.45±19.95 | 82.46±22.16 (n=616) | 0.401 |
| | | | |

Table S1. Characteristics of included and excluded patients in COREA cohort.

Comparison between study groups is based on independent T-test for continuous variables and a χ^2 test for proportions. Data indicate means \pm SDs.

| | Included (n=1843) | Excluded (n=349) | P value |
|--|-------------------|---------------------|---------|
| Age at onset, yr | 38.2±17.3 | 39.9±17.3 (n=349) | 0.096 |
| Body mass index (kg·m ^{-2}) | 24.22±3.52 | 24.54±3.91 (n=320) | 0.134 |
| Smoking (pack-years) | 7.40±13.7 | 13.2±19.5 (n=54) | < 0.001 |
| Atopy, % positive | 59.4 | 60.6 (n=315) | 0.709 |
| PostBD FEV1 (%) | 79.86±22.42 | 79.37±22.14 (n=344) | 0.711 |
| | | | |

Table S2. Characteristics of included and excluded patients in SCH cohort.

Comparison between study groups is based on independent T-test for continuous variables and a χ^2 test for proportions. Data indicate means \pm SDs.