



Early View

Review

## **Pulmonary Complications of Bcr-Abl Tyrosine Kinase Inhibitors**

Jason Weatherald, Louise Bondeelle, Marie-Camille Chaumais, Christophe Guignabert, Laurent Savale, Xavier Jaïs, Olivier Sitbon, Philippe Rousselot, Marc Humbert, Anne Bergeron, David Montani

Please cite this article as: Weatherald J, Bondeelle L, Chaumais M-C, *et al.* Pulmonary Complications of Bcr-Abl Tyrosine Kinase Inhibitors. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.00279-2020>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©ERS 2020

## **Pulmonary Complications of Bcr-Abl Tyrosine Kinase Inhibitors**

Jason Weatherald\*, MD<sup>1</sup>, Louise Bondeelle\*, MD<sup>2</sup>, Marie-Camille Chaumais, PharmD, PhD<sup>3,4,5</sup>,  
Christophe Guignabert, PhD<sup>5,6</sup>, Laurent Savale, MD, PhD<sup>5,6,7</sup>, Xavier Jaïs, MD<sup>5,6,7</sup>, Olivier Sitbon,  
MD, PhD<sup>5,6,7</sup>, Philippe Rousselot, MD, PhD<sup>8</sup>, Marc Humbert, MD, PhD<sup>5,6,7</sup>, Anne Bergeron, MD,  
PhD<sup>2</sup>, David Montani, MD, PhD<sup>5,6,7</sup>

\* JW et LB contributed equally

*1 - University of Calgary, Department of Medicine & Libin Cardiovascular Institute of Alberta, Calgary, Alberta, Canada*

*2 – Université de Paris, Assistance Publique - Hôpitaux de Paris (AP-HP), Service de Pneumologie, Hôpital Saint-Louis, Paris, France*

*3 - Université Paris-Saclay, Faculté de Pharmacie, Université Paris-Saclay, Châtenay-Malabry, France.*

*4 - Assistance Publique - Hôpitaux de Paris (AP-HP), Service de Pharmacie, Hôpital Bicêtre, Le Kremlin-Bicêtre, France.*

*5 - INSERM UMR\_S 999 «Pulmonary Hypertension: Pathophysiology and Novel Therapies», Hôpital Marie Lannelongue, Le Plessis-Robinson, France*

*6 - Université Paris-Saclay, School of Medicine, Le Kremlin-Bicêtre, France*

*7 - Assistance Publique - Hôpitaux de Paris (AP-HP), Department of Respiratory and Intensive Care Medicine, Pulmonary Hypertension National Referral Center, Hôpital Bicêtre, Le Kremlin-Bicêtre, France*

*8 – Service d’Hématologie, Centre hospitalier de Versailles André Mignot, Le Chesnay, France*

### **Corresponding author:**

Montani David, MD, PhD

Centre de Référence de l’Hypertension Pulmonaire, Service de Pneumologie, Hôpital Bicêtre, Assistance Publique Hôpitaux de Paris

78 rue du Général Leclerc, 94270 Le Kremlin-Bicêtre, France.

Tel: (33) 1 45 21 79 72; Fax: (33) 1 45 21 79 71;

e-mail: david.montani@aphp.fr

**Running head:** Pulmonary complications of tyrosine kinase inhibitors

## **ABSTRACT**

Tyrosine kinase inhibitors (TKIs) targeting the Bcr-Abl oncoprotein revolutionized the treatment of chronic myelogenous leukemia. Following the success of imatinib, second and third generation molecules were developed. Different profiles of kinase inhibition and off-target effects vary between TKIs, which leads to a broad spectrum of potential toxicities.

Pulmonary complications are most frequently observed with dasatinib but all other Bcr-Abl TKIs have been implicated. Pleural effusions are the most frequent pulmonary complication of TKIs, usually associated with dasatinib and bosutinib. Pulmonary arterial hypertension is an uncommon but serious complication of dasatinib, which is often reversible upon discontinuation. Bosutinib and ponatinib have also been associated with pulmonary arterial hypertension, while imatinib has not. Rarely, interstitial lung disease has been associated with TKIs, predominantly with imatinib.

Mechanistically, dasatinib affects maintenance of normal pulmonary endothelial integrity by generating mitochondrial oxidative stress, inducing endothelial apoptosis and impairing vascular permeability in a dose-dependent manner. The mechanisms underlying other TKI-related complications are largely unknown. Awareness and early diagnosis of the pulmonary complications of Bcr-Abl TKIs is essential given their seriousness, potential reversibility, and impact on future treatment options for the underlying chronic myelogenous leukemia.

**Key-words:** tyrosine kinase inhibitor; pulmonary; side-effect; complication; adverse effects

## KEY MESSAGES

1. Bcr-Abl tyrosine kinase inhibitors have been associated with certain pulmonary complications, including exudative pleural effusions, chylothorax, interstitial lung disease and pulmonary arterial hypertension. Due to its broader range of off-target protein kinase inhibition and its effects on endothelial cell function, dasatinib is the tyrosine kinase inhibitor most frequently associated with pleural effusion and pulmonary arterial hypertension.
2. Clinicians and patients should be aware of the potential adverse pulmonary effects of tyrosine kinase inhibitors prior to initiation. Unexplained respiratory symptoms require further evaluation including chest x-ray and/or computed tomography of the chest, transthoracic echocardiography, pulmonary function tests and cardiac biomarkers such as NT-proBNP.
3. Many pulmonary complications are reversible upon discontinuation of the offending medication and usually do not recur if another tyrosine kinase inhibitor is introduced. However, multidisciplinary management and close clinical follow-up is necessary to ensure resolution and detect early recurrences.

## INTRODUCTION

Chronic myeloid leukemia (CML) is a myeloproliferative disorder characterized by a cytogenetic hallmark in the Philadelphia chromosome and its associated t(9;22)(q34;q11.2) translocation. This translocation results in a *BCR-ABL1* fusion gene which codes for a Bcr-Abl1 oncoprotein with constitutively active tyrosine kinase activity[1]. The ability of Bcr-Abl to induce a similar disease in mice confirmed its pathogenic role and led to the development of tyrosine kinase inhibitors (TKIs), a new class of anticancer agents. The discovery and clinical use of TKIs, led by imatinib and subsequently enriched with second-generation (dasatinib, nilotinib, and bosutinib) and third-generation (ponatinib) TKIs, has revolutionized the treatment of chronic CML[2]. In the TKI era, a patient diagnosed with CML in the chronic phase may anticipate a normal life expectancy[3]. Thus, the prevalence of CML is increasing in western countries. Most patients will experience a long-enduring treatment with an “optimal” or an intermediate “warning” response as defined by European LeukemiaNet criteria[4]. In this situation, no difference in survival can be demonstrated between imatinib and second generation TKIs used as first-line[5, 6]. In patients who achieve a sustained deep molecular response, defined by a Bcr-Abl1<sup>IS</sup> quantification less than 0.01% for at least 1 year after a treatment duration of at least 3 years, more than half can successfully stop their TKI to enter a treatment-free remission period[7]. Second generation TKIs may achieve faster and deeper molecular response than imatinib and are increasingly proposed as first line options for patients, despite the recognized higher risk of adverse events, including notable pulmonary complications. The objectives of this narrative review are to summarize current knowledge on the epidemiology, clinical features, pathophysiology and management of pulmonary complications of Bcr-Abl TKIs.

## **SEARCH STRATEGY**

References included this narrative review were identified by searching PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/>) for articles published up to June 1, 2019 and an updated search was done on April 21, 2020, using the terms “pulmonary hypertension”, “pleural effusion”, “pneumonitis”, OR “interstitial lung disease”, AND “kinase inhibitor”, “bcr-abl”, “imatinib”, “dasatinib”, “nilotinib”, “bosutinib”, or “ponatinib”, which resulted in 2190 articles. Articles resulting from these searches and relevant references cited in those articles were screened. Articles in English or French were reviewed in full. Articles in other languages with abstracts in English were reviewed if sufficient detail was present in the abstract. Articles relating to non-Bcr-Abl protein kinase inhibitors were not evaluated.

## **Bcr-Abl TYROSINE KINASE INHIBITORS**

Imatinib is the first Bcr-Abl protein tyrosine kinase inhibitor (TKI) developed for CML. Imatinib is a competitive inhibitor of adenosine triphosphate (ATP) targeting the inactive form of Bcr-Abl. When bound in the catalytic site in place of ATP, imatinib maintains Bcr-Abl in its inactive form inhibiting both proliferation and promoting apoptosis of Bcr-Abl cells[8]. However, mutations in the *Abl* gene leading to imatinib inefficacy or resistance have been observed. These mutations involve modifications on the catalytic site and decrease imatinib binding. More than 80 specific types of substitutions have been reported of which T315I mutation is the most resistant. Therefore, a second generation of Bcr-Abl TKIs (dasatinib, nilotinib, and bosutinib) were developed, followed by a third generation with ponatinib. Nilotinib and ponatinib have the same mechanism of action as imatinib in blocking the inactive form of Bcr-Abl (type II inhibitor), whereas dasatinib and bosutinib block the active form (Type I inhibitors)[9]. Dasatinib, nilotinib

and bosutinib have a higher potency than imatinib and could be active on the majority of imatinib of Bcr-Abl resistant cells[10]. However, only ponatinib is capable in the case of a T315I mutation[11].

Imatinib, nilotinib, bosutinib and dasatinib are indicated in the first-line treatment of CML in the chronic phase[12]. Regarding benefit, there is no significant survival difference reported between these three TKIs. However, in France for example, due to an insufficient clinical benefit judged by authorities, dasatinib is not reimbursed as a first-line treatment for CML. In addition to the issue of reimbursement, choice of TKI should be based on a patient's characteristics (age, comorbidities, associated medications, etc.) and the risk of adverse events for each TKI. Bcr-Abl TKIs have many potential adverse effects. Some of them are common, such as skin and subcutaneous tissue reactions or hepato-biliary disorders. However, certain TKIs have a predilection for causing specific complications. For example, dasatinib can promote the development of pleural effusions, pulmonary arterial hypertension (PAH) and hemorrhagic complications; nilotinib may induce metabolism and nutrition disorders and vascular events, and ponatinib is associated with venous and arterial thrombosis. The characteristics of the Bcr-Abl TKIs are detailed in **Table 1**.

Despite their monikers as Bcr-Abl targeted therapies, imatinib, nilotinib, dasatinib, bosutinib and ponatinib are actually multi-kinase inhibitors, with off-target effects depending on the particular drug and dosage. The spectrum of off-target inhibition, in addition to Bcr-Abl, ranges from 6 (nilotinib) to 28 (dasatinib) other protein targets (**Table 2**). For example, imatinib also blocks the platelet derived growth factor (PDGF) receptors and the mast/stem cell growth factor receptor Kit; dasatinib inhibits the proto-oncogene tyrosine-protein kinase Src, and

ponatinib inhibits the vascular endothelial growth factor receptor 2 and fibroblast growth factor receptors 1 to 4[13]. The other targets for each Bcr-Abl TKI are depicted in **Table 2**. Modulation of these targets may have a clinical impact in terms of tolerance of each drug and need to be known by prescribing clinicians. It is important to emphasize that since TKIs are multi-kinase inhibitors, off-target inhibition of Bcr-Abl TKIs overlaps with those of TKIs used for other indications. For example, the Bcr-Abl TKIs used to treat CML share several off-target kinase inhibitory with epidermal growth factor receptor [EGFR] inhibitors used to treat non-small cell lung cancer (e.g. erlotinib, gefitinib) [14]. Examination of overlapping inhibition profiles could potentially shed some light on the underlying mechanisms of toxicities common to both groups of Bcr-Abl and non-Bcr-Abl TKIs, such as interstitial lung disease (ILD) or pneumonitis[15, 16].

### **PATHOPHYSIOLOGY OF Bcr-Abl TKI TOXICITIES**

In terms of pathophysiology of pulmonary toxicity, most of the available data involves dasatinib. Dasatinib is a highly effective multi-kinase inhibitor approved for first-line treatment of Philadelphia-positive CML and for all phases of Philadelphia-positive CML with resistance or intolerance to prior therapy, including imatinib. However, pleural effusions and PAH are two undesirable pulmonary effects observed with dasatinib use[17–22]. Although many of the exact cellular events and specific signaling pathways are incompletely understood, these side effects are often partially or completely reversible after drug withdrawal. As a result, we lack effective marker-based screening strategies and mechanism-based treatment options for PAH occurring in dasatinib-treated patients. In addition, a better understanding of mechanisms involved in



dasatinib-induced PAH would not only help to ameliorate the management of dasatinib-treated patients, but also would improve our knowledge about the pathophysiologic mechanisms underlying the development of PAH. An understanding the mechanisms of cardiopulmonary toxicities is also critical for developing future evidence-based preventive strategies[23, 24]. Similarly, mechanisms underlying TKI-related ILD remain unclear. Drug hypersensitivity reactions or a pharmacological effect of tyrosine kinase inhibition are possibilities but there are no mechanistic data available.

#### Endothelial cell injury and dysfunction

The maintenance of the integrity of the endothelial lining is essential for a myriad of biological molecules and functions under its control[25], and some *in vivo* and *in vitro* evidence indicates that certain TKIs might contribute to the weakening of some specific endothelial functions. For example, rats treated with high doses of dasatinib exhibit increases in Evans blue dye extravasation into the lung parenchyma, supporting that dasatinib can alter endothelial integrity in rat lungs[22]. Consistent with these *in vivo* observations, it has been demonstrated that dasatinib leads to a rapid and reversible increase in paracellular permeability in monolayers of human pulmonary endothelial cells or umbilical vein endothelial cells (HUVECs)[22, 26]. Indeed, when confluent endothelial monolayers are exposed to increasing doses of dasatinib, there is an increase in the passage of macromolecules, loss of VE-cadherin and *zonula occludens* (ZO)-1 from cell-cell junctions, and development of actin stress fibers[22]. These phenomena are not found with imatinib[22]. Remarkably, this increase in endothelial permeability was dependent on reactive oxygen species (ROS) production induced by dasatinib.

Indeed, co-treatment with the antioxidant agent N-acetyl-L-cysteine (NAC) was found to protect pulmonary endothelial cells against this loss of integrity both *in vivo* and *in vitro*[22]. Using immunofluorescence and confocal microscopy analyses, these authors also noted dilation of the lymphatic vessels in rat lungs treated with high doses of dasatinib compared to vehicle or high-dose imatinib-treated rats[22].

It is an open question whether these specific effects of dasatinib on pulmonary vessels are related to vascular cell sensitivity to oxidative stress or to their specific expression/activity of certain protein kinases. Nevertheless, it has been demonstrated that chronic dasatinib therapy causes pulmonary endothelial damage in humans and rodents[21]. Indeed, at high dose, dasatinib treatment attenuated hypoxic pulmonary vasoconstriction responses and increased susceptibility to experimental pulmonary hypertension (PH) in rats[21]. It has been shown that pretreatment of rats with high doses of dasatinib, but not imatinib, increases their response to known inducers of PH, which closely mimics the findings observed in humans[21]. It has been shown that dasatinib induces pulmonary endothelial cell apoptosis in a dose-dependent manner, while imatinib does not. Among the processes of endothelial damage, induction of ER stress, and mitochondrial ROS production were shown to play a central role (**Figure 1**), a phenomenon that was demonstrated to be independent of Src family kinases[21]. High levels of ROS cause damage to proteins, nucleic acids, lipids, and membranes, but also cause DNA damage. When repair of DNA damage is sufficient, this can result in altered transcription of specific vital genes, followed by cellular dysfunction and, ultimately, apoptosis[27]. Consistent with these findings, elevations in markers of endothelial dysfunction and vascular damage are more abundant in the serum of CML patients who were treated with dasatinib, compared with

CML patients treated with imatinib[21]. Furthermore, these data are consistent with a report from Daccord and colleagues who described the presence of irreversible pulmonary vascular lesions in explanted lungs of a case of severe dasatinib-induced PAH[28].

### Smooth muscle ion channels and regulation of vascular tone

Other mechanisms for the pulmonary complications of TKIs have been proposed such as inhibition of different potassium ( $K^+$ ) channels[29], activation of the Rho kinase pathway[22, 26] or alterations of smooth muscle contractions and dysfunction[29–33]. However, further studies are required to elucidate the underlying mechanisms of pulmonary vascular dysfunction, as several TKIs such as imatinib [30, 32, 33], nilotinib[30, 32] and sorafenib[32] have been shown to have potent vasorelaxant effects. In a recent study, Rieg et al. reported that the vasorelaxation mechanisms of imatinib and nilotinib in precision-cut lung slices (PCLS) are linked to the generation of cAMP and mainly dependent on activations of  $K_{ATP}$ -,  $BK_{Ca}^{2+}$ - and  $K_v$ -channels[30]. Nagaraj and colleagues reported that TWIK-related acid-sensitive potassium channel 1 (TASK-1, encoded by KCNK3) and Src co-localize in the plasma membrane of human pulmonary artery smooth muscle cells (PA-SMCs) and that Src is required for the functioning of TASK-1 channels[28]. In isolated perfused and ventilated rat lungs, Fazakas and colleagues also found that 100 nM of dasatinib increased pulmonary pressure, a phenomenon totally abolished by the Rho-kinase inhibitor Y27632 (10  $\mu$ M). However, co-administration of Y27632 with 100 nM of dasatinib did not have a positive effect on pulmonary edema in this *ex vivo* model[25]. These observations are consistent with the results of Phan and colleagues who have

demonstrated that this endothelial cell permeability is not sufficiently explained by ROCK activation or Lyn inhibition[21].

#### Other proposed mechanisms for TKI toxicities

The signaling mechanisms underpinning TKI toxicities are complex, and may include alterations in the BMP [34], NF- $\kappa$ B [35, 36], and Akt/mTOR [37, 38] signaling pathways. However, it has also been shown that the dose, dosing regimen and interpatient variability in drug exposure play a role in the occurrence of some of these undesirable effects of TKIs[39–43], strongly supporting the concept that a pharmacokinetic-guided dose individualization strategy may help to improve their safety profile. In a Phase III dose finding trial, Shah *et al.* have indeed shown that the incidence of pleural effusions was lower in patients receiving a dose of 100 mg once daily than in those given 70 mg twice-daily dosing regimen[44]. Consistent with these observations, it has been reported that rats treated with high dose of dasatinib ( $10 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ , which is the equivalent to 10 $\times$  the clinically relevant dose) developed pleural effusion over the period of 8 weeks, starting from week-5[20]. Remarkably, this phenomenon was not observed in rats treated with dasatinib at the clinically relevant dose (1 $\times$ ) or in rats treated with high dose of imatinib[21]. This dose–response relationship observed in humans and rats strongly suggests that these adverse events are likely to be due to the loss/inhibition of some of the low affinity target kinases of dasatinib, either alone or in combination with inhibition of Src or other Src family kinases such as Lyn, Yes, and Fyn[34, 45, 46].

In summary, it is likely that several overlapping mechanisms contribute to TKI toxicities, however direct cell toxicity, endothelial dysfunction, excess cellular levels of ROS, or indirect inflammatory mediated events are strongly suspected to play a role.

## **PLEURAL EFFUSIONS AND INTERSTITIAL LUNG DISEASES**

### *Epidemiology and Clinical Features*

Pleural effusions are the most common respiratory complications that may complicate the administration of TKIs. Their incidence varies according to the TKI, with pleural effusions being most frequently associated with dasatinib. Due to the more recent approval of both bosutinib and ponatinib, fewer data exist regarding their long-term toxicity profiles, including respiratory complications. A better knowledge of the incidence, the risk factors and the clinical characteristics of TKI-induced pleural effusions allowed the optimization of their management over time. Although rarer, interstitial lung diseases have also been associated with TKI, with more or less acute clinical presentation and corresponding to various histological patterns. Both effusions and interstitial diseases should be suspected with the onset of nonspecific respiratory symptoms such as dyspnea, cough or chest pain, or they may be diagnosed fortuitously on follow-up chest imaging.

### Dasatinib

The occurrence of pleural effusions complicating the administration of dasatinib has been described in the earliest studies evaluating the efficacy of dasatinib in CML[47]. Further studies confirmed that up to 28% of patients treated with dasatinib for CML for a 5-year period and

33% for a 7-year period developed pleural effusions[5, 19] with an incidence of new cases that was similar each year[5, 48]. Dasatinib regimen, including both dosage and the number of dose per day, was quickly noted to be associated with the occurrence of pleural effusions[49], leading to a modification of drug administration to 100 mg once daily. A recent study further found a 6% incidence of pleural effusion in newly diagnosed chronic-phase CML treated with dasatinib 50 mg/day[50]. Other risk factors were identified for TKI-induced pleural effusions such as CML phase (with an increase in patients with advanced phases)[51], presence of a cardiac history, hypertension, hypercholesterolemia, and a history of autoimmune disease[52–54]. The plurality of risk factors found probably accounts for different aetiologies of effusions in addition to dasatinib-induced, such as cardiac dysfunction, infectious or malignant causes, including related to the underlying hematological disease. However, in clinical studies, the most significant risk factor for developing pleural effusions is age, which was associated with dasatinib exposure (i.e. high minimum concentration value, C<sub>min</sub>)[5, 42, 48]. Of note, pleural effusion-associated with dasatinib was not found to be associated with fluid retention[52]. Pleural effusions can be unilateral or bilateral and can be of variable size ranging from minimal to large volume. Analysis of pleural fluid mostly demonstrated exudates with a predominance of lymphocytes; however, some transudates have also been reported[53, 55]. Chylothorax has also been reported[53, 55–58]. The incidence of chylothorax has probably been overlooked because pleural fluid has not always been macroscopically chylous and the diagnosis of chylothorax requires systematic pleural fluid triglyceride quantification. A case of fibrothorax, characterized by a diffusely thick firm fibrous pleura on pleurectomy, has been reported in association with dasatinib following a history of recurrent chylothoraces[59]. In some cases,

pleural effusion was reported to be associated with other side effects such as pericardial effusion, interstitial pneumonia, or pulmonary hypertension[5, 18, 53, 55]. Overall response to dasatinib, progression-free survival, and overall survival were similar between patients who developed pleural effusion and those who did not[48].

More rarely, dasatinib has been associated with parenchymal lung abnormalities either alone or associated with pleural effusion (**Figure 2**)[53, 60–63]. In a retrospective study, Bergeron et al. identified that 9 of 40 (22.5%) patients with chronic-phase CML who received dasatinib developed lung abnormalities[53]. Among them, seven patients had parenchymal abnormalities such as ground glass opacities, consolidations and/or septal thickening associated with pleural effusions for four of these patients. Respiratory symptoms included dyspnea, cough, and chest pain with a median time between dasatinib treatment initiation and respiratory symptoms of 229 days (range, 20–510 d). Two patients presented with fever and myalgia. Bronchoalveolar lavage shows lymphocytic alveolitis in most cases[53, 63]. After dasatinib interruption, lung parenchymal manifestations usually resolve without treatment but corticosteroids have been instituted in some cases [53, 61–63]. Pneumonitis relapsed in only one of the patients in whom dasatinib was reintroduced at a lower dose[53].

### Imatinib

Although peripheral and periorbital edema are the most frequent imatinib-associated adverse events, occurring in 60% of the patients, pleural effusions were reported in only 0-0.8% of the patients treated with imatinib for 5 years[5, 64]. Few cases of pleural effusion, mostly in association with pericardial effusion, have been reported, often in a context of peripheral fluid

retention[65, 66]. Several cases of imatinib-associated interstitial lung diseases have been reported[67–71]. Ohnishi and colleagues reported 27 cases of drug-induced interstitial disease (ILD) associated with imatinib[71]. All patients presented with dyspnea and 16 had hypoxemia with a median time from imatinib initiation of 49 days (range: 10–282 days). The median daily dose of imatinib was 400 mg (range: 200–600 mg). The authors found no clear correlation between the development of imatinib-related pneumonitis and either the dose or duration of imatinib therapy. CT scan patterns include diffuse ground glass opacities, nodules consolidative patterns consistent with organizing pneumonia, with or without signs of pulmonary fibrosis. Transbronchial biopsies found nonspecific histologic patterns including inflammation and fibrosis. Imatinib was discontinued in all patients and 24 patients were treated with corticosteroids. After treatment, ILD either resolved completely (26%), improved (59%), did not improve (15%) while one patient suffered from pulmonary fibrosis. After ILD improved, imatinib was reintroduced at a lower dose (range: 100–400 mg/day) in 11 patients leading to recurrence in four patients. Another case, in which imatinib was reintroduced, develop irreversible pulmonary fibrosis[70]. A case of pulmonary alveolar proteinosis, characterized by accumulation of alveolar proteinaceous material and a ‘crazy paving’ pattern on chest CT, has been reported in association with imatinib, which was exacerbated when switching to dasatinib[72].

### Nilotinib

Nilotinib-associated pulmonary side effects have rarely been reported[73]. Only scattered case reports of pleural effusion and pneumonitis have been published[73–75]. In a retrospective



study in 120 patients receiving nilotinib as second-line therapy, pleural effusion was reported in 2% of patients[73]. A case of biopsy-proven organizing pneumonia was described 3 years after nilotinib initiation[74]. The patient presented with cough for 4 weeks, without other respiratory symptoms, and alveolar consolidation in both lower lobes on CT scan. After improvement following corticosteroid administration, nilotinib was reintroduced at a lower dose with no pneumonitis recurrence.

### Bosutinib

The incidence of bosutinib associated pleural effusions was less than 5% of the patients treated up to 5 years with bosutinib[76, 77]. Pleural effusions were more frequently reported in older patients[77, 78]. One case of bosutinib-related pneumonitis associated with pleural effusion has also been reported[79]. In this case, the CT pattern suggested organizing pneumonia and the lung histology confirmed organizing pneumonia with fibrinous exudate, interstitial eosinophilic infiltrate and small pulmonary arteries abnormalities[79].

### Ponatinib

No cases of pleural effusions or interstitial pneumonitis have been specifically associated with ponatinib to date, with the exception of one case of interstitial lung disease reported in the Phase 1 study evaluating the effect of ponatinib in refractory CML[11]. Since ponatinib and bosutinib are newer and less widely used, the actual risks of pulmonary complications are likely not yet known.

### *Management of TKI-associated pleural effusion and interstitial lung disease*

The treatment of TKI associated-pulmonary side effects should take into account the severity of the respiratory involvement but also the control and prognosis of the underlying CML, including the available therapeutic alternatives. This management plan must involve a close collaboration with hematologists and should be discussed on a case by case basis.

As different causes of pleural effusions have been identified, detailed investigations are required. If the effusion has a sufficient volume, a diagnostic thoracentesis should be performed to differentiate exudate from transudate, to culture and exclude pleural infection, and to analyze the cellularity differential to narrow the differential diagnosis. The administration of empiric diuretics and / or corticosteroids has been reported[80]; however, this strategy has never been evaluated in an interventional study. It is likely that the efficacy of any treatment will depend on the underlying cause of the pleural effusion. Specific management will be different depending on the size and consequences of the pleural effusion. In cases with minimal fluid, continuation of the TKI with close clinical and radiological monitoring may be considered, whereas in cases of moderate or large pleural effusions, dose reduction, withdrawal or change of TKI should be considered according to alternative CML treatment options. However, in a recent study, dasatinib dose reduction after a first episode of pleural effusion did not prevent recurrence of this adverse event[80]. In the 5-year analysis from a phase III Dasatinib Versus Imatinib Study in CML (DASISION) trial, 22% of the patients who developed pleural effusion required drug interruption[5]. Therapeutic thoracentesis may rarely be necessary[5]. In an uncontrolled case series of 6 patients with refractory dasatinib-induced pleural effusions and severe dyspnoea, Aoyama et al. reported that tolvaptan, a

vasopressin V2-antagonist, in addition to diuretics improved symptoms and pleural effusions in 5 of 6 patients, such that dasatinib could be continued or reintroduced in these patients[81]. Therapeutic drug monitoring of TKI may be helpful to minimize the incidence of pleural effusions but is not widely used in practice[82].

Of note, patients who developed dasatinib-associated pleural effusion and subsequently received bosutinib may develop the same side-effect with bosutinib[76, 83]. In a retrospective study including 20 CML patients developing pleural effusion during dasatinib and who subsequently received bosutinib, only 30% had recurrence of pleural effusion[84].

The occurrence of pulmonary parenchymal opacities in a patient treated with TKIs must also lead to detailed assessment. Since ILD appears to be a rare complication of TKIs, other causes such as atypical infections, concomitant medications, exposures, and other unrelated medical conditions should be considered. The evolution of TKI related pneumonitis appears to be favorable with spontaneous resolution with TKI withdrawal[67, 79] or after corticosteroids[53, 69, 70, 74]. Any re-challenge with the suspected offending TKI should be considered according to the perceived risk-benefit balance for the individual patient. Based on limited data, imatinib-associated interstitial lung disease may not recur when switched to nilotinib[85, 86].

## **PULMONARY ARTERIAL HYPERTENSION**

### *Epidemiology*

PAH is characterised by elevated mean pulmonary arterial pressure (mPAP >20 mmHg) and pulmonary vascular resistance (PVR  $\geq$ 3 Wood units) with normal left heart filling pressures (pulmonary artery wedge pressure [PAWP]  $\leq$ 15 mmHg)[87]. The first case reports of reversible PAH with dasatinib were published in 2009[88, 89]. Then, in 2012, Montani et al. described a

series of 9 patients with severe PAH (mPAP ranged from 30-59 mmHg) in patients treated with dasatinib, which was reversible in some cases following dasatinib discontinuation[16]. From the Montani et al. study, the estimated minimum annual incidence of PAH in dasatinib-treated patients in France was 0.45%[16]. However, a recent multi-centre study of 212 dasatinib-treated patients from Australia found a much higher estimated prevalence of pulmonary hypertension on echocardiogram at 5%, although none of them had confirmation with right heart catheterization[90]. There were over 440 cases of PAH associated with protein kinase inhibitors reported to the World Health Organisation's pharmacovigilance database (Vigibase®) as of December 31, 2017[34]. Dasatinib is by far the most frequently implicated Bcr-Abr TKI associated with PAH and is now considered a "definite" cause of drug-induced PAH in the recent 6<sup>th</sup> World Symposium on PH[87]. New onset PAH and/or PAH worsening has been reported with other Bcr-Abl TKIs such as bosutinib[17, 91, 92], ponatinib[93], nilotinib[94]. However, the association between these molecules and PAH is less established than for dasatinib, as many patients in these reports were incompletely characterized with right heart catheterization or had been previously treated with dasatinib. In the Vigibase® study by Cornet et al., the reporting odds ratio (ROR: the proportion of PAH cases associated with a TKI compared to the proportion of PAH cases associated with all other drugs) was highest for dasatinib (28.6), followed by bosutinib (13.4) and ponatinib (3.88)[34]. Imatinib, the first most frequently used TKI for CML, has not been associated with the development of PAH. In fact, preclinical data and a phase II study suggested that imatinib may be an effective treatment for PAH[95]. However, despite improvements in exercise capacity and hemodynamics in the phase III randomized, placebo-controlled trial, the safety profile of imatinib was unacceptable with

high rates of discontinuation and unexpected adverse events compared to placebo, including subdural hematoma[96].

### *Clinical Features and Outcomes*

In a long-term follow-up study of 21 patients from the French PH Registry with right heart catheterization-confirmed dasatinib-induced PAH, 71% of patients were female and the median age was 52 years old[18]. The median exposure to dasatinib before PAH diagnosis was 42 (8–74) months. There were concurrent pleural effusions in 62% and a pericardial effusion was present in six patients (29%). All patients had dasatinib discontinued and 11/21 received PAH targeted therapies such as sildenafil, bosentan or calcium channel blockers, with significant improvements in clinical and haemodynamic parameters at the time of a follow-up right heart catheterization (**Figure 3**). Interestingly, although the median mPAP decreased from 45 to 26 mmHg and PVR decreased from a median of 6.1 to 2.6 Wood units, there was persistent PAH in 37% of cases[18]. Furthermore, two patients had abnormally haemodynamic exercise responses consistent with exercise PH, likely indicating some residual pulmonary vascular disease. Up to 68% of PAH patients have concurrent pleural effusions, which is much higher than the 25% rate in dasatinib clinical trials in CML[97].

While reversible cases could be caused by intense pulmonary arterial vasoconstriction, the frequent persistence of PAH suggests increased susceptibility to the development of irreversible pulmonary arterial remodelling in some dasatinib-treated patients. This was subsequently demonstrated in a histological analysis of explanted lungs from a patient with severe, progressive dasatinib-induced PAH who required lung transplantation[27]. Typical PAH

histopathological features of severe arteriopathy such as arteriolar obliteration, plexiform lesions, medial hypertrophy and intimal thickening were found in this patient. Two recent case reports lend further credence to a 'second-hit' hypothesis of dasatinib-induced PAH in susceptible humans. The first case described dasatinib-induced PAH in a patient with systemic sclerosis[98], a disease in which patients are predisposed to PAH, with up to a 24% lifetime risk[99]. Another reported reversible PAH during dasatinib treatment for CML in the context of an occult anomalous pulmonary venous connection and sinus venosus atrial septal defect, a lesion associated with pulmonary vascular disease but which more likely contributed to dasatinib-induced PAH in light of the complete reversibility in this case[100]. Of the 62 right heart catheterization-confirmed cases from the two largest studies of dasatinib-induced PAH, no deaths directly attributable to PAH were reported[18, 101]. Survival at 1, 3, and 5-years in the French registry study were 90.5%, 90.5% and 85.7%, respectively, which is very similar to long-term survival in randomized trials of dasatinib for CML[5, 18, 19]. However, one case of fatal dasatinib-induced PAH in a 36-year old woman has been reported[102].

#### *Diagnosis and Management of TKI-associated PAH*

Given the relatively low incidence of symptomatic PAH in TKI-treated patients, systematic screening in asymptomatic patients is not practical and is not recommended. When possible and feasible, a baseline echocardiogram may be performed prior to initiating dasatinib as a reference, in order to document the presence or absence of pre-existing pulmonary hypertension. In patients who have unexplained dyspnoea and/or symptoms of right heart dysfunction in the context of TKI treatment, the first step is to assess the likelihood of

pulmonary hypertension by echocardiogram and to perform a chest x-ray, pulmonary function tests, and cardiac biomarkers (such as troponin or N-terminal pro-brain natriuretic peptide [NT-pro-BNP]) to evaluate other potential explanations such as pleural effusions, interstitial or obstructive lung diseases, pneumonia, ischaemic heart disease, or left-sided heart failure (**Figure 4**). According to current PAH guidelines, if an echocardiogram demonstrates an elevated tricuspid regurgitation velocity (TRV)  $>3.4$  m/s, or if the TRV is  $< 3.4$  m/s with secondary signs of PH (such as flattening of the interventricular septum or right heart chamber dilation), then a right heart catheterization is recommended. Other causes of pulmonary hypertension, such as parenchymal lung disease or chronic thromboembolism, should be excluded with a high resolution computed tomography scan and ventilation-perfusion scan, respectively[103]. Although a 6% risk of venous thromboembolic disease has been associated with ponatinib[104], no cases of chronic thromboembolic pulmonary hypertension have been reported in association with TKIs. It must be emphasized that right heart catheterization is crucial for proper diagnosis and to delineate the haemodynamic profile, since patients with TKI-induced PAH may have concurrent bilateral pleural effusions that would otherwise and erroneously suggest post-capillary pulmonary hypertension due to left heart disease. Patients receiving TKIs may also have post-capillary pulmonary hypertension due to elevated left heart pressure (mPAP  $>20$  mmHg and PAWP  $>15$  mmHg) or they may have elevated pulmonary artery pressure due to a hyperdynamic circulation from anaemia or infection, which is characterized by high cardiac output and normal PVR.

In such cases where right heart catheterization confirms post-capillary pulmonary hypertension or hyperdynamic state, it is not necessary to stop the TKI. However, if PAH is confirmed, the

offending TKI should be discontinued immediately.

For patients with intermediate or high-risk features such as severe symptoms (NYHA III-IV), severe haemodynamic derangement (cardiac index  $< 2.5 \text{ L}\cdot\text{min}^{-1}\cdot\text{m}^{-2}$ ), or clinical signs of right heart failure, PAH therapies should be instituted in concordance with PAH treatment guidelines[18, 103, 105]. The currently available treatment options for PAH include phosphodiesterase type-5 inhibitors (e.g. sildenafil, tadalafil), soluble guanylate cyclase stimulators (e.g. riociguat), endothelin receptor antagonists (e.g. bosentan, macitentan), prostacyclin analogues (e.g. epoprostenol, treprostinil) and oral prostacyclin receptor agonists (e.g. selexipag). There have been several reports of successful treatment of TKI-induced PAH with initial combination therapy of an endothelin receptor antagonists and a phosphodiesterase type-5 inhibitor[18, 106, 107]. For low-risk patients with mild symptoms (NYHA I-II), no right heart failure, and low risk hemodynamics (right atrial pressure  $< 8 \text{ mmHg}$  and cardiac index  $> 2.5\text{--}3 \text{ L}\cdot\text{min}^{-1}\cdot\text{m}^{-2}$ ), expectant management without PAH therapies and discontinuation of the TKI can be considered[18]. However, short-term re-evaluation in 3-4 months should be performed in all patients regardless of whether PAH therapies are started, and should include invasive haemodynamic reassessment wherever possible, given the high rate of persistent PAH[108]. A multidisciplinary approach is necessary to determine which alternative TKI is appropriate and when it should commence, depending on severity of PAH and status of the underlying CML. Dasatinib should not be re-introduced and patients who are switched to bosutinib or ponatinib should be closely monitored given the possibility of worsening or recurrent PAH in this setting[17, 92, 93].



## **CONCLUSIONS**

The design and clinical development of Bcr-Abl tyrosine kinase inhibitors have revolutionized the treatment and prognosis of patients with chronic myelogenous leukemia. Nevertheless, these molecules are not specific for Bcr-Abl and have effects on numerous other protein kinases and non-protein kinases. Serious and potentially fatal pulmonary complications can occur from Bcr-Abl TKI therapies, including pleural effusions, ILD and PAH, which requires a comprehensive evaluation and usually requires discontinuation of the TKI. Unexplained respiratory symptoms in a patient receiving a TKI should prompt further investigations including chest x-ray, pulmonary function testing, natriuretic peptides such as NT-proBNP, transthoracic echocardiography, and computed tomography of the chest. Knowledge of the adverse effect profiles of each Bcr-Abl TKI can be helpful for clinicians since, for example, PAH is most frequently associated with dasatinib but has not been associated with imatinib. Thus, the finding of pulmonary hypertension in a patient taking imatinib should prompt an exhaustive search for other causes. A deeper understanding of the mechanisms underlying these complications may help practitioners anticipate the potential risks of new TKIs under development and may guide monitoring and surveillance strategies in the future.

**Table 1: Main characteristics of Bcr-Abl TKI.**

	1 <sup>st</sup> generation	2 <sup>nd</sup> generation			3 <sup>rd</sup> generation
	imatinib	dasatinib	nilotinib	bosutinib	ponatinib
<b>Year of EMA</b>	2001	2006	2007	2013	2013
<b>Type of Bcr-Abl inhibition</b>	II	I	II	I	II
<b>Potency compared with imatinib</b>	–	300x	30x	45x	500x
<b>Indications in CML treatment (chronic phase)</b>					
<b>1<sup>st</sup> line therapy</b>					
	yes	yes*	yes	no	no
<b>2<sup>nd</sup> line therapy</b>					
<b>Intolerance to 1<sup>st</sup> line TKI</b>	yes	yes	yes	yes	no
<b>Failure of 1<sup>st</sup> line imatinib</b>	-	yes	yes	yes	yes
<b>Failure of 1<sup>st</sup> line nilotinib or dasatinib</b>	no	yes	yes	yes	yes
<b>3<sup>rd</sup> line therapy</b>					
<b>Failure and/or intolerance to 2<sup>nd</sup> line TKI</b>		any of the remaining tyrosine kinase inhibitor			
<b>Any line, T315I mutation</b>					
	no	no	no	no	yes
<b>Dose</b>	400-800 mg /day	100 mg /day	300 mg-400 mg twice a day (without food)	500 mg / day (with food)	45 mg / day
<b>Main adverse events</b>	Hepatobiliary disorders, Diarrhea, skin disorders, pleural effusion	Pleural effusion, PAH, Haemorrhage, Hepatobiliary disorders,	skin disorders, hypertension, ischemic heart disease, ischemic cerebrovascular events, Hepatobiliary disorders, metabolism and nutrition disorders	skin disorders, diarrhea, Hepatobiliary disorders, pleural effusion	Hypertension, ischemic heart disease, ischemic cerebrovascular events, venous thrombosis, skin disorders, diarrhea, Hepatobiliary disorders

*CML: chronic myeloid leukemia, EMA: European marketing authorization, PAH: pulmonary arterial hypertension, TKI: tyrosine kinase inhibitor*

**Table 2: Targets of imatinib, dasatinib, nilotinib, bosutinib and ponatinib**

number of common target	Target	imatinib	dasatinib	nilotinib	bosutinib	ponatinib
all	ABL	blue	blue	blue	blue	blue
4	Kit	blue	blue	blue	white	blue
	PDGFRA	blue	blue	blue	white	white
	PDGFRB	blue	blue	blue	white	white
	DDR1	blue	blue	blue	white	white
	CSF1R	blue	blue	blue	white	white
	LCK	blue	white	white	blue	blue
	EGFR	blue	white	white	blue	blue
3	RET	blue	blue	white	white	blue
	FYN	white	blue	white	blue	white
	SRC	white	blue	white	blue	blue
	YES1	white	blue	white	blue	blue
	LYN	white	blue	white	blue	blue
	HCK	white	blue	white	blue	blue
	NTRK1	blue	white	white	blue	blue
2	CA1,2,9,12	blue	white	blue	white	white
	AKT1	white	white	white	white	blue
	CSK	white	blue	white	blue	white
	TNK2	white	blue	white	blue	white
	BTK	white	blue	white	blue	white
	SYK	white	blue	white	blue	white
	FLT3	white	blue	white	white	blue
	EPHA2	white	blue	white	white	blue
	EPHA5	white	blue	white	white	blue
	EPHB1	white	blue	white	white	blue
	FGFR1,2,	white	blue	white	white	blue
	EPHB4	white	blue	white	white	blue
	PTK2B	white	white	white	blue	blue
none	ACVR1B	white	blue	white	white	white
	ACVRL1	white	blue	white	white	white
	ACVR1	white	blue	white	white	white
	TUBA1A	white	blue	white	white	white
	MAPK14	white	blue	white	white	white
	FGFR 3,4	white	white	white	white	blue
	VEGFR1,2,3	white	white	white	white	white

Table 2 describes known targets of imatinib, dasatinib, nilotinib, bosutinib and ponatinib (blue boxes). Number of targets ranges from 7 (nilotinib) to 29 (dasatinib). Twenty-eight targets are common to at least 2 TKI . Among those, Tyrosine-protein kinase ABL is systematically reported explaining their benefit in CML. Targets were extracted from Supertarget Database

(<http://insilico.charite.de/supertarget/index.php>) and article from Greuber et al., Nature Rev Cancer 2013 [13]. Each blue box signifies inhibition by the TKI in the corresponding column.

## FIGURE LEGENDS

**Figure 1:** Mechanisms of dasatinib-induced pleural effusion and pulmonary arterial hypertension. PA-SMC: pulmonary artery smooth muscle cell

**Figure 2:** Various chest computed tomography scan patterns of dasatinib-related pleuro-parenchymal complications. (Top left) ground-glass opacities in the lower right lobe; (Top right) alveolar consolidation in the middle lobe associated with bilateral pleural effusion; (Bottom left) alveolar consolidation in the lower left lobe associated with bilateral pleural effusion; (Bottom right) right-sided pleural effusion.

**Figure 3: Changes in haemodynamic variables and 6-min walk distance (6MWD) by treatment strategy.** Changes in haemodynamics (n=19) from baseline to last evaluation according to initial treatment strategy after the discontinuation of dasatinib: no treatment (n=8), treatment group that received pulmonary arterial hypertension (PAH)-specific treatment (n=9) and calcium channel blocker therapy (n=2). Median (range) values for a) mean pulmonary artery pressure (mPAP) and b) pulmonary vascular resistance (PVR) were not significantly different between treatment groups at baseline or last evaluation, although baseline PVR was higher in the group who received PAH-specific treatment compared with those who received no treatment (8 versus 5.8 Wood units (WU); p=0.14). c) Baseline median cardiac index was significantly lower in the treatment group. \*: p<0.01. d) Baseline median 6MWD was nonsignificantly lower in the treatment group (p=0.15). Reproduced with permission of the © ERS 2019. European Respiratory Journal 50 (1) 1700217; DOI: 10.1183/13993003.00217-2017 Published 27 July 2017

**Figure 4: Diagnosis and management of suspected PAH related to tyrosine kinase inhibitor therapy.**

PH: pulmonary hypertension, TRV: tricuspid regurgitation velocity, RHC: right heart catheterization, PAH: pulmonary arterial hypertension, mPAP: mean pulmonary arterial

pressure, PVR: pulmonary vascular resistance, TKI – tyrosine kinase inhibitor. \$ other causes of pre-capillary PH include obstructive lung diseases, restrictive lung diseases, chronic hypoxemia, and acute or chronic thromboembolic disease. \*A multidisciplinary discussion is recommended to decide which Bcr-Abl tyrosine kinase inhibitor should replace the suspected causative drug. # Risk assessment should be performed using a multidimensional tool such as the REVEAL score or the ESC/ERS 2015 Guidelines Risk assessment tool (Table 13 in reference[103]). ¶ More frequent follow-up evaluations or continuation of PAH specific treatment may be warranted if bosutinib or ponatinib are initiated due to the potential risk of recurrent/worsening PAH with these tyrosine kinase inhibitors.

## **AUTHOR CONTRIBUTIONS**

Literature Search: Jason Weatherald, Louise Bondeelle, Marie-Camille Chaumais, Christophe Guignabert, Philippe Rousselot, Marc Humbert Anne Bergeron, David Montani

Figures & Tables: Jason Weatherald, Louise Bondeelle, Christophe Guignabert, Marie-Camille Chaumais

Writing: All authors contributed to writing the manuscript and critical revisions.

## STATEMENT OF INTERESTS

Dr. Weatherald reports grants, personal fees and non-financial support from Janssen Inc., grants, personal fees and non-financial support from Actelion, personal fees and non-financial support from Bayer, personal fees from Novartis, grants from Alberta Lung Association, grants from Canadian Vascular Network, grants from European Respiratory Society, grants from Canadian Thoracic Society, outside the submitted work. Dr. Bondeelle has nothing to disclose.

Dr. Chaumais reports personal fees from Actelion, non-financial support from Bayer, non-financial support from Boehringer Ingelheim, outside the submitted work. Dr. Guignabert has nothing to disclose. Dr. Savale reports personal fees from Actelion, grants and personal fees from Bayer, grants and personal fees from GSK, outside the submitted work. Dr. Jaïs reports personal fees from Actelion, personal fees from MSD, grants and personal fees from Bayer, grants and personal fees from GSK, outside the submitted work. Dr. Sitbon reports grants, personal fees and non-financial support from Actelion Pharmaceuticals, personal fees and non-financial support from Acceleron Pharmaceuticals, grants, personal fees and non-financial support from Bayer HealthCare, personal fees from Ferrer, grants from GlaxoSmithKline, personal fees from Gossamer Bio, grants, personal fees and non-financial support from MSD, personal fees from United Therapeutics, outside the submitted work. Dr. Rousselot reports grants and personal fees from Pfizer, grants and personal fees from Incyte, grants from Britol Myers Squibb, outside the submitted work. Dr. Humbert reports personal fees from Acceleron, grants and personal fees from Actelion, grants and personal fees from Bayer, grants and personal fees from GSK, personal fees from Merck, personal fees from Morphogen IX, personal fees from United Therapeutics, outside the submitted work. Dr. Bergeron reports grants from



SOS oxygene, personal fees from Shire, personal fees from Pfizer, personal fees from MSD, personal fees from Gilead, outside the submitted work. Dr. Montani reports grants and personal fees from Actelion, grants and personal fees from Bayer, personal fees from GSK, personal fees from Pfizer, personal fees from MSD, personal fees from Chiesi, outside the submitted work

## REFERENCES

1. Goldman JM, Melo JV. Chronic myeloid leukemia--advances in biology and new approaches to treatment. *N. Engl. J. Med.* 2003; 349: 1451–1464.
2. Rosti G, Castagnetti F, Gugliotta G, Baccarani M. Tyrosine kinase inhibitors in chronic myeloid leukaemia: which, when, for whom? *Nat Rev Clin Oncol* 2017; 14: 141–154.
3. Bower H, Björkholm M, Dickman PW, Höglund M, Lambert PC, Andersson TM-L. Life Expectancy of Patients With Chronic Myeloid Leukemia Approaches the Life Expectancy of the General Population. *J. Clin. Oncol.* 2016; 34: 2851–2857.
4. Baccarani M, Deininger MW, Rosti G, Hochhaus A, Soverini S, Apperley JF, Cervantes F, Clark RE, Cortes JE, Guilhot F, Hjorth-Hansen H, Hughes TP, Kantarjian HM, Kim D-W, Larson RA, Lipton JH, Mahon F-X, Martinelli G, Mayer J, Müller MC, Niederwieser D, Pane F, Radich JP, Rousselot P, Saglio G, Sauße S, Schiffer C, Silver R, Simonsson B, Steegmann J-L, et al. European LeukemiaNet recommendations for the management of chronic myeloid leukemia: 2013. *Blood* 2013; 122: 872–884.
5. Cortes JE, Saglio G, Kantarjian HM, Baccarani M, Mayer J, Boqué C, Shah NP, Chuah C, Casanova L, Bradley-Garelik B, Manos G, Hochhaus A. Final 5-Year Study Results of DASISION: The Dasatinib Versus Imatinib Study in Treatment-Naïve Chronic Myeloid Leukemia Patients Trial. *J. Clin. Oncol.* 2016; 34: 2333–2340.
6. Giles FJ, Rosti G, Beris P, Clark RE, le Coutre P, Mahon F-X, Steegmann J-L, Valent P, Saglio G. Nilotinib is superior to imatinib as first-line therapy of chronic myeloid leukemia: the ENESTnd study. *Expert Rev Hematol* 2010; 3: 665–673.
7. Saussele S, Richter J, Guilhot J, Gruber FX, Hjorth-Hansen H, Almeida A, Janssen JJWM, Mayer J, Koskenvesa P, Panayiotidis P, Olsson-Strömberg U, Martinez-Lopez J, Rousselot P, Vestergaard H, Ehrencrona H, Kairisto V, Machová Poláková K, Müller MC, Mustjoki S, Berger MG, Fabarius A, Hofmann W-K, Hochhaus A, Pfirrmann M, Mahon F-X, EURO-SKI investigators. Discontinuation of tyrosine kinase inhibitor therapy in chronic myeloid leukaemia (EURO-SKI): a prespecified interim analysis of a prospective, multicentre, non-randomised, trial. *Lancet Oncol.* 2018; 19: 747–757.
8. Nagar B, Bornmann WG, Pellicena P, Schindler T, Veach DR, Miller WT, Clarkson B, Kuriyan J. Crystal structures of the kinase domain of c-Abl in complex with the small molecule inhibitors PD173955 and imatinib (STI-571). *Cancer Res.* 2002; 62: 4236–4243.
9. Modugno M. New resistance mechanisms for small molecule kinase inhibitors of Abl kinase. *Drug Discov Today Technol* 2014; 11: 5–10.

10. Redaelli S, Piazza R, Rostagno R, Magistroni V, Perini P, Marega M, Gambacorti-Passerini C, Boschelli F. Activity of bosutinib, dasatinib, and nilotinib against 18 imatinib-resistant BCR/ABL mutants. *J. Clin. Oncol.* 2009; 27: 469–471.
11. Cortes JE, Kantarjian H, Shah NP, Bixby D, Mauro MJ, Flinn I, O'Hare T, Hu S, Narasimhan NI, Rivera VM, Clackson T, Turner CD, Haluska FG, Druker BJ, Deininger MWN, Talpaz M. Ponatinib in refractory Philadelphia chromosome-positive leukemias. *N. Engl. J. Med.* 2012; 367: 2075–2088.
12. Hochhaus A, Saussele S, Rosti G, Mahon F-X, Janssen JJWM, Hjorth-Hansen H, Richter J, Buske C, ESMO Guidelines Committee. Chronic myeloid leukaemia: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* 2017; 28: iv41–iv51.
13. Greuber EK, Smith-Pearson P, Wang J, Pendergast AM. Role of ABL family kinases in cancer: from leukaemia to solid tumours. *Nat. Rev. Cancer* 2013; 13: 559–571.
14. Duong-Ly KC, Devarajan K, Liang S, Horiuchi KY, Wang Y, Ma H, Peterson JR. Kinase Inhibitor Profiling Reveals Unexpected Opportunities to Inhibit Disease-Associated Mutant Kinases. *Cell Reports* 2016; 14: 772–781.
15. Suh CH, Park HS, Kim KW, Pyo J, Hatabu H, Nishino M. Pneumonitis in advanced non-small-cell lung cancer patients treated with EGFR tyrosine kinase inhibitor: Meta-analysis of 153 cohorts with 15,713 patients: Meta-analysis of incidence and risk factors of EGFR-TKI pneumonitis in NSCLC. *Lung Cancer* 2018; 123: 60–69.
16. Ding PN, Lord SJ, GebSKI V, Links M, Bray V, Gralla RJ, Yang JC-H, Lee CK. Risk of Treatment-Related Toxicities from EGFR Tyrosine Kinase Inhibitors: A Meta-analysis of Clinical Trials of Gefitinib, Erlotinib, and Afatinib in Advanced EGFR-Mutated Non-Small Cell Lung Cancer. *J Thorac Oncol* 2017; 12: 633–643.
17. Montani D, Bergot E, Günther S, Savale L, Bergeron A, Bourdin A, Bouvaist H, Canuet M, Pison C, Macro M, Poubeau P, Girerd B, Natali D, Guignabert C, Perros F, O'Callaghan DS, Jaïs X, Tubert-Bitter P, Zalcman G, Sitbon O, Simonneau G, Humbert M. Pulmonary arterial hypertension in patients treated by dasatinib. *Circulation* 2012; 125: 2128–2137.
18. Riou M, Seferian A, Savale L, Chaumais M-C, Guignabert C, Canuet M, Magro P, Rea D, Sitbon O, Jaïs X, Humbert M, Montani D. Deterioration of pulmonary hypertension and pleural effusion with bosutinib following dasatinib lung toxicity. *Eur. Respir. J.* 2016; 48: 1517–1519.
19. Weatherald J, Chaumais M-C, Savale L, Jaïs X, Seferian A, Canuet M, Bouvaist H, Magro P, Bergeron A, Guignabert C, Sitbon O, Simonneau G, Humbert M, Montani D. Long-term outcomes of dasatinib-induced pulmonary arterial hypertension: a population-based study. *Eur. Respir. J.* 2017; 50: 1700217.

20. Shah NP, Rousselot P, Schiffer C, Rea D, Cortes JE, Milone J, Mohamed H, Healey D, Kantarjian H, Hochhaus A, Saglio G. Dasatinib in imatinib-resistant or -intolerant chronic-phase, chronic myeloid leukemia patients: 7-year follow-up of study CA180-034. *Am. J. Hematol.* 2016; 91: 869–874.
21. Guignabert C, Phan C, Seferian A, Huertas A, Tu L, Thuillet R, Sattler C, Le Hiress M, Tamura Y, Jutant E-M, Chaumais M-C, Bouchet S, Manéglier B, Molimard M, Rousselot P, Sitbon O, Simonneau G, Montani D, Humbert M. Dasatinib induces lung vascular toxicity and predisposes to pulmonary hypertension. *J. Clin. Invest.* 2016; 126: 3207–3218.
22. Phan C, Jutant E-M, Tu L, Thuillet R, Seferian A, Montani D, Huertas A, Bezu J van, Breijer F, Vonk Noordegraaf A, Humbert M, Aman J, Guignabert C. Dasatinib increases endothelial permeability leading to pleural effusion. *Eur. Respir. J.* 2018; 51: 1701096.
23. Aghel N, Delgado DH, Lipton JH. Cardiovascular toxicities of BCR-ABL tyrosine kinase inhibitors in chronic myeloid leukemia: preventive strategies and cardiovascular surveillance. *Vasc Health Risk Manag* 2017; 13: 293–303.
24. Moslehi JJ, Deininger M. Tyrosine Kinase Inhibitor-Associated Cardiovascular Toxicity in Chronic Myeloid Leukemia. *J. Clin. Oncol.* 2015; 33: 4210–4218.
25. Huertas A, Guignabert C, Barberà JA, Bärtsch P, Bhattacharya J, Bhattacharya S, Bonsignore MR, Dewachter L, Dinh-Xuan AT, Dorfmueller P, Gladwin MT, Humbert M, Kotsimbos T, Vassilakopoulos T, Sanchez O, Savale L, Testa U, Wilkins MR. Pulmonary vascular endothelium: the orchestra conductor in respiratory diseases: Highlights from basic research to therapy. *Eur. Respir. J.* 2018; 51: 1700745.
26. Fazakas C, Nagaraj C, Zabini D, Végh AG, Marsh LM, Wilhelm I, Krizbai IA, Olschewski H, Olschewski A, Bálint Z. Rho-Kinase Inhibition Ameliorates Dasatinib-Induced Endothelial Dysfunction and Pulmonary Hypertension. *Front Physiol* 2018; 9: 537.
27. Roos WP, Kaina B. DNA damage-induced cell death: from specific DNA lesions to the DNA damage response and apoptosis. *Cancer Lett.* 2013; 332: 237–248.
28. Daccord C, Letovanec I, Yerly P, Bloch J, Ognà A, Nicod LP, Aubert J-D. First histopathological evidence of irreversible pulmonary vascular disease in dasatinib-induced pulmonary arterial hypertension. *Eur. Respir. J.* 2018; 51: 1701694.
29. Nagaraj C, Tang B, Bálint Z, Wygrecka M, Hrzenjak A, Kwapiszewska G, Stacher E, Lindenmann J, Weir EK, Olschewski H, Olschewski A. Src tyrosine kinase is crucial for potassium channel function in human pulmonary arteries. *Eur. Respir. J.* 2013; 41: 85–95.
30. Di Salvo J, Steusloff A, Semenchuk L, Satoh S, Kolquist K, Pfitzer G. Tyrosine kinase inhibitors suppress agonist-induced contraction in smooth muscle. *Biochem. Biophys. Res. Commun.* 1993; 190: 968–974.

31. Rieg AD, Bunting NA, Cranen C, Suleiman S, Spillner JW, Schnöring H, Schröder T, von Stillfried S, Braunschweig T, Manley PW, Schälte G, Rossaint R, Uhlig S, Martin C. Tyrosine kinase inhibitors relax pulmonary arteries in human and murine precision-cut lung slices. *Respir. Res.* 2019; 20: 111.
32. Maihöfer NA, Suleiman S, Dreytmüller D, Manley PW, Rossaint R, Uhlig S, Martin C, Rieg AD. Imatinib relaxes the pulmonary venous bed of guinea pigs. *Respir. Res.* 2017; 18: 32.
33. Abe K, Toba M, Alzoubi A, Koubsky K, Ito M, Ota H, Gairhe S, Gerthoffer WT, Fagan KA, McMurtry IF, Oka M. Tyrosine kinase inhibitors are potent acute pulmonary vasodilators in rats. *Am. J. Respir. Cell Mol. Biol.* 2011; 45: 804–808.
34. Pankey EA, Thammasiboon S, Lasker GF, Baber S, Lasky JA, Kadowitz PJ. Imatinib attenuates monocrotaline pulmonary hypertension and has potent vasodilator activity in pulmonary and systemic vascular beds in the rat. *Am. J. Physiol. Heart Circ. Physiol.* 2013; 305: H1288-1296.
35. Cornet L, Khouri C, Roustit M, Guignabert C, Chaumais MC, Humbert M, Revol B, Despas F, Montani D, Cracowski JL. Pulmonary Arterial Hypertension associated with Protein Kinase Inhibitors: A pharmacovigilance-pharmacodynamic study. *Eur. Respir. J.* 2019; 53: 1802472.
36. Davis RE, Ngo VN, Lenz G, Tolar P, Young RM, Romesser PB, Kohlhammer H, Lamy L, Zhao H, Yang Y, Xu W, Shaffer AL, Wright G, Xiao W, Powell J, Jiang J-K, Thomas CJ, Rosenwald A, Ott G, Muller-Hermelink HK, Gascoyne RD, Connors JM, Johnson NA, Rimsza LM, Campo E, Jaffe ES, Wilson WH, Delabie J, Smeland EB, Fisher RI, et al. Chronic active B-cell-receptor signalling in diffuse large B-cell lymphoma. *Nature* 2010; 463: 88–92.
37. Darvishi B, Farahmand L, Eslami-S Z, Majidzadeh-A K. NF- $\kappa$ B as the main node of resistance to receptor tyrosine kinase inhibitors in triple-negative breast cancer. *Tumour Biol.* 2017; 39: 1010428317706919.
38. Liu C, Mu X, Wang X, Zhang C, Zhang L, Yu B, Sun G. Ponatinib Inhibits Proliferation and Induces Apoptosis of Liver Cancer Cells, but Its Efficacy Is Compromised by Its Activation on PDK1/Akt/mTOR Signaling. *Molecules* 2019; 24: E1363.
39. Sajithlal GB, Hamed HA, Cruickshanks N, Booth L, Tavallai S, Syed J, Grant S, Poklepovic A, Dent P. Sorafenib/regorafenib and phosphatidylinositol 3 kinase/thymoma viral proto-oncogene inhibition interact to kill tumor cells. *Mol. Pharmacol.* 2013; 84: 562–571.
40. Larson RA, Druker BJ, Guilhot F, O'Brien SG, Riviere GJ, Krahnke T, Gathmann I, Wang Y, IRIS (International Randomized Interferon vs STI571) Study Group. Imatinib pharmacokinetics and its correlation with response and safety in chronic-phase chronic myeloid leukemia: a subanalysis of the IRIS study. *Blood* 2008; 111: 4022–4028.

41. Picard S, Titier K, Etienne G, Teilhet E, Ducint D, Bernard M-A, Lassalle R, Marit G, Reiffers J, Begaud B, Moore N, Molimard M, Mahon F-X. Trough imatinib plasma levels are associated with both cytogenetic and molecular responses to standard-dose imatinib in chronic myeloid leukemia. *Blood* 2007; 109: 3496–3499.
42. Larson RA, Yin OQP, Hochhaus A, Saglio G, Clark RE, Nakamae H, Gallagher NJ, Demirhan E, Hughes TP, Kantarjian HM, le Coutre PD. Population pharmacokinetic and exposure-response analysis of nilotinib in patients with newly diagnosed Ph+ chronic myeloid leukemia in chronic phase. *Eur. J. Clin. Pharmacol.* 2012; 68: 723–733.
43. Wang X, Roy A, Hochhaus A, Kantarjian HM, Chen T-T, Shah NP. Differential effects of dosing regimen on the safety and efficacy of dasatinib: retrospective exposure-response analysis of a Phase III study. *Clin Pharmacol* 2013; 5: 85–97.
44. Mita A, Abumiya M, Miura M, Niioka T, Takahashi S, Yoshioka T, Kameoka Y, Takahashi N. Correlation of plasma concentration and adverse effects of bosutinib: standard dose or dose-escalation regimens of bosutinib treatment for patients with chronic myeloid leukemia. *Exp Hematol Oncol* 2018; 7: 9.
45. Shah NP, Kim D-W, Kantarjian H, Rousselot P, Llacer PED, Enrico A, Vela-Ojeda J, Silver RT, Khoury HJ, Müller MC, Lambert A, Matloub Y, Hochhaus A. Potent, transient inhibition of BCR-ABL with dasatinib 100 mg daily achieves rapid and durable cytogenetic responses and high transformation-free survival rates in chronic phase chronic myeloid leukemia patients with resistance, suboptimal response or intolerance to imatinib. *Haematologica* 2010; 95: 232–240.
46. Mahé J, de Campaigno EP, Chené A-L, Montastruc J-L, Despas F, Jolliet P. Pleural adverse drugs reactions and protein kinase inhibitors: Identification of suspicious targets by disproportionality analysis from VigiBase. *Br J Clin Pharmacol* 2018; 84: 2373–2383.
47. Han J, Zhang G, Welch EJ, Liang Y, Fu J, Vogel SM, Lowell CA, Du X, Cheresch DA, Malik AB, Li Z. A critical role for Lyn kinase in strengthening endothelial integrity and barrier function. *Blood* 2013; 122: 4140–4149.
48. Talpaz M, Shah NP, Kantarjian H, Donato N, Nicoll J, Paquette R, Cortes J, O'Brien S, Nicaise C, Bleickardt E, Blackwood-Chirchir MA, Iyer V, Chen T-T, Huang F, Decillis AP, Sawyers CL. Dasatinib in imatinib-resistant Philadelphia chromosome-positive leukemias. *N. Engl. J. Med.* 2006; 354: 2531–2541.
49. Hughes TP, Laneuville P, Rousselot P, Snyder DS, Rea D, Shah NP, Paar D, Abruzzese E, Hochhaus A, Lipton JH, Cortes JE. Incidence, outcomes, and risk factors of pleural effusion in patients receiving dasatinib therapy for Philadelphia chromosome-positive leukemia. *Haematologica* 2019; 104: 93–101.

50. Shah NP, Kantarjian HM, Kim D-W, Réa D, Dorlhiac-Llacer PE, Milone JH, Vela-Ojeda J, Silver RT, Khoury HJ, Charbonnier A, Khoroshko N, Paquette RL, Deininger M, Collins RH, Otero I, Hughes T, Bleickardt E, Strauss L, Francis S, Hochhaus A. Intermittent target inhibition with dasatinib 100 mg once daily preserves efficacy and improves tolerability in imatinib-resistant and -intolerant chronic-phase chronic myeloid leukemia. *J. Clin. Oncol.* 2008; 26: 3204–3212.
51. Naqvi K, Jabbour E, Skinner J, Yilmaz M, Ferrajoli A, Bose P, Thompson P, Alvarado Y, Jain N, Takahashi K, Burger J, Estrov Z, Borthakur G, Pemmaraju N, Paul S, Cortes J, Kantarjian H. Early Results of Lower dose Dasatinib 50mg Daily as Frontline Therapy in Newly Diagnosed Chronic Phase Chronic Myeloid Leukemia. *Cancer* 2018; 124: 2740–2747.
52. Khoury HJ, Guilhot F, Hughes TP, Kim D-W, Cortes JE. Dasatinib treatment for Philadelphia chromosome-positive leukemias: practical considerations. *Cancer* 2009; 115: 1381–1394.
53. de Lavallade H, Punnialingam S, Milojkovic D, Bua M, Khorashad JS, Gabriel IH, Chaidos A, Olavarria E, Goldman JM, Apperley JF, Marin D. Pleural effusions in patients with chronic myeloid leukaemia treated with dasatinib may have an immune-mediated pathogenesis. *Br. J. Haematol.* 2008; 141: 745–747.
54. Bergeron A, Réa D, Levy V, Picard C, Meignin V, Tamburini J, Bruzzoni-Giovanelli H, Calvo F, Tazi A, Rousselot P. Lung abnormalities after dasatinib treatment for chronic myeloid leukemia: a case series. *Am. J. Respir. Crit. Care Med.* 2007; 176: 814–818.
55. Rea D, Bergeron A, Fieschi C, Bengoufa D, Oksenhendler E, Dombret H. Dasatinib-induced lupus. *Lancet* 2008; 372: 713–714.
56. Quintás-Cardama A, Kantarjian H, O’Brien S, Borthakur G, Bruzzi J, Munden R, Cortes J. Pleural effusion in patients with chronic myelogenous leukemia treated with dasatinib after imatinib failure. *J. Clin. Oncol.* 2007; 25: 3908–3914.
57. Ferreiro L, San-José E, Suárez-Antelo J, Valdés L. Dasatinib-induced pleural effusion: Chylothorax, an option to consider. *Ann Thorac Med* 2016; 11: 289–293.
58. Chen B, Wu Z, Wang Q, Li W, Cheng D. Dasatinib-induced chylothorax: report of a case and review of the literature. *Invest New Drugs* 2020; doi: 10.1007/s10637-020-00932-3. [Epub ahead of print].
59. Al-Abcha A, Iftikhar MH, Abu Rous F, Laird-Fick H. Chylothorax: complication attributed to dasatinib use. *BMJ Case Rep* 2019; 12: e231653.
60. Adrianza AM, Jimenez CA, Aldana GL, Pozo L, Cortes JE, Sagar AES. Tyrosine Kinase Inhibitor-associated Fibrothorax. *Am. J. Respir. Crit. Care Med.* 2019; 199: 1432.

61. Sato M, Watanabe S, Aoki N, Asakawa K, Moriyama H, Oshima Y, Koya T, Okazuka K, Kikuchi T. [A Case of Drug-Induced Organizing Pneumonia Caused by Dasatinib]. *Gan To Kagaku Ryoho* 2018; 45: 851–854.
62. Radaelli F, Bramanti S, Fantini NN, Fabio G, Greco I, Lambertenghi-Delilieri G. Dasatinib-related alveolar pneumonia responsive to corticosteroids. *Leuk. Lymphoma* 2006; 47: 1180–1181.
63. Jasielec JK, Larson RA. Dasatinib-Related Pulmonary Toxicity Mimicking an Atypical Infection. *J. Clin. Oncol.* 2016; 34: e46-48.
64. Ito Y, Tanigawa M, Iwamoto K, Nigi A, Itani H, Kondo S, Tokui T, Usui E, Tamaki S. Interstitial pneumonitis associated with dasatinib: two case reports and literature review. *Respir Investig* 2019; 57: 506–509.
65. Druker BJ, Guilhot F, O'Brien SG, Gathmann I, Kantarjian H, Gattermann N, Deininger MWN, Silver RT, Goldman JM, Stone RM, Cervantes F, Hochhaus A, Powell BL, Ghabrilove JL, Rousselot P, Reiffers J, Cornelissen JJ, Hughes T, Agis H, Fischer T, Verhoef G, Shepherd J, Saglio G, Gratwohl A, Nielsen JL, Radich JP, Simonsson B, Taylor K, Baccarani M, So C, et al. Five-year follow-up of patients receiving imatinib for chronic myeloid leukemia. *N. Engl. J. Med.* 2006; 355: 2408–2417.
66. Banka R, Udwadia Z. Imatinib-induced pleural effusion: A case report. *J Postgrad Med* 2017; 63: 55–57.
67. Breccia M, D'Elia GM, D'Andrea M, Latagliata R, Alimena G. Pleural-pericardic effusion as uncommon complication in CML patients treated with Imatinib. *Eur. J. Haematol.* 2005; 74: 89–90.
68. Bergeron A, Bergot E, Vilela G, Ades L, Devergie A, Espérou H, Socié G, Calvo F, Gluckman E, Ribaud P, Rousselot P, Tazi A. Hypersensitivity pneumonitis related to imatinib mesylate. *J. Clin. Oncol.* 2002; 20: 4271–4272.
69. Rajda J, Phatak PD. Reversible drug-induced interstitial pneumonitis following imatinib mesylate therapy. *Am. J. Hematol.* 2005; 79: 80–81.
70. Isshiki I, Yamaguchi K, Okamoto S. Interstitial pneumonitis during imatinib therapy. *Br. J. Haematol.* 2004; 125: 420.
71. Zhang P, Huang J, Jin F, Pan J, Ouyang G. Imatinib-induced irreversible interstitial lung disease: A case report. *Medicine (Baltimore)* 2019; 98: e14402.
72. Ohnishi K, Sakai F, Kudoh S, Ohno R. Twenty-seven cases of drug-induced interstitial lung disease associated with imatinib mesylate. *Leukemia* 2006; 20: 1162–1164.




73. Yoshimura M, Kojima K, Tomimasu R, Fukushima N, Hayashi S, Sueoka E, Kimura S. ABL tyrosine kinase inhibitor-induced pulmonary alveolar proteinosis in chronic myeloid leukemia. *Int. J. Hematol.* 2014; 100: 611–614.
74. Suh KJ, Lee JY, Shin D-Y, Koh Y, Bang S-M, Yoon S-S, Park S, Kim I, Lee J-O. Analysis of adverse events associated with dasatinib and nilotinib treatments in chronic-phase chronic myeloid leukemia patients outside clinical trials. *Int. J. Hematol.* 2017; 106: 229–239.
75. Go S-I, Lee WS, Lee G-W, Kang JH, Kang MH, Lee J-H, Kim H-G. Nilotinib-induced interstitial lung disease. *Int. J. Hematol.* 2013; 98: 361–365.
76. Usküdar Teke H, Akay OM, Gören Şahin D, Karagülle M, Gündüz E, Andıç N. Pleural effusion: a rare side effect of nilotinib—a case report. *Case Rep Med* 2014; 2014: 203939.
77. Gambacorti-Passerini C, Cortes JE, Lipton JH, Kantarjian HM, Kim D-W, Schafhausen P, Crescenzo R, Bardy-Bouxin N, Shapiro M, Noonan K, Leip E, DeAnnuntis L, Brümmendorf TH, Khoury HJ. Safety and efficacy of second-line bosutinib for chronic phase chronic myeloid leukemia over a five-year period: final results of a phase I/II study. *Haematologica* 2018; 103: 1298–1307.
78. Gambacorti-Passerini C, Brümmendorf TH, Kim D-W, Turkina AG, Masszi T, Assouline S, Durrant S, Kantarjian HM, Khoury HJ, Zaritskey A, Shen Z-X, Jin J, Vellenga E, Pasquini R, Mathews V, Cervantes F, Besson N, Turnbull K, Leip E, Kelly V, Cortes JE. Bosutinib efficacy and safety in chronic phase chronic myeloid leukemia after imatinib resistance or intolerance: Minimum 24-month follow-up. *Am. J. Hematol.* 2014; 89: 732–742.
79. Brümmendorf TH, Cortes JE, Khoury HJ, Kantarjian HM, Kim D-W, Schafhausen P, Conlan MG, Shapiro M, Turnbull K, Leip E, Gambacorti-Passerini C, Lipton JH. Factors influencing long-term efficacy and tolerability of bosutinib in chronic phase chronic myeloid leukaemia resistant or intolerant to imatinib. *Br. J. Haematol.* 2016; 172: 97–110.
80. Jutant E-M, Meignin V, Montani D, Tazi A, Rousselot P, Bergeron A. Bosutinib-related pneumonitis. *Eur. Respir. J.* 2017; 50: 1700930.
81. Iurlo A, Galimberti S, Abruzzese E, Annunziata M, Bonifacio M, Latagliata R, Pregno P, Ferrero D, Sorà F, Orlandi EM, Fava C, Cattaneo D, Bucelli C, Binotto G, Pungolino E, Tiribelli M, Gozzini A, Gugliotta G, Castagnetti F, Stagno F, Rege-Cambrin G, Martino B, Luciano L, Breccia M, Sica S, Bocchia M, Pane F, Saglio G, Rosti G, Specchia G, et al. Pleural effusion and molecular response in dasatinib-treated chronic myeloid leukemia patients in a real-life Italian multicenter series. *Ann. Hematol.* 2018; 97: 95–100.
82. Aoyama R, Ishikawa J, Harada K. The efficacy of tolvaptan in treating dasatinib-induced pleural effusions in patients with chronic myelogenous leukemia. *J Cardiol* 2020; 75: 203–207.




83. Miura M. Therapeutic drug monitoring of imatinib, nilotinib, and dasatinib for patients with chronic myeloid leukemia. *Biol. Pharm. Bull.* 2015; 38: 645–654.
84. García-Gutiérrez V, Milojkovic D, Hernandez-Boluda JC, Claudiani S, Martin Mateos ML, Casado-Montero LF, González G, Jimenez-Velasco A, Boque C, Martinez-Trillos A, Vázquez IM, Payer ÁR, Senín A, Amustio Díez E, García AB, Carrascosa GB, Ortí G, Ruiz BC, Fernández MÁ, Del Carmen García Garay M, Giraldo P, Guinea JM, De Las Heras Rodríguez N, Hernán N, Pérez AI, Piris-Villaespesa M, Lorenzo JLL, Martí-Tutusaus JMM, Vallansot RO, Ortega Rivas F, et al. Safety and efficacy of bosutinib in fourth-line therapy of chronic myeloid leukemia patients. *Ann. Hematol.* 2019; 98: 321–330.
85. Tiribelli M, Abruzzese E, Capodanno I, Sorà F, Trabacchi E, Iurlo A, Luciano L, Binotto G, Bonifacio M, Annunziata M, Crugnola M, Fanin R. Efficacy and safety of bosutinib in chronic phase CML patients developing pleural effusion under dasatinib therapy. *Ann. Hematol.* 2019; 98: 2609–2611.
86. Dao K, Védy D, Lopez J, Staneczek O, Buclin T, Livio F. Imatinib-induced dose-dependent interstitial lung disease successfully switched to nilotinib: a case report with concentration exposure data. *Int. J. Hematol.* 2013; 97: 299–300.
87. Delomas T, Darné C, Besson C. Lack of recurrence of imatinib-induced interstitial lung disease with nilotinib. *Leuk. Lymphoma* 2012; 53: 332–333.
88. Simonneau G, Montani D, Celermajer DS, Denton CP, Gatzoulis MA, Krowka M, Williams PG, Souza R. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur. Respir. J.* 2019; 53: 1801913.
89. Rasheed W, Flaim B, Seymour JF. Reversible severe pulmonary hypertension secondary to dasatinib in a patient with chronic myeloid leukemia. *Leuk. Res.* 2009; 33: 861–864.
90. Mattei D, Feola M, Orzan F, Mordini N, Rapezzi D, Gallamini A. Reversible dasatinib-induced pulmonary arterial hypertension and right ventricle failure in a previously allografted CML patient. *Bone Marrow Transplant.* 2009; 43: 967–968.
91. Fox LC, Cummins KD, Costello B, Yeung D, Cleary R, Forsyth C, Tatarczuch M, Burbury K, Motorna O, Shortt J, Fleming S, McQuillan A, Schwarzer A, Harrup R, Holmes A, Ratnasingam S, Chan K-L, Hsu W-H, Ashraf A, Putt F, Grigg A. The incidence and natural history of dasatinib complications in the treatment of chronic myeloid leukemia. *Blood Adv* 2017; 1: 802–811.
92. Hickey PM, Thompson AAR, Charalampopoulos A, Elliot CA, Hamilton N, Kiely DG, Lawrie A, Sabroe I, Condliffe R. Bosutinib therapy resulting in severe deterioration of pre-existing pulmonary arterial hypertension. *Eur. Respir. J.* 2016; 48: 1514–1516.

93. Seegobin K, Babbar A, Ferreira J, Lyons B, Cury J, Seeram V. A case of worsening pulmonary arterial hypertension and pleural effusions by bosutinib after prior treatment with dasatinib. *Pulm Circ* 2017; 7: 808–812.
94. Quilot F-M, Georges M, Favrolt N, Beltramo G, Foignot C, Grandvilllemin A, Montani D, Bonniaud P, Camus P. Pulmonary hypertension associated with ponatinib therapy. *Eur. Respir. J.* 2016; 47: 676–679.
95. Zakrzewski D, Seferynska I, Warzocha K, Hryniewiecki T. Elevation of pulmonary artery pressure as a complication of nilotinib therapy for chronic myeloid leukemia. *Int J Hematol* 2012; 96: 132–135.
96. Ghofrani HA, Morrell NW, Hoeper MM, Olschewski H, Peacock AJ, Barst RJ, Shapiro S, Golpon H, Toshner M, Grimminger F, Pascoe S. Imatinib in pulmonary arterial hypertension patients with inadequate response to established therapy. *Am. J. Respir. Crit. Care Med.* 2010; 182: 1171–1177.
97. Hoeper MM, Barst RJ, Bourge RC, Feldman J, Frost AE, Galié N, Gómez-Sánchez MA, Grimminger F, Grünig E, Hassoun PM, Morrell NW, Peacock AJ, Satoh T, Simonneau G, Tapson VF, Torres F, Lawrence D, Quinn DA, Ghofrani H-A. Imatinib mesylate as add-on therapy for pulmonary arterial hypertension: results of the randomized IMPRES study. *Circulation* 2013; 127: 1128–1138.
98. Shah NP, Guilhot F, Cortes JE, Schiffer CA, le Coutre P, Brümmendorf TH, Kantarjian HM, Hochhaus A, Rousselot P, Mohamed H, Healey D, Cunningham M, Saglio G. Long-term outcome with dasatinib after imatinib failure in chronic-phase chronic myeloid leukemia: follow-up of a phase 3 study. *Blood* 2014; 123: 2317–2324.
99. Toya T, Nagatomo Y, Kagami K, Adachi T. Dasatinib-induced pulmonary arterial hypertension complicated with scleroderma: a case report. *Eur Heart J Case Rep* 2019; 3: ytz025.
100. Nihtyanova SI, Schreiber BE, Ong VH, Rosenberg D, Moinzadeh P, Coghlan JG, Wells AU, Denton CP. Prediction of pulmonary complications and long-term survival in systemic sclerosis. *Arthritis & Rheumatol* 2014; 66: 1625–1635.
101. Orlikow E, Weatherald J, Hirani N. Dasatinib-induced pulmonary arterial hypertension. *Can J Cardiol* 2019; 35: 1604.e1-1604.e3
102. Shah NP, Wallis N, Farber HW, Mauro MJ, Wolf RA, Mattei D, Guha M, Rea D, Peacock A. Clinical features of pulmonary arterial hypertension in patients receiving dasatinib. *Am. J. Hematol.* 2015; 90: 1060–1064.
103. Morishita S, Hagihara M, Itabashi M, Ishii Y, Yamamoto W, Numata A, Motohashi K, Matsumoto K, Fujisawa S, Nakajima H. Development of pulmonary arterial hypertension


during oral dasatinib therapy for chronic myelogenous leukemia. *Rinsho Ketsueki* 2016; 57: 999–1003.

104. Galiè N, Humbert M, Vachiery J-L, Gibbs S, Lang I, Torbicki A, Simonneau G, Peacock A, Vonk Noordegraaf A, Beghetti M, Ghofrani A, Gomez Sanchez MA, Hansmann G, Klepetko W, Lancellotti P, Matucci M, McDonagh T, Pierard LA, Trindade PT, Zompatori M, Hoeper M. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension: The Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). *Eur. Respir. J.* 2015; 46: 903–975.
105. Cortes JE, Kim D-W, Pinilla-Ibarz J, le Coutre PD, Paquette R, Chuah C, Nicolini FE, Apperley JF, Khoury HJ, Talpaz M, DeAngelo DJ, Abruzzese E, Rea D, Baccarani M, Müller MC, Gambacorti-Passerini C, Lustgarten S, Rivera VM, Haluska FG, Guilhot F, Deininger MW, Hochhaus A, Hughes TP, Shah NP, Kantarjian HM. Ponatinib efficacy and safety in Philadelphia chromosome-positive leukemia: final 5-year results of the phase 2 PACE trial. *Blood* 2018; 132: 393–404.
106. Galiè N, Channick RN, Frantz RP, Grünig E, Jing ZC, Moiseeva O, Preston IR, Pulido T, Safdar Z, Tamura Y, McLaughlin VV. Risk stratification and medical therapy of pulmonary arterial hypertension. *Eur. Respir. J.* 2019; 53: 1801889.
107. Jose A, Rafei H, Ahari J. Combination targeted pulmonary hypertension therapy in the resolution of Dasatinib-associated pulmonary arterial hypertension. *Pulm Circ* 2017; 7: 803–807.
108. Nishimori M, Honjo T, Kaihotsu K, Sone N, Yoshikawa S, Imanishi J, Nakayama K, Emoto N, Iwahashi M. Dasatinib-Induced Pulmonary Arterial Hypertension Treated with Upfront Combination Therapy. *Case Rep Cardiol* 2018; 2018: 3895197.
109. Weatherald J, Chaumais M-C, Montani D. Pulmonary arterial hypertension induced by tyrosine kinase inhibitors. *Curr Opin Pulm Med* 2017; 23: 392–397.







**A.**



**B.**



**C.**



**D.**

